


# Building up of a nested granite intrusion: magnetic fabric, gravity modelling and fluid inclusion planes studies in Santa Eulália Plutonic Complex (Ossa Morena Zone, Portugal)

H. SANT'OVAIA<sup>a1</sup>  <http://orcid.org/0000-0002-1884-7440>, P. NOGUEIRA<sup>a2</sup>, J. CARRILHO LOPES<sup>a3</sup>, C. GOMES<sup>a4</sup>, M.A. RIBEIRO<sup>a1</sup>, H.C.B. MARTINS<sup>a1</sup>, A. DÓRIA<sup>a1</sup>, C. CRUZ<sup>a1</sup>, L. LOPES<sup>a5</sup>, R. SARDINHA<sup>a5</sup>, A. ROCHA<sup>a4</sup> and F. NORONHA<sup>a1</sup>

<sup>a1</sup> Geology Centre of Porto University, Department of Geosciences, Environment and Spatial Planning, Faculty of Sciences of Porto University, Rua do Campo Alegre, 4169-007 Porto, Portugal

<sup>a2</sup> Geology Centre of Porto University, Geosciences Department of Évora University, Largo dos Colegiais 2, 7004-516 Évora, Portugal

<sup>a3</sup> Geology Centre of Lisbon University, Geosciences Department of Évora University, Largo dos Colegiais 2, 7004-516 Évora, Portugal

<sup>a4</sup> Geophysics Centre of University of Coimbra, Department of Earth Sciences, Faculty of Sciences and Technology of University of Coimbra, Largo Marquês de Pombal, 3000-272 Coimbra, Portugal

<sup>a5</sup> Geosciences Department of Évora University, Largo dos Colegiais 2, 7004-516 Évora, Portugal

## Abstract

The Santa Eulália Plutonic Complex (SEPC), located in the Ossa Morena Zone (south Portugal), is composed of a medium- to coarse-grained pink granite (G0-type) and a central grey medium-grained biotite granite (G1-type). Available Rb–Sr data indicates an age of 290 Ma. An emplacement model for the SEPC is proposed, taking into account magnetic fabric, 2D gravity modelling and fluid inclusion planes studies. The G0 and G1 types demonstrate different magnetic behaviour: G0 is considered a magnetite-type granite and G1 is an ilmenite-type granite. The formation of G0 required oxidized conditions related to the interaction of mafic rocks with a felsic magma. The 2D gravity modelling and subvertical magnetic lineations show that the feeder zone of the SEPC is located in the eastern part of the pluton, confirming the role of the Assumar and Messejana Variscan faults in the process of ascent and emplacement. The magma emplacement was controlled by ENE–WSW planar anisotropies related to the final brittle stages of the Variscan Orogeny. The emplacement of the two granites was almost synchronous as shown by their gradational contacts in the field. The magnetic fabric however suggests emplacement of the G0-type first, closely followed by emplacement of the G1-type, pushing the G0 laterally which becomes more anisotropic towards the margin. The G1-type became flattened, acquiring a dome-like structure. The SEPC is a nested pluton with G0-type granite assuming a tabular flat shape and G1-type forming a rooted dome-like structure. After emplacement, SEPC recorded increments of the late Variscan stress field documented by fluid inclusion planes in quartz.

(Received December 27 2013)

(Accepted September 10 2014)

## Keywords:

- anisotropy of magnetic susceptibility;
- isothermal remanent magnetization;
- fluid inclusion planes;
- 2D gravity modelling;
- magma emplacement;
- Variscan Orogeny

## How to cite this article

H. SANT'OVAIA, P. NOGUEIRA, J. CARRILHO LOPES, C. GOMES, M.A. RIBEIRO, H.C.B. MARTINS, A. DÓRIA, C. CRUZ, L. LOPES, R. SARDINHA, A. ROCHA and F. NORONHA. Building up of a nested granite intrusion: magnetic fabric, gravity modelling and fluid inclusion planes studies in Santa Eulália Plutonic Complex (Ossa Morena Zone, Portugal). Geological Magazine, available on CJO2014. doi:10.1017/S0016756814000569.

## Link to this article

<http://dx.doi.org/10.1017/S0016756814000569>