

PUBLISHED BY

INTECH

open science | open minds

World's largest Science,
Technology & Medicine
Open Access book publisher



2800+
OPEN ACCESS BOOKS



96,000+
INTERNATIONAL
AUTHORS AND EDITORS



90+ MILLION
DOWNLOADS



BOOKS
DELIVERED TO
151 COUNTRIES

AUTHORS AMONG
TOP 1%
MOST CITED SCIENTIST



12.2%
AUTHORS AND EDITORS
FROM TOP 500 UNIVERSITIES



Selection of our books indexed in the
Book Citation Index in Web of Science™
Core Collection (BKCI)

Chapter from the book *Worldwide Wound Healing - Innovation in Natural and Conventional Methods*

Downloaded from: <http://www.intechopen.com/books/worldwide-wound-healing-innovation-in-natural-and-conventional-methods>

Interested in publishing with InTechOpen?
Contact us at book.department@intechopen.com

Nursing Interventions in Prevention and Healing of Leg Ulcers: Systematic Review of the Literature

César Fonseca , Manuel Lopes , Ana Ramos ,
Vitor Santos , Antonio Esquinas and Pedro Parreira

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/65719>

Abstract

Aim: The purpose of this study was to define nursing interventions for patients with venous, arterial or mixed leg ulcers.

Methodology: A survey was conducted in EBSCO (CINAHL Plus with Full Text, MEDLINE with Full Text), MedicLatina, Academic Search Complete, with full text articles, published between 2008/01/01 and 2015/01/31, with the following keywords: [(MM "leg ulcer") OR (wound care) OR (wound healing)] AND [(nursing) OR (nursing assessment) OR (nursing intervention)].

Results: The different leg ulcer etiologies require different therapeutic approach to prevention and treatment. Predictive factors were identified associated with healing: patient-centred care, interpersonal relationship, pain control, control of the exudate, education for health self-management, self-care, therapeutic compliance, implementation of guidelines, auditing and feedback on the practices.

Conclusion: Evidence-based practice helps to improve efficiency, safety and quality of nursing care directed to people with leg ulcers or at risk of developing this type of wounds.

Keywords: nursing interventions, leg ulcer, prevention and treatment

1. Introduction

Currently, the needs of healthcare clients are increasingly demanding and complex, as a result of the increase in average life expectancy [1, 2], and the resulting prevalence of chronic

illnesses, such as leg ulcers. The possibility of ulceration increases with age, due to which an exponential increase in its incidence and prevalence is expected [3]. Leg ulcers can be defined as ulceration below the knee on any part of the leg [4], including the foot, and is classified as a chronic wound, that is, a wound that remains stuck in any of the phases of the healing process for a period of 6 weeks or more, or that requires a structured intervention of nursing care [5, 6]. There are several known leg ulcer aetiologies, among which those of venous origin are the most common, at 70% of the cases, followed by those that are arterial in origin, at 10–20% of the cases, and those of a mixed aetiology, at 10–15% of the cases [7]. The main causes of the appearance of leg ulcers are chronic venous hypertension, arterial disease or a combination of the two [5, 6]. The less-frequent causes are neuropathy, infection, vasculitis, neoplasia, blood and metabolic disorders, or lymphedema and disorders that are iatrogenic in origin [5, 6]. The relevance of this problem is supported by statistics, in which 1.5–3 in every 1000 individuals have leg ulcers, with increased prevalence at higher ages, leading to 20 for every 1000 individuals aged more than 80 years [2, 8]. The literature mentions that leg ulcers are interpreted as “a forever healing experience” [9, 10], associated with the fact that 50% of the time, primary healthcare nurses are involved in treating leg ulcers, with the presence of pain in 49–90% of cases, where 50% of venous ulcers heal after 4 months, 20% of venous ulcers heal in between 4 months and 1 year, 20% require a period greater than 2 years to heal, 8% remain unhealed even after 8 years and 69–26% recur in the first year [9].

The repercussions of leg ulcers also have an effect on healthcare expenditure, wherein 1–2% of the total health budget in Western countries [11], of which Portugal is part, is directed to this type of chronic wound [4, 12, 13]. In European countries, an economical investment of approximately 6.5 million Euros/year is estimated in the treatment of leg ulcers [14]. Its impact interferes significantly with human lifestyles, affecting the performance of day-to-day activities, quality of life, functional capacity and self-esteem, leading to workplace absenteeism, financial problems, isolation, sleep disorders and the development of mental illness [15, 16].

Hence, a systematisation is proposed of nursing interventions aimed at persons with venous, arterial or mixed leg ulcers.

2. Methodology

Carried out research in the EBSCO search engine: CINAHL Plus with Full Text, MEDLINE with Full Text, MedicLatina, Academic Search Complete, sought full-text articles, published between 2008/01/01 and 2015/01/31, with the following keywords: [(MM "leg ulcer") OR (wound care) OR (wound healing)] AND [(nursing) OR (nursing assessment) OR (nursing intervention)], filtered through initial question in PICO format. As a starting point in the systematic review of the literature, the following question was formulated in PICO [16] format: “In relation to persons with venous, arterial and mixed leg ulcers (Population), what are the nursing interventions (Intervention) that can influence healing (Outcomes)?” The EBSCO search engine was queried, with access to two databases: CINAHL Plus with Full Text,

MEDLINE with Full Text, MediciLatina, Academic Search Complete, sought full text articles, published between 2008/01/01 and 2015/01/31, with the following keywords: [(MM "leg ulcer") OR (wound care) OR (wound healing)] AND [(nursing) OR (nursing assessment) OR (nursing intervention)].

The process for searching and selecting material for analysis is explained in **Table 1**.

	Protocol
↓	Identification:
	<ul style="list-style-type: none"> • No. of cases identified: CINAHL -310 • No. of cases identified: MEDLINE – 433 • No. of cases identified: MediciLatina - 31 • No. of cases identified: Academic Search Complete - 24
	Screening:
	<ul style="list-style-type: none"> • No. of duplicated cases that were removed – 384 • No. of cases selected – 414
	Inclusion Criteria (complete reading):
	<ul style="list-style-type: none"> • No. of full text articles with inclusion criteria - 11 • No. of full text articles without inclusion criteria - 403
	Articles Included (levels of evidence[16]): Level I – 2; Level II – 1; Level IV – 3; Level V – 4; Level VI – 1

Table 1. Process for searching and selection for the systematic review of the literature.

3. Results

In order to make the methodology used easy to understand and transparent, the list of 12 articles selected is explained (**Table 2**) for the body of analysis, which formed the basis for the preparation of the discussion and the corresponding conclusions, having been subjected to classification by levels of evidence.

Level of evidence/ articles	Method	Participants	Interventions	Results
Level of evidence – V [17]	Systematic review of the literature	Review of three guidelines	Critical review of the guidelines in order to	Recommendations are prepared on how to assess leg ulcers, as well as

Level of evidence/ articles	Method	Participants	Interventions	Results
		(Medline and Cochrane)	prepare a set of recommendations	how to treat the different aetiologies: venous, arterial and mixed
Level of evidence— V [8]	Systematic review of the literature	31 articles relating to people with venous and/or mixed leg ulcers	The articles were analysed in order to be able to perceive failure To comply with the treatment	Pain, discomfort and different lifestyles are some of the reasons why leg ulcer patients did not comply with the treatment. Healthcare professionals must focus on problems reported by the patients in order to be able to help them overcome these problems and to motivate them to take the treatment
Level of evidence— V [14]	Systematic review of the literature	Publications on social support and persons with leg ulcers	Several studies are compared in order to establish a relationship between the effect of social support on the healing of venous leg ulcers as well as on recurrence	Social support is important for persons with venous leg ulcers, as this support is necessary during as well as after the wound is healed, in order to prevent recurrence
Level of evidence— V [18]	RCT	All the people with leg ulcers in the region of Skaraborg (Sweden).	Identification of people with leg ulcers, their aetiology, prevalence and ongoing treatment	It was observed that venous leg ulcers continue to be the most prevalent, followed by arterial leg ulcers. In general, there is a reduction in prevalence in relation to previous studies
Level of evidence— I [19]	Systematic review of the literature	325 database articles (Cinahl, Medline and Cochrane)	Critical review of the articles found in order to prepare a set of recommendations	Recommendations are prepared on how to assess and intervene among patients with arterial leg ulcers, on the following points: debridement, dressing selection, infection control, nutrition, pain control
Level of evidence— I [20]	Systematic review of the literature	180 database articles (Medline and Cochrane)	Critical review of the articles found in order to prepare a set of recommendations	Recommendations are prepared on how to assess, prevent and treat people with venous leg ulcers
Level of evidence— IV [6]	Retrospective study	Eight people with mixed leg ulcers treated by two specialist nurses	Identify the role of the specialist nurse in controlling ulceration	The ulcers heal in between 6 and 30 weeks after the first application of an inelastic bandage system. This intervention was well tolerated by all

Level of evidence/ articles	Method	Participants	Interventions	Results
				the patients, and no adverse effects were recorded
Level of evidence— IV [12]	Prospective study	Five persons with leg ulcers, with assistance at home and over the telephone, for 12 weeks	Identify the strategies in the promotion of therapeutic compliance	Individualisation of information, training/instruction increases therapeutic compliance
Level of evidence— V [13]	Systematic review of the literature	Five articles resulting from searches on MEDLINE, British Nursing Index and Cumulative Index to Nursing and Allied Health Literature (CINAHL)	Control the level of exudate in leg ulcers	Proper control of the amount of exudate minimises the impact on quality of life, damage to the wound bed, on the perilesional skin, reduces the risk of infection, days required for healing of the wound and health expenditure
Level of evidence— VI [3]	Case study	A person with a venous leg ulcer	Analyse the influence of self-care on leg ulcer healing	The capacity for self-care stimulated by the patient's empowerment reduces the need for seeking health care
Level of evidence— IV [21]	Case study— control	11 persons with pressure ulcers and 20 with leg ulcers, in primary care	Verify the advantages of the use of absorbent dressings in controlling the exudate of venous ulcers	The control of exudate promotes healing of chronic wounds, control of pain and negative psychosocial effects associated with the smell and change of dressings

Table 2. Body of analysis.

4. Discussion of the data

The evaluation of the awareness, expectations, quality of social support, need for information enables the adaptation of education, training and instruction on healthy lifestyles, wound care, physical exercise and elevation of lower limbs in venous aetiology. The patient, on feeling involved, with the use of an easy-to-understand language, increases his or her motivation, self-efficacy and capacity for self-care [12].

The creation of spaces (Leg Clubs) has proved to be a fundamental strategy in therapeutic compliance [10], where nurses with specific training in the area of leg ulcers promote social

interaction between patients with the same type of ulcer, evaluate the support required by each individual, provide training aimed at self-care and case management, provide the corresponding treatment and constant monitoring [4–9]. The result of the implementation of this project was the reduction in pain intensity, significant progress in healing and an increase in quality of life, specifically at the workplace, in moods, in mobility, sleep patterns and other aspects [10]. The positive effect of this model is also reflected at the social level [9], given that more extended social contact with people who have or have had the same problem, reduces social isolation and provides effective coping mechanisms for dealing with the crisis situation - the illness [3, 4, 16].

Venous ulcer	Arterial ulcer	Mixed ulcer
<ul style="list-style-type: none"> Apply compressive therapy if the Ankle-Brachial Index (ABI) is greater than 0.8 Select the type of compression to be used: multi-layer compression system below the knee (general treatment); reduced compression system (in case the patient does not tolerate higher compressions); compression stockings (after the ulcer's healing); intermittent compression system (can be used in isolation or together with another compression system, in order to increase venous return) Select the compressive therapy material to be used: elastic bandages (long-stretch), considered more effective; non-elastic bandages (short-stretch), which cause less discomfort/pain; multi-layer elastic bandage systems (two, three or four layers); elastic stockings; multi-layered elastic stockings; zinc-impregnated bandage (Unna Boot) together with elastic bandages Apply compression stockings with customised measurement, if the ulcer has healed Refer for vascular surgery in the following situations: no reduction in the size of the ulcer after 30 days of treatment; ulcer present for more than 6 months; intolerance to compressive therapy; ineffectiveness in pain control; frequent recurrence 	<ul style="list-style-type: none"> Assess the lower limbs, frequently, in relation to: functional capacity; colouring, temperature; capillary reperfusion; sensitivity; presence of dorsalis pedis, posterior tibial pulse; signs of neuropathy Perform cleaning of the wound with non-cytotoxic products Do not debride dry and stable necrotic tissue without proper assessment of the perfusion through vascular surgery, not applying any moisture retaining bandage material Debride the necrotic tissue, based on a multi-professional decision, using autolytic and enzymatic debridement Provide training in the following structural areas: control of associated pathologies; giving up smoking and drinking; encouraging the ingestion of food rich in vitamin B6 (increases HDL-C and reduces triglycerides), such as potato, banana, chicken breast, sunflower seeds, salmon, tuna, avocado, and others; prevention of chemical, thermal or mechanical trauma on the lower limbs; skin care; use of sui footwear and non-compressive stockings Institute a regular physical exercise programme, for patients with intermittent claudication, based on 30–60-min walks (3 days a week at least), wherein the patient may stop and rest in case of pain Refer to vascular surgery if: ABI is lower than 0.8; signs and symptoms of infection; continued pain at rest, even with the limb dropped; absence of both, pedis and posterior tibial pulses If ABI is lower than 0.5, refer urgently for observation by vascular surgery 	<ul style="list-style-type: none"> Apply reduced compression (between 23 and 30 mmHg) in the presence of oedema Refer for vascular surgery, if there is no healing development and ABI is lower than 0.5

Table 3. Nursing interventions in venous, arterial and mixed leg ulcers.

On initial assessment, it is crucial to cover the history of health: associated co-morbidities, habitual therapy, psycho-emotional state, influence of odour on social life, nutritional state, presence and intensity of pain and individual treatment preference. In evaluating the amount of exudate, it is recommended to document the saturation of the absorbent dressing and support/compression bandage, instead of rating it as minimum (+), moderate (++) and high (+++), in order to increase the record's objectivity. If the perilesional skin is macerated, this indicates that the dressings must be applied more frequently, or that the selected material is not the most suited for controlling the exudate. In leg ulcers subjected to compression therapy, in venous aetiology, dressings that allow for evaporation of the exudate through their semi-permeable covering cease to be effective. In light of the above, the use of hydrofibre and alginates is recommended. The application of negative pressure on the wound bed and the use of protective sprays/creams on the surrounding area are measures to be considered in the case of ulcers with hard-to-control exudate [13, 14].

The application of compression bandages is considered an essential element in the treatment of venous leg ulcers. The effect of compression on mixed leg ulcers can be beneficial in reducing local oedema, in improving microcirculation, contributing to improving arterial flow and improving venous as well as lymphatic drainage. Still, the application of compression on legs with mixed aetiology requires strong additional care, as it is generally contraindicated for the reduction of arterial flow, causing greater tissue damage, as shown in study [6], with the use of inelastic bandage. The number of layers to be applied is normally decided based on the malleolar circumference, based on the manufacturer's instructions (one layer is applied when the malleolar circumference is ≤ 25 cm, and two layers when it is > 25 cm). However, in mixed aetiology, it is recommended to start the treatment with only one layer, for less-elevated compression levels. The interventions provided by specialist nurses when dealing with chronic wounds increase gains in health, due to their expertise in using tools for assessing compressive therapy: Doppler ultrasound and ankle-brachial index (ABI).

Concerning nursing interventions in the prevention and treatment of venous, arterial or mixed leg ulcers, it is fundamental to know the patient's clinical history (personal background, chronic pathologies, current state of the client) and the history of the ulcer (source, time, treatments performed) [6, 12, 14–16, 18, 22–24]. On meticulously evaluating the characteristics of the wound (size, depth, exudate, wound bed, type of tissues, perilesional skin, pain) [4, 18], the decision must be made in partnership with the client (**Table 3**) in order to establish common goals [15, 17–21].

Thus, the treatment must involve pain prevention [10], preparation of the wound bed [6–9], wound cleaning [14–16], management of products to be applied to the bed and perilesional skin [7], joint selection of the type of material for application of compressive treatment and preparation of a physical exercise plan [8, 14–16, 18, 22], continuous client empowerment [17], referral to specialities in case of allergic reactions [13], need for supplementary therapies and/or non-effective treatments carried out in which the ulcer/state of the client deteriorates.

5. Conclusions and implications for the practice of nursing

The presence of social support was the aspect most mentioned by the people as essential in their process of adaptation, whether provided by significant persons, or through contact with people in similar situations (self-help groups) or by the nurse. The education for health self-management was considered of utmost importance in controlling other associated chronic illnesses, in the reduction of other existing risk factors and for creating physiological conditions that favoured better healing. The monitoring of the ulcer's characteristics, physical activity, nutritional diet, favoured healing and improved the perceived quality of life. Continuous and up-to-date training of nurses providing care to leg ulcer patients emerged as another aspect positively associated with the effectiveness and excellence of the interventions carried out.

An approach centred on the patient and on pain control, mainly when the therapeutic plan involves compression on the wound bed, is key determinants in increasing participation and involvement in the therapeutic plan.

Conflicts of interest

The author declares no conflict of interest.

Author details

César Fonseca^{1*}, Manuel Lopes¹, Ana Ramos², Vitor Santos¹, Antonio Esquinas³ and Pedro Parreira⁴

*Address all correspondence to: cesar.j.fonseca@gmail.com

1 University of Évora, Évora, Portugal

2 Hospital Middle Tejo Center, Tomar, Portugal

3 Intensive Care Unit, Hospital Morales Meseguer, Murcia, Spain

4 Coimbra Nursing School, Coimbra, Portugal

References

- [1] Instituto Nacional de Estatística. *Projeções de População Residente em Portugal 2008–2060*. Instituto Nacional de Estatística – Ministério da Saúde.. Portugal. Lisboa. 2014.
- [2] Van Hecke A, Grypdonck M, Defloor T. A review of why patients with leg ulcers do not adhere to treatment. *Journal of Clinical Nursing* [serial on the Internet]. (2009, Feb),

- [cited January 4, 2015]; 18(3): 337–349. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=105629598&site=ehost-live>
- [3] Yarwood-Ross L, Haigh C. Managing a venous leg ulcer in the 21st century, by improving self-care. *British Journal Of Community Nursing* [serial online]. October 2012;17(10):460–465. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=108112049&site=ehost-live>
- [4] Brown A. Does social support impact on venous ulcer healing or recurrence?. *British Journal of Community Nursing* [serial on the Internet]. (2008, Mar 2), [cited January 4, 2015]; 13(3): S6. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=105909037&site=ehost-live>
- [5] Werchek S. Diagnosis and treatment of venous leg ulcers. *The Nurse Practitioner* [serial on the Internet]. (2010, Dec), [cited January 4, 2015]; 35(12): 46–53. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=104972282&site=ehost-live>
- [6] Neill K, Turnbull K. Use of specialist knowledge and experience to manage patients with mixed aetiology leg ulcers. *Journal Of Wound Care* [serial online]. April 2012;21(4):168–174. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=104422923&site=ehost-live>
- [7] Vowden P. Leg ulcers: assessment and management. *Independent Nurse* [serial on the Internet]. (2010, Nov 22), [cited January 4, 2015]; 12(3): 30–33. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=104956673&site=ehost-live>
- [8] Hecke A, Grypdonck M, Defloor T. Interventions to enhance patient compliance with leg ulcer treatment: a review of the literature. *Journal of Clinical Nursing* [serial on the Internet]. (2008, Jan), [cited January 4, 2015]; 17(1): 29–39. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=106013847&site=ehost-live>
- [9] Chamanga E. How can community nurses improve quality of life for patients with leg ulcers?. *Nursing Times* [serial online]. March 16, 2010;106(10):15–17. <http://search.ebscohost.com/login.aspx?direct=true&db=mnh&AN=20426295&site=ehost-live>
- [10] Ebbeskog B, Emami A. Older patients' experience of dressing changes on venous leg ulcers: more than just a docile patient. *Journal of Clinical Nursing* [serial on the Internet]. (2005, Nov), [cited January 4, 2015]; 14(10): 1223–1231. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=106385326&site=ehost-live>
- [11] EUROSTAT. Population projections 2008–2060: from 2015, deaths projected to outnumber births in the EU27. S.1: Eurostat Press Office; Lanzieri: Eurostat News. 2008:1–4. Available from: <http://ec.europa.eu/eurostat>
- [12] Van Hecke A, Beeckman D, Grypdonck M, Meuleneire F, Hermie L, Verhaeghe S. Knowledge Deficits and Information-Seeking Behavior in Leg Ulcer Patients: An Exploratory Qualitative Study. *Journal Of Wound, Ostomy & Continence Nursing*

- [serial online]. July 2013;40(4):381–387. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=107958718&site=ehost-live>
- [13] Menon J. Managing exudate associated with venous leg ulceration. *British Journal Of Community Nursing* [serial online]. June 2, 2012; 25(4):S6–s16. <http://search.ebscohost.com/login.aspx?direct=true&db=mnh&AN=22875033&site=ehost-live>
- [14] Martin F. Educational challenges and requirements for managing leg ulcers in the communi. *British Journal Of Community Nursing* [serial online]. June 2, 2014; 12(2):S32–6. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=103960614&site=ehost-live>
- [15] Edwards H, Courtney M, Finlayson K, Lindsay E, Lewis C, Chang A, et al. Chronic venous leg ulcers: effect of a community nursing intervention on pain and healing. *Nursing Standard* [serial on the Internet]. (2005, Sep 7), [cited January 4, 2015]; 19(52): 47–54. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=106533450&site=ehost-live>
- [16] Melnyk B, Fineout-Overholt E. *Evidence-based practice in nursing & healthcare: a guide to best practice* [monograph on the Internet]. Philadelphia, Pennsylvania: Lippincott Williams & Wilkins; 2005. <http://file.zums.ac.ir/ebook/208-Evidence-Based%20Practice%20in%20Nursing%20&%20Healthcare%20-%20A%20Guide%20to%20Best%20Practice,%20Second%20Edition-Be.pdf>
- [17] Guideline for management of wounds in patients with lower-extremity venous disease: complete summary [monograph on the Internet]. U.S. Department of Health and Human Services. U.S. Department of Health and Human Services, 2005. <https://www.guideline.gov/summaries/summary/38249/guideline-for-management-of-wounds-in-patients-with-lowerextremity-venous-disease>
- [18] Forssgren A, Fransson I, Nelzén O. Leg Ulcer Point Prevalence can be Decreased by Broad-scale Intervention: a Follow-up Cross-sectional Study of a Defined Geographical Population*. (Cover story). *Acta Dermato-Venereologica* [serial on the Internet]. (2008, May), [cited January 4, 2015]; 88(3): 252–256. <http://search.ebscohost.com/login.aspx?direct=true&db=mnh&AN=18480924&site=ehost-live>
- [19] Assessment and management of venous leg ulcers: complete summary [monograph on the Internet]. US National Guideline Clearinghouse U.S. Department of Health and Human Services; 2008. http://www.gacguidelines.ca/site/GAC_Guidelines/assets/pdf/LEGU05-6_Venous_Leg_Ulcer_Summary.pdf
- [20] Guideline for management of wounds in patients with lower-extremity arterial disease: complete summary [monograph on the Internet]. US National Guideline Clearinghouse U.S. Department of Health and Human Services; 2008. <https://www.guideline.gov/summaries/summary/38249/guideline-for-management-of-wounds-in-patients-with-lowerextremity-venous-disease>
- [21] Yarwood-Ross L, Haigh C. Managing a venous leg ulcer in the 21st century, by improving self-care. *British Journal Of Community Nursing* [serial online].

October 2012;17(10):460–465. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=108112049&site=ehost-live>

- [22] Van Hecke A, Grypdonck M, Defloor T. The clinical nursing competences and their complexity in Belgian general hospitals. *Journal of Advanced Nursing* [serial on the Internet]. 2006;56(6):669–678. [cited January 13, 2011]. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=106290657&site=ehost-live>
- [23] Domeij D, Flodén M. POPULATION AGING AND INTERNATIONAL CAPITAL FLOWS. *International Economic Review* [serial on the Internet]. (2006, Aug), [cited January 4, 2015]; 47(3): 1013–1032. <http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=21678955&site=ehost-live>
- [24] Heinen M, Evers A, Van Uden C, CJM, PCM, Van Achterberg T. Sedentary patients with venous or mixed leg ulcers: determinants of physical activity. *Journal of Advanced Nursing* [serial on the Internet]. (2007, Oct), [cited January 4, 2015]; 60(1): 50–57. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=106190961&site=ehost-live>

INTECH

