


Actividades e interesses do Instituto de Ciências da Terra relacionados com a Previsão Numérica do Tempo

Rui Salgado com

ICT – Grupo 1 – Atmospheric Sciences, Water and Climate



Models in use in Evora (ICT)

- NWP in mesoscale: BRAMS, WRF, AROME, MESO-NH
- Surface (coupled or stan-alone): SURFEX platform
- Lake models: FLake and LAKE
- Radiation: Ecrad and LibRadtran

Observations

- Atmosphere
 - Évora Observatory, Meteorological and radiometric networks, Field campaigns
- Lake (Alqueva reservoir)
 - floating platform, field campaigns

Computational infrastructure

- Terra Cluster:
 - 128 processing cores (32 per node)
 - 157.1GB. total RAM memory
 - The processors are Xeon(R) CPU E5-2660 0 @ 2.20GHz
- Khromeleque Cluster:
 - 16 nodes with 4 AMD Opteron 6376 (64 cores 64 bits a 2.3 GHz)
 - 1 head node (2 x Intel Xeon E5 2650, 32 GB)
 - node 2 x Intel Xeon E5 2640v2, 64 GB, com 2 Xeon Phi 7210
 - node 2 x Intel Xeon E5 2690v2, 128 GB, com 1 nVidia Tesla K20
 - Infiniband backbone FDR 4x (56 Gbps)
 - distributed filesystem 100 TB




- A state-of-art "non-hydrostatic non-hydrostatic mesoscale atmospheric model developed by the Laboratoire d'Aérologie and by CNRM/Météo-France.
- Incorporates a non-hydrostatic system of equations, for dealing with scales ranging from large (synoptic) to small (large eddy) scales;
- Has a complete set of physical parameterizations, which are particularly advanced for the representation of clouds and precipitation;
- Allows a multi-scale approach through a grid-nesting technique;
- Is a versatile code, vectorized, parallelized, operating in 1D, 2D or 3D designed to handle real situations as well as academic cases;
- Is coupled with a chemistry module and a lightning module;
- Is coupled to a Forest Fire propagation model (FOREFIRE)
- Has observation operators that compare model output directly with satellite observations, radar, lidar and GPS.

Reference: Lac, C. et al., 2018

- Is coupled to the SURFEX platform of surface models for the representation of surface atmosphere interactions
 - SURFEX is used in AROME, HARMONIE, ...
- Share the *Physics* with AROME (runs operationally in IPMA)
- Share the radiative code of IFS (now EcRAD)
- Is the research model of Méteo-France: new developments may be included in the operational versions.
- participation in the steering committee and in biennial meetings



- SURFEX (Surface Externalisée, in French, Masson *et al.*, 2013) is a surface modelling platform developed by Météo-France in cooperation with the scientific community.
- SURFEX is composed of various physical models for **natural land surface, urbanized areas, lakes and oceans**.
- It also simulates chemistry and aerosols surface processes and can be used for assimilation of surface and near surface variables.
- SURFEX can be used in
 - stand alone mode
 - coupled to an atmospheric model.



- SURFEX is used in the NWP models of the ALADIN - HIRLAM consortiums


SURFEX basis

Land cover database




Tile %

Tiling approach



Flux calculation




Models

Seas and oceans:
 Prescribed SST, Charnock formulation
 ECUME (multi-campaign parameterization), 1D Ocean Mixed Layer model



Lakes :
 Prescribed LST, Charnock formulation
 FLake lake model




Soil and vegetation: ISBA
 Force restore or diffusion for heat and water transfers in the soil




Town: TEB
 Canyon concept, detailed radiative scheme
 Vegetated buildings, impact of trees in canyon

Model parameters




Patrick Le Moigne et al., 2018

- Water temperature profile, two layers:
 - Mixed Layer (Constant Temperature)
 - thermocline (temperature profile is defined by a shape factor)
 - 4 variables (T_s , T_b , h and c)
 - important parameters:
 - lake depth
 - **Extinction Coefficient**
- Computationally cheap and good representation of Surface temperature evolution
- In use in SURFEX / AROME / Meso-NH and in ECMWF, DWD and MetOffice



- Flake was tested with data from Alqueva, measured on a floating platform
- The tests were used to link FLake into SURFEX
Salgado and Le Moigne (2010)
- Collaboration in IFS implementation
Balsamo et al. (2012)



Extinction Coefficient: measurements




- Development of a FieldSpec UV/VNIR da ASD coupled to an optical cable and a cosine receptor
- Apparatus developed by Miguel Potes (PhD thesis)

- Measurements of:
 - Underwater irradiation
 - Extinction coefficient
- Locals:
 - Alqueva reservoir (Potes et al., 2013, 2017)
 - Thau lagoon (Thaumex, LeMoigne et al., 2013)




- Global land cover map of ecosystems (273 in version 2)
- Dataset of surface parameters associated for each grid mesh in tabular form:
 - albedo, LAI, % of vegetation cover, z_0 , minimum stomatal resistance, and root zone.
- Now (Second Generation) is at 300m resolution
- Used by NWP models to produce physiographic maps ("climatologies")



Ecoclimap-II map over Europe
Faroux et al., 2013

- Operational version used at IPMA don't included Alqueva
- The ALqueva was inserted in the Ecoclimap database by Carlos Policarpo et al (2017)
- and was implemented and tested by Sónia Assunção (Master thesis, 2017) e Maria Monteiro in a collaboration ICT/IPMA (see poster)



Ecoclimap-II map over Europe
Faroux et al., 2013








Figura 3 – Representação da base de dados ECOCLIMAP_II_v2.3 sem a fisiografia da albufeira de Alqueva (à esquerda) e com a fisiografia (à direita). As áreas a verde representam superfícies com água.

- library for radiative transfer <http://libradtran.org>
- Set of C and Fortran routines to compute solar and thermal radiation in the atmosphere (3D). Free under GNU GPL.
- Several examples of works done in ICT

Global irradiation: aerosols radiative forcing







Remote sensing of cloud properties

clear sky irradiation at the surface

diffuse irradiation in the circumsolar region



sensitivity tests to: aerosols and water vapour




Applications

Climate impact of Alqueva

- Effects on Temperature and humidity (Miranda et al., 1995, Salgado, 2006)
- Effects on fog (Salgado, 2006, Policarpo et al., 2017)
- Characterization of the Lake Breeze (Iakunin et al., 2018)
- Effects in the atmospheric electrical field Lopes et al., 2016; Nicol et al., 2018
- Heat, water vapour and CO_2 fluxes (Potes et al., 2017)





ESTUDO DE IMPACTE AMBIENTAL DO ALQUEVA
CLIMA
Relatório Final



Pedro M. A. Miranda
Francisco Abreu
Rui Salgado

Relatório Técnico do ICAT
Janeiro 1995



Representation of lakes in NWP



⌚ Improvement of lake parameterization in MF models

Due to the increase of horizontal resolution in models

Need to improve the diurnal cycle over lake areas

A step forward to data assimilation




⌚ SURFEX implementation of FLake model

Salgado and Le Moigne, 2010

⌚ Field Campaigns validations




THAUMEX, South-France : Le Moigne et al., 2013



- A monthly climatology of FLake variables has been created using a Lake Planet experiment
 - This climatology is used by the ECMWF model
 - Example: Climatological Ice Depth (Above water)

Water quality forecast


- Forecast of water quality parameters using NWP
 - In-line or post-processing
- Using a Flake type approach (EcoFlake)
- Try a more physical (and chemical) based model (LAKE)
 - see Maksim's presentation



pattern
of the
profiles
are
similar.

- The use of solar energy in electricity generation is growing
- There are many new projects in Portugal, namely in Alentejo
- There is a need to improve solar radiation predictions based on NWP
 - Global Radiation
 - Direct Normal Radiation, for

- Concentrating solar photovoltaic (CPV) and thermal (CST) technologies require direct normal irradiance (DNI) forecasts
- DNI is a new ECMWF output, not validated
- New Radiative code (2017): EcRad
- 1st step: Access the quality of ECMWF forecast of DNI
 - Francisco Lopes et al. (2018)
 - See his presentation



Short-term forecasts of GHI and DNI for solar energy systems operation: assessment of the ECMWF integrated forecasting system in southern Portugal


Francis M. Lopes^{a,b,*}, Hugo G. Silva^{a,b,c}, Rui Salgado^{b,c}, Afonso Cavaco^d, Paulo Canhoto^{b,c}, Manuel Collares-Pereira^{a,b,c,d}

^a Renewable Energies Chair, University of Évora, IIFA, Palácio do Vimioso, Largo Marquês de Marialva, Apart. 94, 7002-554 Évora, Portugal

^b Earth Sciences Institute, University of Évora, Rua Romão Ramalho, 59, 7000-671 Évora, Portugal


^c Department of Physics, School of Sciences and Technology, University of Évora, Rua Romão Ramalho, 59, 7000-671 Évora, Portugal

^d Portuguese Solar Energy Institute, IIFA, Palácio do Vimioso, Largo Marquês de Marialva, Apart. 94, 7002-554 Évora, Portugal



Improve Global Radiation prediction

- Study and development of a solar radiation predicting algorithm based on ECMWF's forecasts and ANNs
 - Sara Pereira et al. (2018)
(See her presentation)
- Post-processing technique
- ANN was trained and validated using data from ICT and IPMA stations



Orographic Precipitation - Madeira



1. Introduction


Starting point

HPE in the 2009/2010 winter and the disaster on February 2010.

LARGE SCALE

A cyclone and frontal systems affecting the island.

Example: 20 February 2010




Couto et al. (2012)

COUTO FT, SALGADO R, COSTA MJ (2012). Analysis of intense rainfall events on Madeira Island during the 2009/2010 winter. *Natural Hazards and Earth System Sciences*. 12: 2225-2240.

MESOSCALE

MESO-NH simulations at 1 km resolution showed the **orographic effect** for HPE over the island, and maximum of accumulated precipitation in the highlands.




3. Results


Large scale environment

The meridional water vapour transport through the Atmospheric Rivers (ARs) may be identified from the Aqua-AIRS or ECMWF analyses, as shown for the episode of 20 February 2010.

AIRS observations



ECMWF analyses



From an operational point of view, this information helps the HPE forecast, since this transport may be easily identified over the ocean several hours before it reaches the island.


Some notes on the forecast of heavy precipitation events for Madeira

3. Results

Local effects: Part I – AROME results for precipitation distribution

=> The model captures the **orographic signature** in the accumulated precipitation, as well as the **different rainfall patterns**. This result is evidenced by comparing the AROME outputs with those obtained with the MESO-NH model.

AROME outputs at 2.5 km resolution for four periods considered by Couto et al. (2016).



Some notes on the forecast of heavy precipitation events for Madeira



4. Conclusions and perspective

Conclusions

The satellite data may support the monitoring of water vapour transport over the Atlantic Ocean.

From the AROME analysis, at least when the orographic effects lead to quasi-stationary orographically induced precipitating systems, it is able to reproduce satisfactorily the different rainfall patterns in a short-range forecast.

For the season, the AROME captured the main significant precipitation periods in the three regions considered, despite some under or overestimation of the local values.

Perspective

Besides the orographic signature or location of the accumulated precipitation, a more detailed analysis about the AROME performance is suggested for future works, also considering other situations.



Some notes on the forecast of heavy precipitation events for Madeira

- Flavio Couto slides
- based on Couto et al. (2012, 2015, 2016 and 2017)
- Last work in collaboration with IPMA



Dust emission, diffusion and transport

- Simulation of dust events from the emission to the effects on cloud
- Case studies using Meso-NH
- Work done by Dina Santos et al. (2011, 2013)



Simulated aerosol optical depth



Pollen monitoring and forecast

- AutoPollen: Automatic Pollen Detection Network
 - support of EUMETNET and MeteoSwiss
- We want to participate
- It should be good if IPMA join us
- More than 100 million Europeans suffer from allergic rhinitis and 70 million from asthma
- Pollen forecasting are necessary to prevent pollen allergy
- It will be a new forecast product




Benefits of real-time pollen monitoring

- Timely information
- Input for forecasts and models
- Improve the quality, e.g. of information and forecasts
- Partnerships, convergence with aerosol and air quality measurements
- New products and services



- **Meso-NH - ForeFire** coupled model can be used to simulate and predict fine- to large-scale effects of wildfires
 - *Fillipi et al. (2018)*
- Operational at Corsica
- Maybe coupled to AROME
- To be tested in near future during CILIFO (ITERREG Project)



Atmospheric Science Observatory



Sun-sky photometer
CIMEL



Temperature and humidity
profilerMicrowave Radiometer



Raman Lidar System




IRGASON, integrated
eddy covariance
system to access
fluxes of momentum,
energy, vapour and
CO₂

Ceilometer
VAISALA
CL31



ALOP Field Experiment

- Observação simultânea da qualidade da água e do estado da atmosfera



Agradecimentos

COMPETE 2020 através do ICT (UID / GEO / 04683/2013) com a referência POCI-01-0145-FEDER-007690 e dos projectos ALOP (ALT20-03-0145-FEDER-000004) e DNI-A (ALT20-03-0145-FEDER-000011).



References

References

- Balsamo, G., R. Salgado, E. Dutra, S. Boussetta, T. Stockdale, M. Potes, 2012: On the contribution of lakes in predicting near-surface temperature in a global weather forecasting model, *Tellus-A*, 64, 15829, DOI: 10.3402/tellusa.v64i0.15829
- Couto, F. T., Salgado, R., and Costa, M. J., 2012: Analysis of intense rainfall events on Madeira Island during the 2009/2010 winter, *Nat. Hazards Earth Syst. Sci.*, 12, 2225-2240, doi:10.5194/nhess-12-2225-2012
- Couto, F. T., Salgado, R., Costa, M. J. and Prior, V., 2015: Precipitation in the Madeira Island over a 10-year period and the meridional water vapour transport during the winter seasons. *International Journal of Climatology*, 35(13), 3748–3759. doi: 10.1002/joc.4243.
- Couto, F. T., Ducrocq, V., Salgado, R., Costa, M. J., 2016: Numerical simulations of significant orographic precipitation in Madeira island. *Atmospheric Research*, 169, Part A, 102-112, doi:10.1016/j.atmosres.2015.10.002.
- Couto, F. T., Ducrocq, V., Salgado, R. and Costa, M. J. (2017), Understanding significant precipitation in Madeira island using high-resolution numerical simulations of real cases. *Q.J.R. Meteorol. Soc.*, 143: 251–264. doi:10.1002/qj.2918
- Costa, M.J., R. Salgado, D. Santos, V. Levizzani, D. Bortoli, A. M. Silva and P. Pinto, 2010: Modelling of orographic precipitation over Iberia: a springtime case study. *Adv. Geosci.*, 25, 103–110. doi:10.5194/adgeo-25-103-2010
- Faroux, S., Kaptué Tchuenté, A. T., Roujean, J.-L., Masson, V., Martin, E., and Le Moigne, P.: ECOCLIMAP-II/Europe: a twofold database of ecosystems and surface parameters at 1 km resolution based on satellite information for use in land surface, meteorological and climate models, *Geosci. Model Dev.*, 6, 563-582, doi:10.5194/gmd-6-563-2013, 2013.
- Filippi, J-B., Bosseur, F., Mari, C. and Lac, C. (2018). Simulation of a Large Wildfire in a Coupled Fire-Atmosphere Model. *Atmosphere*, 9, 218.
- Iakunin, M., Salgado, R., and Potes, M., 2018: Breeze effects at a large artificial lake: summer case study. *Hydrol. Earth Syst. Sci.*, 22, 5191-5210, <https://doi.org/10.5194/hess-22-5191-2018>.
- Lac, C. et al., 2018: Overview of the Meso-NH model version 5.4 and its applications, *Geosci. Model Dev.*, 11, 1929-1969.
- Le Moigne, P., D. Legain, F. Lagarde, M. Potes, D. Tzanos, E. Moulin, J. Barrié, R. Salgado, G. Messiaen, A. Fiandrino, S. Donier, O. Traullé, and M. J. Costa, 2013: Evaluation of the lake model FLake over a coastal lagoon during the THAUMEX field campaign, *Tellus A*, 65, 20951. doi: 10.3402/tellusa.v65i0.20951.
- Lopes, F., Silva, H., Salgado, R., Potes, M., Nicoll, K., & Harrison, R. (2016). Atmospheric electrical field measurements near a fresh water reservoir and the formation of the lake breeze. *Tellus A*, 68. doi:<http://dx.doi.org/10.3402/tellusa.v68.31592>

References

Lopes, F. M. Silva, H. G., Salgado, R., Cavaco, A., Canhoto, P., Collares-Pereira, M., 2018: Short-term forecasts of GHI and DNI for solar energy systems operation: assessment of the ECMWF integrated forecasting system in southern Portugal, . Solar Energy,, Volume 170, 14-30, <https://doi.org/10.1016/j.solener.2018.05.039>

Masson, V. et al. 2013.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of earth surface variables and fluxes, Geosci. Model Dev., 6, 929-960, doi: 10.5194/gmd-6-929-2013.

Miranda, P.M.A., F. Abreu e R. Salgado, 1995; Estudo de Impacte Ambiental do Alqueva - Clima; Relatório Final. Relatório Técnico do ICAT, 186pp.

Nicoll, K. A.; R.G. Harrison, H.G. Silva, R. Salgado, M. Melgao, D. Bortoli, 2018: Electrical sensing of the dynamical structure of the planetary boundary layer. Atmos, 2018, doi: 10.1016/j.atmosres.2017.1 1.009

Pereira, S., Canhoto, P., Salgado, R., Costa, M.J., 2018. Study and development of a solar radiation predicting algorithm based on ecmwf ' s for ecasts and ANNs, in: grand renewable energy 2018. Yokohama.

Policarpo, C., Salgado, R. and Costa, M. J. (2017): Numerical Simulations of Fog Events in Southern Portugal. Advances in Meteorology, vol. 2017, Article ID 1276784, 16 pages, 2017. doi:10.1155/2017/1276784

Potes, M. Salgado, R., Costa M.J., Morais, M., Bortoli, D., Kostadinov, I. and Mammarella, I., 2017: Lake–atmosphere interactions at alqueva reservoir: a case study in the summer of 2014. Tellus A 2016, 00, 1272787, <http://dx.doi.org/10.1080/16000870.2016.1272787>

Potes, M., Costa, M. J., Salgado, R., Bortoli, D., Serafim, A. Le Moigne, P., 2013: Spectral measurements of underwater downwelling radiance of inland water bodies. Tellus A. 65, DOI: 10.3402/tellusa.v65i0.20774.

Salgado, R., 2006: Interacção solo - atmosfera em clima semi-árido. Dissertação apresentada à Universidade de Évora para a obtenção do grau de Doutor em Física.

Salgado, R. and P. Le Moigne, 2010: Coupling of the FLake model to the Surfex externalized surface model. *Boreal Env. Res.* 15: 231–244.

Santos, D., Costa, M.J., Silva, A.M., Salgado, R., Domingues, A. and Bortoli, D. (2011): ‘Saharan desert dust radiative effects: a study based on atmospheric modelling’, Int. J. Global Warming, Vol. 3, Nos. 1/2, pp.88–102.

Santos, D., M.J. Costa, A.M. Silva, R. Salgado, 2013: Modeling Saharan desert dust radiative effects on clouds, Atmospheric Research, Volume 127, June 2013, Pages 178-194, ISSN 0169-8095, 10.1016/j.atmosres.2012.09.024.