



Role of QseG membrane protein in beneficial enterobacterial interactions with plants and *Mesorhizobia*

Denise Pereira Torres, Ana Paço, Esther Menéndez, Pedro F. Mateos & Clarisse Brígido

To cite this article: Denise Pereira Torres, Ana Paço, Esther Menéndez, Pedro F. Mateos & Clarisse Brígido (2021) Role of QseG membrane protein in beneficial enterobacterial interactions with plants and *Mesorhizobia*, Journal of Plant Interactions, 16:1, 510-521, DOI: 10.1080/17429145.2021.2000050

To link to this article: <https://doi.org/10.1080/17429145.2021.2000050>



© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



View supplementary material [↗](#)



Published online: 13 Nov 2021.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Role of QseG membrane protein in beneficial enterobacterial interactions with plants and *Mesorhizobia*

Denise Pereira Torres ^{a*}, Ana Paço ^{a*}, Esther Menéndez ^a, Pedro F. Mateos ^b and Clarisse Brígido ^a

^aMED – Mediterranean Institute for Agriculture, Environment and Development, Institute for Advanced Studies and Research, Universidade de Évora, Évora, Portugal; ^bDepartamento de Microbiología y Genética, Centro Hispano Luso de Investigaciones Agrarias, Unidad Asociada CSIC-USAL, Universidad de Salamanca, Salamanca, Spain

ABSTRACT

Homologs of *qseG* gene (coding for the membrane protein QseG), along with the *qseEF* genes, are present in many *Enterobacteriaceae*; however, its role in non-pathogenic strains is still unknown. To fill this knowledge gap, we investigated the role of QseG protein of a plant-associated enterobacterium in the interactions with its legume host and in the benefits induced by this enterobacterium in the *Mesorhizobium*–chickpea symbiosis. Here, we showed that QseG of *Kosakonia* sp. MH5 is involved in the following processes: (i) the evasion of the plant immune system and (ii) the efficient colonization of chickpea root cells. Furthermore, these features are essential for the beneficial effects of this strain on the *Mesorhizobium*–chickpea symbiosis. This study demonstrates that the role of QseG is transversal to pathogenic and non-pathogenic enterobacteria and is a step forward to better understanding the molecular bases of plant–bacteria interactions established between legume and beneficial endophytic enterobacteria.

ARTICLE HISTORY

Received 16 June 2021
Accepted 26 October 2021

KEYWORDS

Plant defence response; bacterial infection ability; plant growth-promoting bacteria; *Cicer arietinum* L; symbiosis; bacterial endophytes

Introduction

Enterobacteriaceae is the largest family of the order *Enterobacterales* and contains many relevant genera and species, including the model animal pathogenic bacteria *Escherichia coli* (Adeolu et al. 2016) and also, species described as plant-associated bacteria, such as the ones within the genera *Enterobacter*, *Phytobacter*, and *Kosakonia* (Brígido, Menéndez et al. 2019; Brígido, Singh et al. 2019; Jha et al. 2011; Madhaiyan et al. 2020), amongst others. However, these genera also harbor strains with pathogenic potential for plants (García-González et al. 2018) and for humans (Mertschnigg et al. 2020; Pillonetto et al. 2018; Salimiyan Rizi et al. 2020), revealing the thin line between pathogenic and mutualistic relationships within this family.

Regardless of the taxonomic and phenotypic diversity within the *Enterobacteriaceae* family, most of our knowledge of the quorum-sensing (QS) systems in this family comes from the *E. coli* model. QS is the ability of bacterial cells to regulate their gene expression according to population growth. This mechanism affects the behavior of bacterial groups, and it is involved in the regulation of traits relevant for the outcome of the interactions between bacteria, either pathogenic or beneficial, and their plant host cells. For example, the QS system of different plant-associated bacteria coordinate biofilm formation, virulence factors expression, and bacteria motility (Kai 2018; Loh et al. 2002; van Loon et al. 2008). QS is subject to regulatory cascades, especially two-component systems that often respond to environmental stimulation. In *E. coli*, the QS regulator proteins E and F (QseEF, also known as GlrKR or YfhKA) are a two-component system involved in the regulation of virulence, metabolism,


and cell envelope homeostasis of this bacteria (Reading et al. 2010; Reading et al. 2007; Reichenbach et al. 2009).

Interestingly, genes coding for the two-component system QseEF are co-transcribed in an operon with QseG (also known as YfhG), an outer membrane protein showing homology with α -helical proteins (Reading et al. 2009; Reading et al. 2007). Recently, it was shown that QseG is indispensable for QseEF activity in *E. coli* K-12. QseE was incompetent to phosphorylate/activate the response regulator QseF in the *E. coli* K-12 Δ qseG mutants (Göpel and Görke 2018). Like the co-transcribed proteins QseE and QseF, QseG also plays a role in regulating the virulence of enteric pathogens (Cameron et al. 2018; Reading et al. 2009; Xiao et al. 2012). For instance, mutation in this gene severely affects the survival of the fish pathogen *Edwardsiella tarda* in murine macrophages as well as its virulence towards zebra fish (Xiao et al. 2012). Similarly, enterohemorrhagic *E. coli* (EHEC) Δ qseG mutants show lower capacity to colonize the rabbit intestinal cells comparing with the wild-type (WT) strain (Cameron et al. 2018). Although QseG is important for the virulence of different pathogenic bacteria, its mechanisms of action differ according to bacteria species. For instance, studies point out a role of QseG in the regulation of the type III secretion system function in EHEC (Cameron et al. 2018; Reading et al. 2009). On the other hand, in *Salmonella enterica* serovar *Typhimurium*, QseG have been associated to flagella phase variation and, consequently, to regulate flagellin recognition. Moreover, its deletion confers less virulence and reduced intestinal colonization in an infant rabbit model than the WT strain (Cameron et al. 2018).

Despite the functional characterization of the *qseEGF* operon in a few genera of intestinal pathogenic bacteria

CONTACT C. Brígido  ccb@uevora.pt  MED – Mediterranean Institute for Agriculture, Environment and Development, Institute for Advanced Studies and Research, Universidade de Évora, Pólo da Mitra, Ap.94, Évora 7006-554, Portugal

*D.P. Torres and A. Paço contributed equally to this work.

 Supplemental data for this article can be accessed <https://doi.org/10.1080/17429145.2021.2000050>.

© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

belonging to the *Enterobacteriaceae* and *Hafniaceae* families (Cameron et al. 2018; Göpel and Görke 2018; Reading et al. 2010; Reading et al. 2009; Xiao et al. 2012), its role in non-pathogenic strains, including plant-associated enterobacteria, whose genome contains putative *qseEGF* operon orthologs (Göpel et al. 2011), is still unknown. Therefore, the characterization of the QseG protein of a plant-associated enterobacterial bacterium presents an opportunity to probe the molecular bases of the plant–bacteria interaction and to better understand the functionality of this protein within *Enterobacteriaceae* family, whose interaction with their hosts varies from pathogenic to beneficial.

In this work, we carried out several experiments to investigate the role of QseG of *Kosakonia* sp. MH5 on the symbiotic *Mesorhizobium ciceri* LMS-1 –chickpea (*Cicer arietinum* L.) association. MH5 strain is an endophytic bacterium isolated from the chickpea roots that exhibits various *in vitro* plant growth-promoting mechanisms (Brígido, Singh et al. 2019) and promotes chickpea growth under salinity when co-inoculated with LMS-1 *in planta* assays (Brígido, Menéndez et al. 2019). Our work presents evidence that the QseG of MH5 strain maintains its function by positively interfering in the host of colonization and infection processes in a similar way to that is observed in *Enterobacteriaceae* pathogens in animals. Furthermore, invasion and a proper establishment within the roots and/or root nodules are essential for this strain to exert beneficial effects on the symbiotic chickpea–*Mesorhizobium* association under salinity.

Materials and methods

Bacterial strains and growth conditions

The bacterial strains and plasmids used in this study are listed in Table S1. The rhizobial strain, *Mesorhizobium ciceri* LMS-1, was grown in tryptone yeast medium (Beringer 1974), at 28°C for routine use as described in (Brígido et al. 2012). The strain *Kosakonia* sp. MH5 and its derivatives, $\Delta qseG$ mutant and complemented strains, were grown in tryptic soy broth (TSB, Liofilchem) or tryptic soy agar (TSA, Liofilchem) medium at 28°C. The *Escherichia coli* strains were grown in Luria–Bertani (LB) broth (Sambrook and Russell 2001), at 37°C. To allow the growth of the conjugation strain *E. coli* WM3064 (Dehio and Meyer 1997), meso-diaminopimelic acid was added to the medium at final concentration of 300 μ M. The biosensor strains *Agrobacterium tumefaciens* NTL4(pZRL4) (Cha et al. 1998) and *Chromobacterium violaceum* CV026 (McClellan et al. 1997) were cultured also at 28°C in AT minimal medium (Morton and Fuqua 2012) and LB medium, respectively. When necessary, 100 μ g/mL ampicillin, 50 μ g/mL kanamycin, 30 μ g/mL gentamycin, 10 μ g/mL tetracycline, or 25 μ g/mL chloramphenicol were added to the medium.

DNA methods and construction of *Kosakonia* sp. MH5 derivatives

The extraction of bacterial genomic DNA was conducted using the E.Z.N.A. bacterial DNA kit (Omega Bio-Tek, Norcross, GA, USA), and the plasmid DNA isolation was performed using the Zippy™ Plasmid Miniprep kit (Zymo Research, Freiburg, Germany). PCR products and gel bands were purified using the kit DNA Clean & Concentrator or Zymoclean™

Gel DNA Recovery kits (Zymo Research, Freiburg, Germany). Unless otherwise specified, molecular techniques were performed using standard protocols (Sambrook and Russell 2001). The primers used in this study are listed in Table S2.

Based on the genome sequence of *Kosakonia* sp. MH5 (accession number ASM993594v1), the flanking regions of the putative *qseG* gene were amplified by following the same approach as described in Paço et al. (2020) using the corresponding primer pairs (Table S2). All PCR reactions were performed in total volume of 50 μ L containing 1 \times buffer, 0.2 mM dNTP mix, 1.5 mM MgCl₂, 15 pmol of each forward and reverse primers, 0.02 U/ μ L of KOD Hot Start DNA Polymerase (Novagen, Winsconsin, USA), and 20 ng of DNA. The PCR programs for amplification the *qseG* flanking regions were as follows: 95°C for 2 min followed by 30 cycles of 95°C for 20 s, 56°C for 10 s, and 70°C for 15 s. To fuse the *qseG* flanking regions, another PCR reaction using a mixture of upstream and downstream PCR products as DNA template and the external primers set (Table S2) was performed under the following program conditions: 95°C for 2 min, 30 cycles of 95°C for 20 s, 68°C for 10 s, and 70°C for 36 s.

The amplified PCR product was isolated and purified from an agarose gel and cloned in pNZY28-blunt vector (NZYtech, Lisbon, Portugal). The fused-flanking region of the *qseG* gene was then subcloned into the EcoRV sites of the suicide plasmid pNPTS138-R6KT (Lassak et al. 2010) and transformed into *E. coli* DH5 α -pir competent cells. Then, the plasmids were purified from positive clones and cloned into *E. coli* WM3064. The pNPTS138-R6KT- $\Delta qseG$ was mobilized into the MH5 cells by biparental mating. Double recombinants were selected as previously described (Ekandjo et al. 2017). Gene knockout was confirmed by PCR in kanamycin-sensitive colonies using the flanking primers, as well as an internal primer pair (Table S2).

To complement the in-frame deletion of the *qseG* gene, replacement of the deletion *qseG* gene by the complete *qseG* gene in its original genomic location was performed. For this, a DNA fragment, containing the complete *qseG* gene and part of its flanking regions, was amplified from the genomic DNA of MH5 WT strain using the flanking primers pair (Table S2). The PCR program was as follows: 95°C for 2 min followed by 30 cycles of 95°C for 20 s, 58°C for 10 s, and 70°C for 50 s. The PCR product was purified and cloned into pNZY28, treated with the EcoRI and HindIII and subcloned into the suicide plasmid pNPTS138-R6KT. The resulting recombinant suicide plasmid was mobilized into the MH5 cells by biparental mating and the in-frame deletion in the $\Delta qseG$ mutant strain was replaced by the WT copy of the gene by homologous recombination as described previously. The positive complemented strain was hereafter denominated as $\Delta qseG$ -KI.

Fluorescently tagged strains were generated by introducing the plasmid pHc60 (GFP) (Cheng and Walker 1998) into $\Delta qseG$ mutant strain and the plasmid pBHR-mRFP (RFP) (Smit et al. 2005) into MH5 WT strain by biparental or triparental mating, respectively, as earlier described (Nascimento et al. 2012; Smit et al. 2005).

In vitro phenotypic characterization of the $\Delta qseG$ mutant strain

Acyl-homoserine lactones production

To determine the range of acyl-homoserine lactones (AHLs) produced by MH5 WT and its derivatives, two biosensor

strains able to detect long or short chain AHL were used. The transformed biosensor strain *A. tumefaciens* NTL4(pZRL4) was used for long side-chain AHLs (C8-HSL-C14-HSL) detection while strain *C. violaceum* CV026 was used for short side-chain AHLs detection (C4-HSL-C8-HSL). The reporter strains and the MH5 WT and derivatives strains were grown overnight at 28°C in their respective culture medium. For short chain AHL assay, 50 µL of the cell suspension containing the endophytic bacterium ($OD_{600nm} = 1$) were spread onto LB agar plates previously supplemented with 50 µg/mL 5-bromo-4-chloro-3-indolyl-β-D-galactoside (X-Gal, Sigma-Aldrich). The plates were dried, inoculated with four droplets of 3 µL of overnight culture of NTL4 (pZRL4) and incubated for 48 h at 28°C. For long chain AHL assay, each bacterial endophyte was streaked approximately 1 cm apart from the biosensor strain CV026 onto LB agar plates and incubated at 28°C for 96 h.

Swarming motility assay

To evaluate motility of the $\Delta qseG$ mutant strain and compared it with the WT strain, the swarming motility of each strain was evaluated in plates containing TSB supplemented with 0.5% (w/v) agar. Five microliters of a bacterial suspension grown overnight in TSB medium with an OD_{590nm} of 2 were spotted in the center of the plates. Plates were incubated at 28°C for 14 days, and differences in motility were determined by the rate of radial expansion from the inoculation spot. Five replicas per treatment were performed.

Biofilm formation

Biofilm formation on microtiter PVC plates was performed according to (O'Toole 2011), with few modifications. Briefly, strains were grown in TSB medium at 28°C. After overnight incubation, the OD_{600nm} of the cultures were adjusted to 0.1 in M9 broth medium (Duan et al. 2009). One hundred microliters of cells suspensions were pipetted into individual wells of a 96-well PVC plate (Corning Costar 2797, Massachusetts, USA). The plates were sealed and incubated at 28°C without agitation for 8 h. Cell growth was measured in a BioRad Microtiter Plate reader (Model 680) at 565 nm. Then, the media were removed, and surface-attached bacteria biomass was quantified by staining them with 125 µL of 0.1% (w/v) solution of crystal violet in water. After incubation for 10–15 min, the wells were gently rinsed with distilled water and dried. The remaining dye was solubilized with 80% ethanol: 20% acetone (v/v) solution and quantified by measuring the absorbance at 565 nm. Biofilm formation was normalized by cell growth. For each treatment, 8 wells of the 96-well PVC plate were measured.

The biofilm formation was also evaluated on sand (Fujishige et al. 2006). After overnight growth in TSB medium at 28°C, the OD_{590nm} were adjusted to 0.4 and 500 µL of the bacterial solution or the same volume of TSB medium were inoculated in 1.25 mL tubes containing 600 mg of sterile sand. After 72 h of incubation at 28°C with 200 rpm agitation, the media were removed, and the sand washed once with 1× phosphate-buffered saline (PBS). The cells were finally suspended in 1 mL of 1× PBS and the tubes were vortexed intensively to release attached cells. Cells were quantified by flow cytometry using a CyFlow Cube8 (Sysmex Partec, Görlitz, Germany).

Tolerance to salinity and manganese toxicity

To evaluate the $\Delta qseG$ mutant strain tolerance to salinity and manganese and compared it with the WT strain, bacterial growth was determined by measuring the OD_{565nm} during 28 h of growth in the liquid medium. For all treatments, bacterial cultures with an initial OD_{565nm} of 0.1 were prepared after an overnight growth in TSB. After growth, MH5 WT and derivative strains were grown in M9 broth medium supplemented with 1% (w/v) NaCl or 1 mM manganese as stress conditions and grown in regular M9 broth medium as control condition. Three replicas per treatment were performed.

Effect of the $\Delta qseG$ mutant strain on the mesorhizobia-legume symbiosis

Pot experiment setup

To evaluate the effects the $\Delta qseG$ mutant strain on the rhizobium-legume symbiosis, a plant growth assay was conducted using the symbiotic chickpea-*Mesorhizobium* model under salinity conditions in a growth chamber.

The well-characterized *M. ciceri* LMS-1 was used as rhizobial microsymbiont and was prepared as previously described by Brígido et al. (2013) while the MH5 WT and derivatives were prepared as described by Brígido, Menéndez et al. (2019).

Surface sterilization and pre-germination of the chickpea seeds were carried out as earlier described (Paço et al. 2016). The setup of the pots trial was performed as following a similar protocol to that described by Brígido, Menéndez et al. (2019) with the exception that the bacterial suspensions used to inoculate each germinated seedling contained a final OD_{565nm} of 1 of each bacterium. A nitrogen-free nutrient solution (Broughton and Dilworth 1971) was used to water the plants three times a week. To induce salinity stress, 0.075% of NaCl was added to nitrogen-free nutrient solution after the second week of plants growth until the end of the experiment. Four treatment combinations were prepared as follows: (i) chickpea single inoculated with LMS-1; (ii) chickpea co-inoculated with LMS-1 and MH5 WT; (iii) chickpea co-inoculated with LMS-1 and $\Delta qseG$ mutant strain; and (iv) chickpea co-inoculated with LMS-1 and $\Delta qseG$ -KI strain. Five pots per treatment were used. The plants were harvested after 6 weeks, and the number of nodules, nodules dry weight, root dry weight, and shoot dry weight (SDW) were registered.

Nodule histology

Chickpea nodules were excised from roots of 6-week-old plants and processed for microscopy as previously described (da-Silva et al. 2019). A pool of nodules from at least three representative plants from each treatment (chickpea single inoculated with LMS-1 or co-inoculated with LMS-1 and MH5 WT or co-inoculated with LMS-1 and $\Delta qseG$ mutant strains) were processed for microtome sectioning. Toluidine blue-stained cross sections of nodules were examined by brightfield microscopy under a Leica DM6000B microscope.

Gene expression analyses by qRT-PCR

RNA extraction

Expression levels of specific genes in the MH5 WT and $\Delta qseG$ mutant strain were evaluated during the early stages of interaction with the legume host. The strains were

grown in a mixture of TSB medium and chickpea root exudates (1:1, V/V) at 28°C for 24 h. Chickpea root exudates were previously obtained as described by Srivastava et al. (1999). To ensure a complete break of bacterial cell walls, the cells were mechanically lysed, and the total bacterial RNA was extracted using the Absolutely RNA Miniprep Kit (Agilent Technologies, Santa Clara, USA). Three biological replicates of each strain were used.

To evaluate changes in chickpea response to inoculation with MH5 WT or $\Delta qseG$ mutant strain, a plant growth assay was performed as described previously. Chickpea seedlings were co-inoculated with LMS-1 and MH5 WT or co-inoculated with LMS-1 and $\Delta qseG$ mutant strain. Chickpea seedlings single inoculated with LMS-1 were used as control. The expression of plant defense-response-related genes in roots was assessed 24 h after bacterial inoculation. The entire root system was gently and thoroughly washed with distilled water. Then, roots were separated from stem, frozen in liquid nitrogen, and ground to a fine powder using mortar and pestle and stored at -80°C for further analysis.

Three seedlings were pooled per replicate and three biological replicates were used per treatment. Total RNA of chickpea seedlings was extracted using the kit Maxwell® 16 LEV simplyRNA Tissue Kit (Promega, Madison, USA).

RNA analysis and cDNA synthesis

The concentration and purity of the RNA extracted from both bacteria and plants were evaluated using a NanoDrop 800 UV-Vis spectrophotometer (Thermo Fisher Scientific, Waltham, USA). The RNA integrity was checked on 1% agarose gel electrophoresis. Approximately, 2 µg of total RNA from bacteria and 1 µg of total RNA from chickpea seedlings were subjected to reverse transcription for cDNA synthesis using the NZY First-strand cDNA synthesis kit (NZYtech, Lisbon, Portugal).

qRT-PCR

To evaluate changes in the expression levels of genes related to plant-microbe interactions, the expression of *ompA*, *tagJ*, and *virK* genes were evaluated in the MH5 WT and $\Delta qseG$ mutant strains after exposition to chickpea root exudates. On the other hand, gene expression profile of genes related to plant defense-response, namely *CaRBOH*-like, *CaPR1*, and *CaFLS2*, were assessed in chickpea seedlings co-inoculated with LMS-1 and either with MH5 WT or $\Delta qseG$ mutant strain. The primers for these target genes were designed using the Primer-Blast NCBI tool (Ye et al. 2012). The 16S rRNA gene was used to normalize the expression of the bacterial target genes while *Ca18SrRNA* and *CaGAPDH* genes were used to normalize the expression of the chickpea target genes. All primers used herein are listed in Table S2. Primer pairs were tested on cDNA samples to ensure amplification of the target region. The specificity of amplification reactions was verified by a melting curve analysis performed at 60–95°C. qRT-PCR was conducted in a 96-well plate format on a 7500 real-time PCR detection system (Applied Biosystems) using the Green Master Mix (2×) ROX kit (Nzytech). The cDNA used was diluted 5× with Dnase/Rnase free water and used 800 ng in a reaction volume of 20 µL and performed as outlined by the manufacturer (Applied Biosystems). For each gene, three technical replicates of three biological replicates were performed. No template controls to detect DNA contamination or primer dimers were also

performed. Expression levels for the target were determined by the $2^{-\Delta Ct}$ method, where ΔCt is the difference between the Ct of a target gene and the Ct of the reference gene, and transcript levels were calculated and normalized according to Willems et al. (2008).

Evaluation of root colonization ability by MH5 WT and $\Delta qseG$ mutant strain

Colony-forming unit quantification of bacterial cells in different rhizocompartments

The standard dilution plating method was used to evaluate the ability of root colonization by MH5 WT and $\Delta qseG$ mutant strain. For that purpose, a plant growth assay with chickpea seedlings co-inoculated with LMS-1 and MH5 WT or $\Delta qseG$ mutant strain was performed as described previously. The roots sampling was carried out at 24, 48, and 72 h after seedlings bacterial inoculation. After thoroughly washing the roots with sterile water, they were cut transversally into two sections of equal size and the seed-proximal section was kept. For each treatment, roots from a pool of three plants were harvested. To obtain the bacteria attached to rhizoplane, the root sections were submerged in 3X Ringer's solution (Surette et al. 2003) and vortexed for 2 min at max velocity, and then bacterial cells were recovered by centrifugation and then suspended in 3X Ringer's solution. To obtain the bacteria within root cells, the root sections were surface-disinfected as described in Rashid et al. (2012) and crushed with a sterile pestle in 3X Ringer's solution. Serial dilutions of the bacteria suspension were prepared and 100 µL of each dilution was plated in TSA plates supplemented with 20 µg/mL of ampicillin. Plates were incubated at 28°C for 48 h. Three biological replicates with three plants each were performed.

Fluorescence microscopy

To visualize chickpea root colonization, RFP-tagged MH5 WT and GFP-tagged $\Delta qseG$ mutant strains were used to inoculated surface-disinfected and pregerminated chickpea seeds. Briefly, surface-disinfected and pregerminated chickpea seedlings were transferred to a 0.75% water-agar plate (square petri dish, 140 × 140 mm) and inoculated with 400 µL/plant of each of the bacterial suspensions at OD₆₀₀ of 0.4 prepared as described above. Plants were overlaid with double filter paper and incubated in a growth chamber for 5 days. Then, plant roots were excised and carefully removed from the plates to a microscope slide. Roots were visualized using a confocal laser scanning microscope (Leica TCS SPE). For each treatment, at least 2 plates of 3 seedlings/each were observed plus an uninoculated control (chickpea seeds without bacterial inoculum).

Bioinformatics analyses

Analyses of the gene synteny of the *qseEGF* region in *Enterobacterales* were performed using the MicrobesOnline tool (Dehal et al. 2010) accessed in November 2020 and NCBI database. For this, sequences of one representative species of each of the seven *Enterobacterales* families (Adeolu et al. 2016), whose genome sequences were available in MicrobesOnline website, were chosen and compared with MH5 WT. The representative species were as follows: *E. coli* str. K-12 substr. MG1655 (*Enterobacteriaceae*), *Erwinia*

amylovora CFBP1430 (*Erwiniaceae*), *Pectobacterium carotovorum* subsp. *carotovorum* PC1 (*Pectobacteriaceae*), *Yersinia enterocolitica* subsp. *enterocolitica* 8081 (*Yersiniaceae*), *Edwardsiella tarda* EIB202 (*Hafniaceae*), *Proteus mirabilis* HI4320 (*Morganellaceae*), and *Pragia* sp. CF-458 (*Budviciaceae*).

Statistical analysis

Data were analyzed using either Student's t-test or one-way ANOVA. The Duncan test was used to detect significant differences between the treatments means ($P < 0.05$). Statistical analysis was carried out using SPSS V.21 software (SPSS Inc.).

Results

qseEGF operon is conserved in Enterobacterales

The *qseEGF* operon is annotated in the genome of several enterobacterial species (Göpel et al. 2011) and it is relatively well-characterized in a few enteric pathogens. To check whether the gene architecture of this operon is conserved in *Enterobacterales*, we analyzed the gene synteny of *qseEGF* region in representative species of the seven families of this order. The genomic analysis revealed that *qseEGF* operon was conserved in this order (Figure S1). The putative *qseEGF* operon (Locus tag: DK872_RS06570-RS06580) of MH5 WT was localized downstream the small RNA *GlmY* (NZ_QODH01000002.1: 32386-32570) and upstream to the gene *glnB* (WP_002438074.1), which encodes a nitrogen regulatory protein P-II (Figure 1(A)). The amino acid identities of the QseE, QseG, and QseF proteins of MH5 WT are of 85%, 58%, and 94% to the respective orthologues in *E. coli*. The putative *qseG* gene (DK872_RS06575) presented a sequence of 738 bp, which likely encodes a protein of 245-amino acid peptides with a predicted molecular weight of 27.9 kDa. An alignment of its predicted amino acid sequence with sequences of characterized proteins from *Enterobacterales* confirmed the identity of QseG in MH5 WT (Figure 1(B)).

In vitro phenotypic characteristics of $\Delta qseG$ mutant strain are not affected by deletion of the *qseG* gene

To characterize the QseG protein of a plant-associated endophytic enterobacterium, a QseG knockout mutant of the MH5 strain was constructed by in-frame deletion of an internal fragment of 645 bp of its *qseG* gene sequence. The $\Delta qseG$ mutant strain genotype was restored when the in-frame *qseG* gene deletion was successfully replaced by the complete *qseG* gene sequence in its original genomic location.

AHLs production in the MH5 WT and $\Delta qseG$ mutant strains was detected using the biosensors strains *A. tumefaciens* NTL4 (pZLR4) and *C. violaceum* CV026. Both MH5 WT and $\Delta qseG$ mutant strains were able to activate the biosensor strain *Agrobacterium tumefaciens* NTL4 (pZLR4). On the other hand, none of the tested enterobacterial showed positive results for the *C. violaceum* CV026 bioassay, indicating that MH5 WT is unable to synthesize AHL-type molecules with short acyl chains. Thus, the ability of the MH5 WT to synthesize AHL with medium to long

fatty-acid chains did not require the action of the QseG protein. Similarly, neither biofilm formation on abiotic surfaces such as microtiter PVC plates (Figure S2A) or sand (Figure S2B) nor swarming motility (Figure S2C) were affected by the *qseG* gene deletion in the $\Delta qseG$ mutant strain.

Our previous results showed that *Kosakonia* sp. MH5 is tolerant to high salt concentrations (5% NaCl) and to manganese (1 mM Mn) (Brígido, Menéndez et al. 2019). To investigate whether QseG plays a role in responding to these abiotic stresses, the growth of the MH5 derivative strains was analyzed in M9 broth medium supplemented with NaCl or manganese. There were no significant differences between the growth curves of the MH5 WT, $\Delta qseG$ mutant and $\Delta qseG$ -KI strains in any of the conditions tested (Figure S2D).

QseG regulates the expression of genes related to plant-microbe interactions

Considering that QseG is implicated in the regulation of the expression of genes critical for the interactions between enterobacterial species and animal hosts (Cameron et al. 2018; Reading et al. 2009; Xiao et al. 2012), we hypothesized that QseG of plant-associated enterobacterial species also plays a role in the evasion of host immune system. To test this hypothesis, the expression levels of homologues of genes well-known for their roles in virulence of pathogenic bacteria were studied in MH5 WT and $\Delta qseG$ mutant strains upon exposition to chickpea root exudates for 24 h. Three genes were selected: the *ompA* (DK872_RS02445), which codes for the outer membrane protein A (OmpA), and contributes to bacterial adherence, biofilm formation, and host defense evasion (Smith et al. 2007); the *virK* (DK872_RS21215) and *tagJ* (also known as *impE*; DK872_RS20860) code for effector proteins secreted for the type II and VI secretion system (T6SS), respectively (Förster et al. 2014; Yamazaki et al. 2008).

Although there was a trend towards a downregulation of the expression levels of *virK* in the $\Delta qseG$ mutant strain, only the expression of *ompA* and *tagJ* genes were statistically lower than the MH5 WT strain (Figure 2). In addition, the degree of downregulation was stronger for the effector protein TagJ, whose expression levels were less than 0.5-fold changes of the expression detected in the WT strain.

QseG plays an essential role in Mesorhizobium-chickpea-Enterobacteriaceae interactions

Since our previous results showed that MH5 WT was able to enhance the symbiotic LMS-1-chickpea association under salinity (Brígido, Menéndez et al. 2019), the involvement of QseG protein in the beneficial effect promoted by MH5 WT in this tripartite association was investigated.

Although the nodulation ability of LMS-1 microsymbiont was not statistically affected by the presence of any of the MH5 derivative strains (Figure 3(A)), the nodule dry weight was significantly higher in plants co-inoculated with LMS-1 and any of the strains containing a functional QseG protein comparing to the remaining plants. Consequently, plant total biomass was 1.6 times higher in plants co-inoculated with the rhizobia and MH5 strains with a complete *qseG* gene (Figure 3(B)). In contrast, the increase in nodule dry weight




Figure 1. The *qseEGF* operon is conserved in *Enterobacteriales*. (A) Schematic representation of the gene synteny of *QseEGF* operon. Primers used for the knockout are shown. Genes are just approximately drawn to scale. (B) Alignment of *QseG* protein amino acid sequences from bacteria belonging to the order *Enterobacteriales*. Sequences were retrieved from NCBI and belong to the following genomes (accession numbers are in parentheses): *Kosakonia radicincitans* DSM 16656 (ARD59616.1), *E. coli* O157:H7 (EFD2029335.1), *Edwardsiella tarda* EIB202 (ACY83621.1), *Salmonella enterica* subsp. *enterica* serovar Typhimurium str. SL1344 (HAD6495497.1), and *Citrobacter rodentium* (WP_012906675.1).

was counteracted when the chickpea plants were co-inoculated with the *Mesorhizobium* strain and the $\Delta qseG$ mutant strain. Altogether, these results indicate that the benefits of MH5 WT in the symbiotic *Mesorhizobium*–chickpea association were dependent of a functional *QseG* protein.

To determine whether the histology of the nodules formed by LMS-1 was affected by the presence of MH5 WT or $\Delta qseG$ mutant strains, we analyzed histological cross sections of the nodules of chickpea plants single inoculated with LMS-1 or co-inoculated with LMS-1 and MH5 WT or $\Delta qseG$ mutant strains. The bright field light microscopy (Figure 3(C–H)) showed that nodules of all

treatments presented a typical histology of indeterminate effective nodules, having differentiated meristematic, infection, and bacteroid zones. Despite the difference in nodules size, no differences on nodule histology were observed among the different treatments.

QseG plays a role in root infection and colonization

Expression of genes related to plant defense evasion is reduced in the $\Delta qseG$ mutant strain, suggesting that the knockout mutant strain elicits broader plant defense responses than the MH5 WT strain. To test this hypothesis,




Figure 2. QseG plays a role in plant–microbe interactions. Expression levels of *ompA* and the secretion system proteins *virK* and *TagJ* genes in MH5 WT (MH5) and $\Delta qseG$ mutant ($\Delta qseG$) after exposure to root exudates for 24 h. Relative gene expression was normalized against 16S rRNA and compared to MH5 WT as control. Each point represents the mean and standard error of the mean of three biological replicates. Asterisks indicate significant differences between the genotypes (Student's *t* test: ** $p < 0.01$, *** $p < 0.001$).

we evaluate changes in chickpea response to co-inoculation with LMS-1 and MH5 WT or $\Delta qseG$ mutant strain, during the early stages of plant–bacteria interactions. Chickpea seedlings single inoculated with LMS-1 were used as control. We studied the expression of a putative respiratory burst oxidase homolog, RBOH (*CaRBOH*-like; XP_004494602.1), related to the production of reactive oxygen species (ROS) in response to environmental constraints, and a pathogenesis-related protein (PR) 1 (*CaPR1*; XP_004485656.1). Additionally, the expression of a putative chickpea flagellin receptor, flagellin sensing (FLS) 2 (*CaFLS2*; XP_012572420.1). Expression levels of *CaFLS2* in the roots were upregulated in plants co-inoculated with the *Mesorhizobium* and any of enterobacterial strains compared with single-inoculated plants at 24 h after inoculation (Figure 4(A)). Even though *CaFLS2* expression was 1.3-fold changes higher in plants co-inoculated with the MH5 WT than in plants co-inoculated with the $\Delta qseG$ mutant strain. Similarly, *CaRBOH*-like expression was differentially induced in co-inoculated plants, being its expression much higher in plants co-inoculated with LMS-1 and $\Delta qseG$ mutant. Hence, the $\Delta qseG$ mutant strain triggered plant defense responses in chickpea roots 3.9-fold changes stronger than the MH5 WT strain. Curiously, compared with roots of single-inoculated plants, *CaPR1* expression was only significantly upregulated in plants inoculated with the MH5 WT.

Following, we evaluated if the capacity for colonization and infection of chickpea roots by the enterobacterial endophyte were altered by the deletion of the *qseG* gene sequence. For that purpose, two different approaches were performed: (i) direct observation of chickpea root colonization under confocal microscope after the inoculation of chickpea seedlings with the RFP-tagged MH5 WT or GFP-tagged $\Delta qseG$ mutant strain or the mixture of both at 5 days post-inoculation and (ii) analyzing colonization and infection on roots by counting the number of viable enterobacterial cells on the surface or within root cells at 24, 48 and 72 h post-inoculation. The first approach revealed that both MH5 WT and $\Delta qseG$ mutant strains were able to effectively colonize chickpea roots (Figure 4(B–D)). When inoculated together, the MH5 WT showed to be more competitive than the $\Delta qseG$ mutant strain. The results of the second approach, which include the strain LMS-1 as compatible

chickpea mesorhizobia, suggested a delay in the proliferation of $\Delta qseG$ mutant strain on the roots at 24 h post-inoculation (Figure 4(E)). In this time point, the number of $\Delta qseG$ cells attached to the chickpea roots was three times higher than with the MH5 WT. But the number of attached bacteria did not differ significantly between these two strains at 48 h post-inoculation. Interestingly, internal root tissue colonization by MH5 WT was higher than by the $\Delta qseG$ mutant strain at all tested time points (Figure 4(F)).

To check if internal root colonization by the $\Delta qseG$ strain was compromised even at later points of this multipartite interaction, a hydroponic assay with chickpea plants grown under salinity stress conditions was performed. The results showed that the growth of chickpea plants co-inoculated with LMS-1 and $\Delta qseG$ were restricted (Figure 4(G)) and consistent with the results from the pot experiment. The colony-forming unit (CFU) counting showed that MH5 WT population inside the chickpea roots was 10-fold superior to the $\Delta qseG$ mutant strain at 28 days post-inoculation (Figure 4(H)). Together, these results show that the internal root infection and colonization ability of the MH5 WT was compromised by the *qseG* gene deletion, indicating that QseG influenced the onset of plant–bacteria associations, especially the endophytic colonization, nevertheless are not very important for epiphytic growth.

Discussion

We observed the involvement of QseG protein in the early stages of the plant–endophytic bacteria interactions, particularly in the internal root tissue colonization and plant's immune system evasion. The involvement of QseG of the endophytic enterobacterium *Kosakonia* sp. MH5 strain in the regulation of expression genes related to host defense evasion is supported by (i) the downregulation of the expression of *ompA* and *tagJ* genes in the $\Delta qseG$ mutant strain and (ii) the defective colonization of the internal root cells from an early to a later stage of the plant– $\Delta qseG$ mutant strain interaction. *OmpA* is a virulence gene that seems to be crucial to mitigation of host defenses by animal pathogens (Confer and Ayalew 2013; Smith et al. 2007) and to the motility and efficient plant root colonization (Cole et al. 2017). *TagJ* (type VI accessory genes J) is an accessory protein present in sub-families 3 and 5 of type VI secretion systems, T6SSs (Boyer et al. 2009), which, in plant-associated bacteria is involved in plant manipulation and interbacterial competition (Bernal et al. 2018). These observations agree with the fact that beneficial microbes are initially recognized as potential invaders, but they are able to short-circuit plant defense responses to enable successful colonization of host roots (Zamioudis and Pieterse 2012). This is particularly apparent for endophytes living within the plant and, therefore, are directly exposed to the host's immune system (Liu et al. 2017). Therefore, it seems plausible that plant-beneficial bacteria, as MH5 strain, need to escape root immune responses in order to establish an effective colonization (Yu et al. 2019).

With the $\Delta qseG$ mutant strain's ability to evade the plant's defense reduced, it is expected that chickpea inoculation with this strain elicits broader plant defense responses than the MH5 WT strain. Consistent with this hypothesis, *CaRBOH*-like expression was higher in chickpea roots inoculated with MH5 $\Delta qseG$ mutant strain. RBOH's is among the major




Figure 3. QseG is required for benefits of *Kosakonia* sp. MH5 in the symbiotic chickpea–*Mesorhizobium* relationship under salinity (0.075% NaCl). Results from 54 days plant growth assay with chickpea plants single inoculated with *M. ciceri* LMS-1 (LMS-1) or co-inoculated with LMS-1 and MH5 WT (LMS-1 + MH5) or ΔqseG (LMS-1 + ΔqseG) or MH5 ΔqseG KI (LMS-1 + KI). (A) Nodule dry weight in bars and number of nodules in line. (B) Root, shoot, and total biomass dry weight. Error bars show standard error of the mean ($n = 5$ plants) and letters the grouping of data sets according to Duncan's test at $P \leq 0.05$. Uppercase letters show the grouping of data sets relative to nodule and root dry weight, while lowercase data sets grouping relative to number of nodules and shoot dry weight uppercase. Underline lowercase letters show the grouping of data sets relative to total biomass dry weight. Histological cross sections (6 μm) of chickpea nodules formed in single inoculated plants (C, F), co-inoculated with LMS-1 + MH5 (D, G), and co-inoculation of LMS-1 + ΔqseG (E, H). Scale bars are 1mm (A, B, C) and 400μm (D, E, F).

class of enzymes involved in the oxidative burst, one of the earliest events following the recognition of microbes by the plant immune system (Torres and Dangl 2005). Transcriptional upregulation of RBOH's genes in response to the presence of phytopathogens and beneficial bacteria is described in several plants, including *Arabidopsis*, barley, maize, and chickpea (Arthikala et al. 2017; Chen et al. 2017; do Amaral et al. 2014; Gupta et al. 2013; Morales et al. 2016; Torres et al. 2017). Moreover, *CaRBOH*-like expression was higher in roots of double-inoculated plants. This result is congruent with studies that suggest rhizobia have evolved to reduce or actively suppress host defence responses (Zamioudis and Pieterse 2012). For instance, inoculation of *Ensifer*

(*Sinorhizobium*) *meliloti* Sm2011 does not trigger expression of *MtFLS2*, while inoculation of the phytopathogen *Pseudomonas syringae* pv. *Tomato* DC3000 or co-inoculation of both bacteria does (Chen et al. 2017).

Lower expression of the salicylic acid (SA)-dependent marker gene *CaPRI* (Breen et al. 2017) in plants co-inoculated with the *Mesorhizobium* and ΔqseG mutant strain may suggest a role of QseG in inducing of disease resistance, but it can also be the result of its inability to activate PR expression for not reaching the required population density within plant cells. Either way, the induction of disease resistance by triggering the expression of PR genes, as observed by some plant growth-promoting bacteria, is reduced in plants




Figure 4. QseG is involved in the processes (i) to evade host immune system and (ii) to efficiently colonize the chickpea root cells. (A) Expression of defense-related genes in roots of chickpea plants single inoculated with *M. ciceri* LMS-1 (LMS-1) or co-inoculated with LMS-1 and MH5 WT (LMS-1 + MH5) or $\Delta qseG$ (LMS-1 + $\Delta qseG$). Relative gene expression was normalized against *Ca18S*rRNA and *CaGAPDH* genes and compared with the expression in single inoculated plants as control. Each point represents the mean and standard deviation of three biological replicates. Different letters correspond to statistically significant differences ($P < 0.05$). (E) Colonization of the surface and (F) internal cells of chickpea roots by the MH5 WT (LMS-1 + MH5) and $\Delta qseG$ (LMS-1 + $\Delta qseG$) in the early stages of multipartite *Kosakonia*–chickpea–*Mesorhizobium* interaction. Plants were grown in vermiculite pots and roots were sampled 24, 48, and 72 h after bacterial inoculation. Each point represents the mean and standard error of the mean of three biological replicates. (G) Shoot dry weight of inoculated chickpea plants grown under salinity in a hydroponic system for 28 days after inoculation. Each point represents the mean and standard error of the mean of 7–8 plants. Different letters correspond to statistically significant differences ($P < 0.05$). (H) Colonization of chickpea roots by the MH5 WT (LMS-1 + MH5) and $\Delta qseG$ (LMS-1 + $\Delta qseG$) in the late stage of multipartite *Kosakonia*–chickpea–*Mesorhizobium* interaction. Each point represents the mean and standard error of the mean of three plants. Asterisks indicate significant differences between the genotypes (Student's *t* test: ** $p < 0.01$).

inoculated with the $\Delta qseG$ mutant. For instance, inoculation of tobacco plants with beneficial *Pseudomonas* spp. strains triggers the expression of *PR3* and induces systemic resistance against the bacterial pathogen *Erwinia carotovora* (van Loon et al. 2008). High *PR* gene expression levels were also associated with induction of resistance against the fungal pathogen *Verticillium dahliae* in *Arabidopsis* plants inoculated with *Paenibacillus alvei* K165 (Tjamos et al. 2005). Similarly, inoculation of *Arabidopsis* plants with *Kosakonia radicincitans* DSM 16656 leads to activation of SA and jasmonate (JA)/ethylene (ET) defense-related genes and increased resistance to chewing and phloem-feeding insects (Brock et al. 2013; Brock et al. 2018). Thus, it can be speculated that priming for chickpea enhanced defense is improved by MH5 WT inoculation. On the other hand, induction of systemic resistance is dependent of the initial

population density of the plant growth-promoting bacteria as was observed in radish (Raaijmakers et al. 1995). For instance, *Gluconacetobacter diazotrophicus* ROS-detoxifying mutants, which are also deficient in root colonization, fail to induce *PR10* expression in rice (Alquéres et al. 2013; Raaijmakers et al. 1995). Nevertheless, induction of resistance in chickpea by this beneficial endophytic bacterium needs to be further investigated.

The hypothesis that QseG is important for the host plant–bacteria interaction is also strengthened by the observation that, despite co-inoculated chickpea seedlings displayed the highest transcript levels of the putative *CaFLS*, its expression decreased in plants inoculated with the $\Delta qseG$ mutant strain. *FLS2* is one of the best characterized plant recognition receptor of the model plant *Arabidopsis thaliana* (Zipfel and Oldroyd 2017). Although *FLS2* orthologue in chickpea was not

yet characterized, its sequence shows 82.5% of identity to MtFLS2 (XP_013457339.1) and conserved domains. FLS2 recognize the flagellin epitope flg22, a highly conserved part of the N terminus of bacterial flagella (Boller and Felix 2009). *Kosakonia* sp. MH5 encodes two flagellin homologues with identical flg22 motifs and 100% identity to flg22 of *Salmonella enterica* (Figure S3), which has recently been shown to triggers FLS2-dependent defence responses in alfalfa (Wahlig et al. 2019). Therefore, the mutation in the *qseG* gene may alter the flagellin recognition as earlier observed in *S. enterica* serovar Typhimurium (Cameron et al. 2018). Despite that flagellar phase variation was not before reported in *Kosakonia* spp., a study revealed that flagellin glycosylation islands are common in the genome of bacteria belonging to this genus (De Maayer and Cowan 2016). Flagellin post-transcriptional modifications may interfere with the recognition of the immunogenic epitopes of the flagellin protein and as consequence helps to evade host defence responses (Verma et al. 2005). Although this is not consistent with a reduced chickpea defence response, the intensity of the immune responses does not always correlate with the expression level of the FLS2 receptor in the root system but rather depends on the expressing tissue (Wyrsh et al. 2015). The FLS2 promoter activity is mainly present in the *Arabidopsis* root stele and expands to the cortex and epidermal region (Beck et al. 2014; Wyrsh et al. 2015). From this perspective, lower levels of expression of FL2 in chickpea roots inoculated with $\Delta qseG$ mutant may be due to its inability to reach those root tissues that priming to a proper balance of defense responses according to the expected exposure to elicitors.

Although the importance of a functional QseG for the benefits of the MH5 in the chickpea–*Mesorhizobium* symbiosis is evident, we do not fully identify the mechanisms of action behind the benefits of this enterobacterium. Looking to the gene synteny of the *qseEGF* operon region of MH5 strain, its genomic context is typical of bacteria assigned to the families *Enterobacteriaceae*, *Erwiniaceae*, and *Pectobacteriaceae* (Göpel et al. 2011). In these families, the gene encoding the PII protein GlnB is localized just downstream of *qseEGF* genes while in *Yersiniaceae*, *Hafniaceae*, *Morganellaceae*, and *Budviciaceae*, the gene *nadE* (NH₃-dependent NAD⁺ synthetase) is inserted between *qseF* and *glnB* (Figure S1). In *E. coli*, the genes encoding QseEGF and GlnB are co-transcribed in an operon. PII proteins are widely distributed small homotrimeric signal transduction proteins that bind to different targets, including NifA in nitrogen fixing bacteria (Sotomaior et al. 2012). It was shown that *Azospirillum brasilense* NifA is only active in the presence of GlnB (Arsene et al. 1996) and the uridylylated form of GlnB is essential for NifA activity, and potentially for interaction of GlnB with the N terminal domain of NifA (Inaba et al. 2015). Curiously, transcription of *glnB* increases in *E. coli* *qseG* mutant, suggesting that QseG act through the promoter upstream of *glnB* to repress its transcription (Reading et al. 2009). It can be speculated that *qseG* mutant changes the transcription of *glnB*, which indirectly may have altered the activation of NifA in the nitrogen fixing MH5 strain. Nevertheless, further characterization of the $\Delta qseG$ mutant through transcriptomic analysis and analysis of the QseG protein-dependent changes in the plant transcriptome will be important future steps toward determining the role QseG membrane protein and their targets play in bacteria–plant interactions.

Conclusion

Our work shows that QseG of *Kosakonia* sp. MH5 is involved in the following processes: (i) the evasion of the plant immune system and (ii) the efficient colonization of chickpea root cells. Furthermore, the invasion and a proper establishment within the roots and/or root nodules are essential for this strain to be able to exert beneficial effects on the symbiotic chickpea–*Mesorhizobium* association under salinity. Our work present evidence that the QseG of an endophytic enterobacterium associated with plants maintains its function by positively interfering in the process of colonization and infection of the host similarly to what is observed in *Enterobacteriaceae* pathogens in animals. This corresponds, as far as we know, to the first study regarding the role of QseG protein of an endophytic enterobacterium in the plant–enterobacterium interaction showing that its role is similar and transversal to enterobacteria regardless their interaction with the host. Our findings provide insights into the molecular bases of beneficial legume endophytic–enterobacteria interactions and explore the fine line between beneficial and pathogenic enterobacteria.

Acknowledgments

This work was supported by FEDER funds through the COMPETE 2020 (POCI), and FCT in the framework of the project POCI-01-0145-FEDER-016810 (PTDC/AGR-PRO/2978/2014). CB acknowledges a post-doctoral fellowship (SFRH/BPD/94751/2013) and a contract from the Individual Call to Scientific Employment Stimulus 2018 (CEE-CIND/00093/2018) from FCT. The authors also thank Professor Dr. Kai Thormann, from the Institute for Microbiology and Molecular Biology at Justus Liebig University Giessen, for kindly providing the *E. coli* WM3064 strain, and Isabel Duarte Maças, Estação Nacional de Melhoramento de Plantas, Elvas, Portugal, for her kindness in providing the chickpea seeds.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This work is funded by National Funds through FCT-Foundation for Science and Technology under the Project UIDB/05183/2020.

ORCID

Denise Pereira Torres  <http://orcid.org/0000-0002-2326-9876>

Ana Paço  <http://orcid.org/0000-0003-0362-1630>

Esther Menéndez  <http://orcid.org/0000-0003-2320-0067>

Pedro F. Mateos  <http://orcid.org/0000-0003-1628-598X>

Clarisse Brígido  <http://orcid.org/0000-0002-4167-5253>

References

- Adeolu M, Alnajjar S, Naushad S, Gupta RS. 2016. Genome-based phylogeny and taxonomy of the ‘*Enterobacteriales*’: proposal for *Enterobacterales* ord. nov. divided into the families *Enterobacteriaceae*, *Erwiniaceae* fam. nov., *Pectobacteriaceae* fam. nov., *Yersiniaceae* fam. nov., *Hafniaceae* fam. nov., *Morganellaceae* fam. nov., and *Budviciaceae* fam. nov. *Int J Syst Evol Microbiol.* 66:5575–5599.
- Alquéres S, Meneses C, Rouws L, Rothballer M, Baldani I, Schmid M, Hartmann A. 2013. The bacterial superoxide dismutase and glutathione reductase are crucial for endophytic colonization of rice roots by *Gluconacetobacter diazotrophicus* PAL5. *Mol Plant Microbe Interact.* 26:937–945.

- Arsene F, Kaminski PA, Elmerich C. 1996. Modulation of NifA activity by PII in *Azospirillum brasilense*: evidence for a regulatory role of the NifA N-terminal domain. *J Bacteriol.* 178:4830–4838.
- Arthikala M-K, Montiel J, Sánchez-López R, Nava N, Cárdenas L, Quinto C. 2017. Respiratory burst oxidase homolog gene a is crucial for *Rhizobium* infection and nodule maturation and function in common bean. *Front Plant Sci.* 8:2003.
- Beck M, Wyrsch I, Strutt J, Wimalasekera R, Webb A, Boller T, Robatzek S. 2014. Expression patterns of FLAGELLIN SENSING 2 map to bacterial entry sites in plant shoots and roots. *J Exp Bot.* 65:6487–6498.
- Beringer J. 1974. R factor transfer in *Rhizobium leguminosarum*. *Microbiology.* 84:188–198.
- Bernal P, Llamas MA, Filloux A. 2018. Type VI secretion systems in plant-associated bacteria. *Environ Microbiol.* 20:1–15.
- Boller T, Felix G. 2009. A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu Rev Plant Biol.* 60:379–406.
- Boyer F, Fichant G, Berthod J, Vandenbrouck Y, Attree I. 2009. Dissecting the bacterial type VI secretion system by a genome wide in silico analysis: what can be learned from available microbial genomic resources? *BMC Genomics.* 10:104.
- Breen S, Williams SJ, Outram M, Kobe B, Solomon PS. 2017. Emerging insights into the functions of pathogenesis-related protein 1. *Trends Plant Sci.* 22:871–879.
- Brígido C, Menéndez E, Paço A, Glick BR, Belo A, Félix MR, Oliveira S, Carvalho M. 2019. Mediterranean native leguminous plants: a reservoir of endophytic bacteria with potential to enhance chickpea growth under stress conditions. *Microorganisms.* 7:24.
- Brígido C, Nascimento FX, Duan J, Glick BR, Oliveira S. 2013. Expression of an exogenous 1-aminocyclopropane-1-carboxylate deaminase gene in *Mesorhizobium* spp. reduces the negative effects of salt stress in chickpea. *FEMS Microbiol Lett.* 349:46–53.
- Brígido C, Robledo M, Menéndez E, Mateos PF, Oliveira S. 2012. A ClpB chaperone knockout mutant of *Mesorhizobium ciceri* shows a delay in the root nodulation of chickpea plants. *Mol Plant Microbe Interact.* 25:1594–1604.
- Brígido C, Singh S, Menéndez E, Tavares MJ, Glick BR, Mdr F, Oliveira S, Carvalho M. 2019. Diversity and functionality of culturable endophytic bacterial communities in chickpea plants. *Plants.* 8:42.
- Brock AK, Berger B, Mewis I, Ruppel S. 2013. Impact of the PGPB *Enterobacter radicincitans* DSM 16656 on growth, glucosinolate profile, and immune responses of *Arabidopsis thaliana*. *Microb Ecol.* 65:661–670.
- Brock AK, Berger B, Schreiner M, Ruppel S, Mewis I. 2018. Plant growth-promoting bacteria *Kosakonia radicincitans* mediate anti-herbivore defense in *Arabidopsis thaliana*. *Planta.* 248:1383–1392.
- Broughton WJ, Dilworth MJ. 1971. Control of leghaemoglobin synthesis in snake beans. *Biochem J.* 125:1075–1080.
- Cameron EA, Gruber CC, Ritchie JM, Waldor MK, Sperandio V. 2018. The QseG lipoprotein impacts the virulence of enterohemorrhagic *Escherichia coli* and *Citrobacter rodentium* and regulates flagellar phase variation in *Salmonella enterica* serovar typhimurium. *Infect Immun.* 86:e00936–17.
- Cha C, Gao P, Chen Y-C, Shaw PD, Farrand SK. 1998. Production of acyl-homoserine lactone quorum-sensing signals by gram-negative plant-associated bacteria. *Mol Plant Microbe Interact.* 11:1119–1129.
- Chen T, Duan L, Zhou B, Yu H, Zhu H, Cao Y, Zhang Z. 2017. Interplay of pathogen-induced defense responses and symbiotic establishment in *Medicago truncatula*. *Front Microbiol.* 8:973.
- Cheng H-P, Walker GC. 1998. Succinoglycan is required for initiation and elongation of infection threads during nodulation of alfalfa by *Rhizobium meliloti*. *J Bacteriol.* 180:5183–5191.
- Cole BJ, Feltcher ME, Waters RJ, Wetmore KM, Mucyn TS, Ryan EM, Wang G, Ul-Hasan S, McDonald M, Yoshikuni Y, et al. 2017. Genome-wide identification of bacterial plant colonization genes. *PLoS Biol.* 15:e2002860.
- Confer AW, Ayalew S. 2013. The OmpA family of proteins: roles in bacterial pathogenesis and immunity. *Vet Microbiol.* 163:207–222.
- da-Silva JR, Menéndez E, Eliziário F, Mateos PF, Alexandre A, Oliveira S. 2019. Heterologous expression of *nifA* or *nodD* genes improves chickpea-*Mesorhizobium* symbiotic performance. *Plant Soil.* 436:607–621.
- Dehal PS, Joachimiak MP, Price MN, Bates JT, Baumohl JK, Chivian D, Friedland GD, Huang KH, Keller K, Novichkov PS. 2010. Microbesonline: an integrated portal for comparative and functional genomics. *Nucleic Acids Res.* 38:D396–D400.
- Dehio C, Meyer M. 1997. Maintenance of broad-host-range incompatibility group P and group Q plasmids and transposition of Tn5 in *Bartonella henselae* following conjugal plasmid transfer from *Escherichia coli*. *J Bacteriol.* 179:538–540.
- De Maayer P, Cowan DA. 2016. Flashy flagella: flagellin modification is relatively common and highly versatile among the *Enterobacteriaceae*. *BMC Genomics.* 17:377.
- do Amaral FP, Bueno JCF, Hermes VS, Arisi ACM. 2014. Gene expression analysis of maize seedlings (DKB240 variety) inoculated with plant growth promoting bacterium *Herbaspirillum seropedicae*. *Symbiosis.* 62:41–50.
- Duan J, Müller KM, Charles TC, Vesely S, Glick BR. 2009. 1-aminocyclopropane-1-carboxylate (ACC) deaminase genes in rhizobia from southern Saskatchewan. *Microb Ecol.* 57:423–436.
- Ekanjio LK, Ruppel S, Remus R, Witzel K, Patz S, Becker Y. 2017. Site-directed mutagenesis to deactivate two nitrogenase isozymes of *Kosakonia radicincitans* DSM16656 T. *Can J Microbiol.* 64:97–106.
- Förster A, Planamente S, Manoli E, Lossi NS, Freemont PS, Filloux A. 2014. Coevolution of the ATPase ClpV, the sheath proteins TssB and TssC, and the accessory protein TagJ/HsiE1 distinguishes type VI secretion classes. *J Biol Chem.* 289:33032–33043.
- Fujishige NA, Kapadia NN, De Hoff PL, Hirsch AM. 2006. Investigations of *Rhizobium* biofilm formation. *FE0MS Microbiol Ecol.* 56:195–206.
- García-González T, Sáenz-Hidalgo HK, Silva-Rojas HV, Morales-Nieto C, Vancheva T, Koebnik R, Ávila-Quezada GD. 2018. *Enterobacter cloacae*, an emerging plant-pathogenic bacterium affecting chili pepper seedlings. *Plant Pathol J.* 34:1.
- Göpel Y, Görke B. 2018. Interaction of lipoprotein QseG with sensor kinase QseE in the periplasm controls the phosphorylation state of the two-component system QseE/QseF in *Escherichia coli*. *PLoS Genet.* 14:e1007547.
- Göpel Y, Lüttmann D, Heroven AK, Reichenbach B, Dersch P, Görke B. 2011. Common and divergent features in transcriptional control of the homologous small RNAs GlnY and GlnZ in *Enterobacteriaceae*. *Nucleic Acids Res.* 39:1294–1309.
- Gupta S, Bhar A, Chatterjee M, Das S. 2013. *Fusarium oxysporum* f. sp. *ciceri* race 1 induced redox state alterations are coupled to downstream defense signaling in root tissues of chickpea (*Cicer arietinum* L.). *PLoS One.* 8(9): e73163.
- Inaba J, Thornton J, Huergo LF, Monteiro RA, Klassen G, Pedrosa F, Merrick M, de Souza EM. 2015. Mutational analysis of GlnB residues critical for NifA activation in *Azospirillum brasilense*. *Microbiol Res.* 171:65–72.
- Jha CK, Aeron A, Patel BV, Maheshwari DK, Saraf M. 2011. *Enterobacter*: role in plant growth promotion. In: Maheshwari D., editor. *Bacteria in agrobiolgy: plant growth responses*. Berlin, Heidelberg: Springer; p. 159–182. https://doi.org/10.1007/978-3-642-20332-9_8.
- Kai K. 2018. Bacterial quorum sensing in symbiotic and pathogenic relationships with hosts. *Biosci Biotechnol Biochem.* 82:363–371.
- Lassak J, Henche A-L, Binnenkade L, Thormann KM. 2010. Arcs, the cognate sensor kinase in an atypical Arc system of *Shewanella oneidensis* MR-1. *Appl Environ Microbiol.* 76:3263–3274.
- Liu H, Carvalhais LC, Crawford M, Singh E, Dennis PG, Pieterse CMJ, Schenk PM. 2017. Inner plant values: diversity, colonization and benefits from endophytic bacteria. *Front Microbiol.* 8:2552.
- Loh J, Pierson EA, Pierson III LS, Stacey G, Chatterjee A. 2002. Quorum sensing in plant-associated bacteria. *Curr Opin Plant Biol.* 5:285–290.
- Madhaiyan M, Saravanan VS, Blom J, Smits TH, Rezzonico F, Kim S-J, Weon H-Y, Kwon S-W, Whitman WB, Ji L. 2020. *Phytobacter palmae* sp. nov., a novel endophytic, N2 fixing, plant growth promoting gammaproteobacterium isolated from oil palm (*Elaeis guineensis* Jacq.). *Int J Syst Evol Microbiol.* 70:841–848.
- McClellan KH, Winson MK, Fish L, Taylor A, Chhabra SR, Camara M, Daykin M, Lamb JH, Swift S, Bycroft BW. 1997. Quorum sensing and *Chromobacterium violaceum*: exploitation of violacein production and inhibition for the detection of N-acyl-homoserine lactones. *Microbiology.* 143:3703–3711.

- Mertschnigg T, Patz S, Becker M, Feierl G, Ruppel S, Bunk B, Spröer C, Overmann J, Zarfel G. 2020. First report of *Kosakonia radicincitans* bacteraemia from Europe (Austria)-Identification and whole-genome sequencing of strain DSM 107547. *Sci Rep.* 10:1–6.
- Morales J, Kadota Y, Zipfel C, Molina A, Torres M-A. 2016. The Arabidopsis NADPH oxidases *RbohD* and *RbohF* display differential expression patterns and contributions during plant immunity. *J Exp Bot.* 67:1663–1676.
- Morton ER, Fuqua C. 2012. Laboratory maintenance of *Agrobacterium*. *Curr Protoc Microbiol.* 24:3D. 1.1–3D. 1.6.
- Nascimento F, Brígido C, Alho L, Glick B, Oliveira S. 2012. Enhanced chickpea growth-promotion ability of a *Mesorhizobium* strain expressing an exogenous ACC deaminase gene. *Plant Soil.* 353:221–230.
- O'Toole GA. 2011. Microtiter dish biofilm formation assay. *J Vis Exp.* 47:2437.
- Paço A, Brígido C, Alexandre A, Mateos PF, Oliveira S. 2016. The symbiotic performance of chickpea rhizobia can be improved by additional copies of the *clpB* chaperone gene. *PLoS One.* 11: e0148221.
- Paço A, da-Silva JR, Torres DP, Glick BR, Brígido C. 2020. Exogenous ACC deaminase is key to improving the performance of pasture legume-rhizobial symbioses in the presence of a high manganese concentration. *Plants.* 9:1630.
- Pillonetto M, Arend LN, Faoro H, D'Espindula HR, Blom J, Smits TH, Mira MT, Rezzonico F. 2018. Emended description of the genus *Phytobacter*, its type species *Phytobacter diazotrophicus* (Zhang 2008) and description of *Phytobacter ursingii* sp. nov. *Int J Syst Evol Microbiol.* 68:176–184.
- Raaijmakers JM, Leeman M, Van Oorschot MM, Van der Sluis I, Schippers B, Bakker P. 1995. Dose-response relationships in biological control of Fusarium wilt of radish by *Pseudomonas* spp. *Phytopathology.* 85:1075–1080.
- Rashid S, Charles TC, Glick BR. 2012. Isolation and characterization of new plant growth-promoting bacterial endophytes. *Appl Soil Ecol.* 61:217–224.
- Reading NC, Rasko D, Torres AG, Sperandio V. 2010. A transcriptome study of the QseEF two-component system and the QseG membrane protein in enterohaemorrhagic *Escherichia coli* O157: H7. *Microbiology.* 156:1167.
- Reading NC, Rasko DA, Torres AG, Sperandio V. 2009. The two-component system QseEF and the membrane protein QseG link adrenergic and stress sensing to bacterial pathogenesis. *Proc Natl Acad Sci USA.* 106:5889–5894.
- Reading NC, Torres AG, Kendall MM, Hughes DT, Yamamoto K, Sperandio V. 2007. A novel two-component signaling system that activates transcription of an enterohemorrhagic *Escherichia coli* effector involved in remodeling of host actin. *J Bacteriol.* 189:2468–2476.
- Reichenbach B, Göpel Y, Görke B. 2009. Dual control by perfectly overlapping σ 54- and σ 70-promoters adjusts small RNA GlmY expression to different environmental signals. *Mol Microbiol.* 74:1054–1070.
- Salimiyan Rizi K, Ghazvini K, Farsiani H. 2020. Clinical and pathogenesis overview of Enterobacter infections. *Rev Clin Med.* 6:146–154.
- Sambrook J, Russell D. 2001. Molecular cloning: a laboratory manual. 3rd ed. New York: Cold Spring Harbor Laboratory Press.
- Smit P, Raedts J, Portyanko V, Debellé F, Gough C, Bisseling T, Geurts R. 2005. NSP1 of the GRAS protein family is essential for rhizobial Nod factor-induced transcription. *Science.* 308:1789–1791.
- Smith SG, Mahon V, Lambert MA, Fagan RP. 2007. A molecular Swiss army knife: OmpA structure, function and expression. *FEMS Microbiol Lett.* 273:1–11.
- Sotomaior P, Araújo LM, Nishikawa CY, Huergo LF, Monteiro RA, Pedrosa FO, Chubatsu LS, Souza EM. 2012. Effect of ATP and 2-oxoglutarate on the in vitro interaction between the NifA GAF domain and the GlnB protein of *Azospirillum brasilense*. *Braz J Med Biol Res.* 2012(45):1135–1140.
- Srivastava P, Sharma P, Dogra R. 1999. Inducers of nod genes of *Rhizobium ciceri*. *Microbiol Res.* 154:49–55.
- Surette MA, Sturz AV, Lada RR, Nowak J. 2003. Bacterial endophytes in processing carrots (*Daucus carota* L. var. sativus): their localization, population density, biodiversity and their effects on plant growth. *Plant Soil.* 253:381–390.
- Tjamos SE, Flemetakis E, Paplomatas EJ, Katinakis P. 2005. Induction of resistance to *Verticillium dahliae* in *Arabidopsis thaliana* by the biocontrol agent K-165 and pathogenesis-related proteins gene expression. *Mol Plant Microbe Interact.* 18:555–561.
- Torres DP, Proels RK, Schempp H, Hüchelhoven R. 2017. Silencing of *RBOHF2* causes leaf age-dependent accelerated senescence, salicylic acid accumulation, and powdery mildew resistance in barley. *Mol Plant Microbe Interact.* 30:906–918.
- Torres MA, Dangl JL. 2005. Functions of the respiratory burst oxidase in biotic interactions, abiotic stress and development. *Curr Opin Plant Biol.* 8:397–403.
- van Loon LC, Bakker PA, van der Heijden WH, Wendehenne D, Pugin A. 2008. Early responses of tobacco suspension cells to rhizobacterial elicitors of induced systemic resistance. *Mol Plant Microbe Interact.* 21:1609–1621.
- Verma A, Arora SK, Kuravi SK, Ramphal R. 2005. Roles of specific amino acids in the N terminus of *Pseudomonas aeruginosa* flagellin and of flagellin glycosylation in the innate immune response. *Infect Immun.* 73:8237–8246.
- Wahlg TA, Bixler BJ, Valdés-López O, Mysore KS, Wen J, Ané J-M, Kaspar CW. 2019. *Salmonella enterica* serovar Typhimurium ATCC 14028S is tolerant to plant defenses triggered by the flagellin receptor FLS2. *FEMS Microbiol Lett.* 366:fny296.
- Willems E, Leyns L, Vandesompele J. 2008. Standardization of real-time PCR gene expression data from independent biological replicates. *Anal Biochem.* 379:127–129.
- Wyrsh I, Domínguez-Ferreras A, Geldner N, Boller T. 2015. Tissue-specific FLAGELLIN-SENSING 2 (FLS2) expression in roots restores immune responses in Arabidopsis fls2 mutants. *New Phytol.* 206:774–784.
- Xiao J, Chen T, Yang M, Zhang Y, Wang Q. 2012. Identification of *qseEGF* genetic locus and its roles in controlling hemolytic activity and invasion in fish pathogen *Edwardsiella tarda*. *Lett Appl Microbiol.* 55:91–98.
- Yamazaki A, Hirata H, Tsuyumu S. 2008. HrpG regulates type II secretory proteins in *Xanthomonas axonopodis* pv. *citri*. *J Gen Plant Pathol.* 74:138–150.
- Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. 2012. Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC Bioinform.* 13:134.
- Yu K, Pieterse CM, Bakker PA, Berendsen RL. 2019. Beneficial microbes going underground of root immunity. *Plant Cell Environ.* 42:2860–2870.
- Zamioudis C, Pieterse CM. 2012. Modulation of host immunity by beneficial microbes. *Mol Plant Microbe Interact.* 25:139–150.
- Zipfel C, Oldroyd GE. 2017. Plant signalling in symbiosis and immunity. *Nature.* 543:328.