

animal

<http://journals.cambridge.org/ANM>

Additional services for **animal**:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Fat area and lipid droplet morphology of porcine oocytes during *in vitro* maturation with *trans*-10, *cis*-12 conjugated linoleic acid and forskolin

E. G. Prates, C. C. Marques, M. C. Baptista, M. I. Vasques, N. Carolino, A. E. M. Horta, R. Charneca, J. T. Nunes and R. M. Pereira

animal / *FirstView* Article / November 2012, pp 1 - 8
DOI: 10.1017/S1751731112001899, Published online:

Link to this article: http://journals.cambridge.org/abstract_S1751731112001899

How to cite this article:

E. G. Prates, C. C. Marques, M. C. Baptista, M. I. Vasques, N. Carolino, A. E. M. Horta, R. Charneca, J. T. Nunes and R. M. Pereira Fat area and lipid droplet morphology of porcine oocytes during *in vitro* maturation with *trans*-10, *cis*-12 conjugated linoleic acid and forskolin. animal, Available on CJO doi:10.1017/S1751731112001899

Request Permissions : [Click here](#)

Fat area and lipid droplet morphology of porcine oocytes during *in vitro* maturation with *trans*-10, *cis*-12 conjugated linoleic acid and forskolin

E. G. Prates^{1,2}, C. C. Marques¹, M. C. Baptista¹, M. I. Vasques¹, N. Carolino^{1,3},
A. E. M. Horta¹, R. Charneca², J. T. Nunes² and R. M. Pereira^{1,3†}

¹L-INIAV Santarém, Quinta da Fonte Boa, 00-048 Vale de Santarém, Portugal; ²Universidade de Évora – Instituto de Ciências Agrárias e Ambientais Mediterrânicas (ICAAM), 7002-554 Évora, Portugal; ³Escola Universitária Vasco da Gama, Mosteiro de S. Jorge de Milrêu, 3040-714 Coimbra, Portugal

(Received 22 February 2012; Accepted 1 August 2012)

Lipid droplets (LD) in porcine oocytes form a dark mass reaching almost all cytoplasm. Herein we investigated changes in fat areas, cytoplasmic tone and LD morphology during in vitro maturation (IVM) of porcine oocytes cultured with 100 μM trans-10, cis-12 conjugated linoleic acid (t10,c12 CLA) or 10 μM forskolin at different time periods. Four groups were constituted: control, excipient, t10,c12 CLA and forskolin, with drugs being supplemented during 44 to 48 h and the initial 22 to 24 h in Experiments 1 and 2, respectively. In Experiment 3, forskolin was supplemented for the first 2 h. Matured oocytes were inseminated with frozen-thawed boar semen and cleavage rate recorded. Before and during IVM, samples of oocytes were evaluated for LD, total and fat areas and fat gray value or for meiotic progression. Results showed that forskolin supplementation during 44 to 48 h or 22 to 24 h inhibits oocyte maturation (exp. 1: forskolin = 5.1 ± 8.0%, control = 72.6 ± 5.0%; exp. 2: forskolin = 24.3 ± 7.4%, control = 71.6 ± 5.6%) and cleavage (exp. 1: forskolin = 0.0 ± 0.0%, control = 55.4 ± 4.1%; exp. 2: forskolin = 8.3 ± 3.3%, control = 54.5 ± 3.0%). Forskolin also reduced oocyte and fat areas. In Experiment 3, forskolin negative effect on oocyte maturation and cleavage disappeared, although minor (P ≤ 0.03) LD and oocyte fat areas were identified at 22 to 24 h of IVM. Oocytes supplemented with t10,c12 CLA during 44 to 48 h presented a lighter (P ≤ 0.04) colour tone cytoplasm than those of control and forskolin. In conclusion, t10,c12 CLA and forskolin were capable of modifying the distribution and morphology of cytoplasmic LD during porcine oocyte maturation, thus reducing its lipid content in a time-dependent manner.

Keywords: lipid droplets, porcine oocyte maturation, lipid content, conjugated linoleic acid, forskolin

Implications

Swine is relevant for agriculture and livestock economy. *In vitro* embryo production can be an effective system for boar fertility evaluation and germoplasm cryopreservation, but also for biomedical research. However, nuclear and cytoplasmic maturation asynchrony and the high intracellular lipid content in porcine oocyte are still limiting factors for the application of such biotechnologies in this species. The present work increases understanding on the effect of two substances, *trans*-10, *cis*-12 conjugated linoleic acid and forskolin, capable of lipid modulating to improve oocyte developmental competence and embryo yield.

Introduction

Pigs are important not only as livestock, but also as an experimental model for humans in biomedical research.

Obesity and its physiological consequences are increasingly prevalent among women of reproductive age and are associated with infertility; porcine oocytes and embryos are excellent models to study this paradigm (Gesink *et al.*, 2007; Wang *et al.*, 2009). The negative effect of an excessive intracellular lipid content of porcine oocytes and embryos on assisted reproductive technologies has been extensively reported (Dobrinsky *et al.*, 1999; Men *et al.*, 2006; Pereira and Marques, 2008). Lipid droplets (LD) are organelles constituted primarily by triacylglycerols and cholesteryl esters surrounded by a monolayer of phospholipids with embedded integral and peripheral proteins, occupying a considerably mass in the eukaryotic cells, particularly in the mammalian oocytes and embryos (Fujihira *et al.*, 2004; Walther and Farese, 2009; Zehmer *et al.*, 2009). LD in porcine oocytes form a dark mass reaching almost all cytoplasm, but their function and importance are scarcely known. This dark appearance of porcine oocytes and embryos, compared with others species, has been attributed to an increased lipid

† E-mail: rosaln@gmail.com

content (McEvoy *et al.*, 2000; Genicot *et al.*, 2005). On the other hand, the intermediate filaments of the cytoskeleton interact directly with these special organelles (Zehmer *et al.*, 2009). Therefore, it is not surprising that during gamete and embryo cryopreservation, morphological variations occurring in LD as consequence of physical changes of lipids lead to irreversible damages in the cytoskeleton (Fujihira *et al.*, 2004; Pereira and Marques, 2008).

Recently, two chemicals, *trans*-10, *cis*-12 conjugated linoleic acid (τ 10, τ 12 CLA) and forskolin, intracellular lipid modulators, were shown to improve porcine and bovine embryo survival after cryopreservation (Men *et al.*, 2006; Pereira *et al.*, 2007 and 2008). Indeed, *in vitro*-produced bovine embryos cultured with 100 μ M τ 10, τ 12 CLA were characterized by a reduced lipid accumulation and an improved blastocyst cryosurvival (Pereira *et al.*, 2007 and 2008). Moreover, alterations in the lipid profile of bovine oocytes matured with this CLA isomer were associated with an enhanced blastocyst quality (Lapa *et al.*, 2011). In pig, τ 10, τ 12 CLA exerts specific effects on adipocytes by diminishing fat deposition through lipogenesis reduction, lipolysis and fatty acid oxidation increase (Corino *et al.*, 2006). Although this CLA isomer has never been tested in porcine oocytes, it is expected to induce alterations in cytoplasmic lipid content during *in vitro* maturation (IVM).

Forskolin, an adenylyl cyclase stimulator, at 10 μ M dose is an effective lipolytic substance capable of improving oocyte and embryo cryosurvival (Men *et al.*, 2006; Fu *et al.*, 2011). However, despite the contribution of forskolin to lipid reduction, it can simultaneously arrest meiotic resumption during porcine oocytes IVM in a dose-dependent response. At 10 μ M dose, the meiotic arrest was fully reversible (Laforest *et al.*, 2005). Moreover, by delaying spontaneous meiotic resumption, forskolin can increase oocyte development potential owing to a better synchronization of nuclear and cytoplasmic maturation (Thomas *et al.*, 2004). Thus, the possibility of embracing both forskolin effects on meiosis synchronization and lipid content reduction in porcine oocytes should be investigated using different supplementation time periods.

The present study investigates the effect of supplementing maturation media of porcine oocytes with 100 μ M τ 10, τ 12 CLA or 10 μ M forskolin, during different time periods, on oocyte intracellular lipid content and maturation process. The developmental potential of those oocytes for cleavage was also assessed.

Material and methods

Oocyte recovery and IVM

Otherwise stated, all products were purchased from Sigma, Sintra, Portugal. Prepubertal gilt ovaries were obtained from a local slaughterhouse. At the laboratory, medium-sized follicles (3 to 6 mm) were aspirated, cumulus–oocyte complexes (COC) collected and washed in Dulbecco's phosphate-buffered saline medium (mDPBS; Gil *et al.*, 2003). Oocytes with a dark, evenly granulated cytoplasm and surrounded by a compact cumulus cell mass were selected and washed in

maturation medium, the BSA-free North Carolina State University medium (NCSU-23; Petters and Wells, 1993) supplemented with 10% (v/v) porcine follicular fluid, 0.1 mg/ml cysteine, 10 ng/ml epidermal growth factor and 100 μ M glutathione (GSH). Oocytes were cultured in pre-equilibrated maturation medium (3 ml) containing 10 IU/ml equine chorionic gonadotrophin (eCG; Intervet, Agualva-Cacem, Portugal) and 10 IU/ml human chorionic gonadotrophin (hCG; Pregnyl, Organon, Portugal) for 22 to 24 h and then for another 22 to 24 h without hormones, at 39°C in an atmosphere of 5% CO₂ in air.

Fertilization and embryo culture

In vitro fertilization and embryo culture were performed according to Gil *et al.* (2003). Briefly, denuded oocytes were washed in modified Tris-buffered fertilization medium. Oocytes were then placed in 50 μ l droplets of this medium under mineral oil (30 oocytes/drop). Frozen-thawed boar sperm (100 μ l) were washed in mDPBS by centrifugation (1900 g/3 min). The sperm pellet was resuspended in fertilization medium to obtain a final concentration of 1×10^6 spz/ml and added to oocytes. After 6 h of co-incubation, presumptive zygotes were washed in embryo culture medium (NCSU-23 with 0.4% BSA) and placed in a 4-well multidish for embryo culture during 48 h at 39°C and 5% CO₂ in humidified air.

Assessment of oocyte maturation and cleavage rates

Meiotic progression assessment was performed at 0 h, between 22 to 24 h and 44 to 48 h of IVM. COC were gently pipetted in a 1% sodium citrate solution to remove cumulus cells. Denuded oocytes were mounted on slides, fixed in an acetic acid:ethanol solution (1:3) for 48 to 72 h, then stained with 1% lacmoid in 45% (v/v) acetic acid before examination under a phase-contrast microscope ($\times 400$). Oocytes were classified according to their chromosome configuration as: germinal vesicle (GV); germinal vesicle breakdown (GVBD) till metaphase-I (MI); and anaphase (Anaph.) to metaphase-II (MII). Oocytes in the latest phase were considered matured. Fixed oocytes presenting a broken oolemma or abnormal cytoplasmic appearance were discharged.

Cleavage rates (cleaved embryos/inseminated oocytes $\times 100$) were assessed at 48 h after oocyte insemination.

Lipid content evaluation

Fresh oocyte samples, before IVM (0 h), at 22 to 24 h and 44 to 48 h of IVM, were examined under Nomarski differential interference contrast microscope for lipid content evaluation (Pereira *et al.*, 2008). Oocytes were photographed using an Olympus camera (Olympus Portugal, Lisboa, Portugal) attached to the microscope and analysed using ImageJ software (version 1.44q, National Institutes of Health, USA). A total of 100 LD randomly distributed at different focal planes were evaluated per oocyte. Total area occupied by LD, denominated fat area, and the gray mean value within fat area of each oocyte were also measured with the same equatorial focal plane, brightness and contrast. This mean value was calculated by converting each pixel to a gray scale by ImageJ software (Figure 1). On the basis of the gray mean value within the fat




Figure 1 Fat areas (arrows) of porcine oocytes evaluated by their gray mean values after Nomarski microscopy observation at the equatorial plan: a dark oocyte from control group after 44 to 48 h of *in vitro* maturation (IVM) (a) and the correspondent gray value histogram (b); a light oocyte from the *trans*-10, *cis*-12 conjugated linoleic acid treatment after 44 to 48 h of IVM (c) and the correspondent gray value histogram (d). Gray scale: 0 pixels/ μm^2 – black to 255 pixels/ μm^2 – white.

area at 44 to 48 h of IVM, oocytes were classified into two categories: dark (<112 pixels/ μm^2) or light (>112 pixels/ μm^2) gray. In addition, cytoplasmic and total (including *zona pellucida*) areas of each oocyte were recorded. Fat oocyte index was calculated (fat area/total oocyte area \times 100).

Experimental design

Three experiments (three replicates each) were designed. *Trans*-10, *cis*-12 octadecadienoic acid (100 μM †10,†12 CLA, Matreya, Pleasant Gap, Pennsylvania, USA; Pereira *et al.*, 2007) and forskolin (10 μM , Men *et al.*, 2006) were first diluted in ethanol because of its hydrophobicity and added to oocytes in the maturation media using different supplementation times according to the following experiments. The final concentration of ethanol was adjusted to 0.1% in all treatment groups.

Experiment 1. The objective of this experiment was to study the effect of †10,†12 CLA and forskolin, administered during total IVM period on porcine oocyte lipid content, maturation and developmental potential. Four experimental groups were designed: (1) Control – COC ($n = 373$) were matured during 44 to 48 h without supplementation; (2) Ethanol – COC ($n = 337$) were matured during 44 to 48 h supplemented with 0.1% ethanol; (3) †10,†12 CLA – COC ($n = 322$) were matured during 44 to 48 h with 100 μM †10,†12 CLA; (4) Forskolin – COC ($n = 353$) were matured during 44 to 48 h with 10 μM forskolin. After maturation, COC were inseminated ($n = 816$) with frozen-thawed boar semen and cleavage rates recorded. At 22 to 24 h and 44 to 48 h of IVM, samples of oocytes were assessed for meiosis progression ($n = 493$) and lipid content evaluation ($n = 76$). Immature oocytes ($n = 97$, 87 for meiosis progression and 10 for lipid content evaluation, respectively) were also evaluated.

Experiment 2. In Experiment 2, supplementation period of 100 μM †10,†12 CLA and 10 μM forskolin was reduced to the first 22 to 24 h followed by 22 to 24 h of standard protocol of IVM and their effect on porcine oocyte lipid content, maturation and developmental potential also studied. Four experimental groups were also established: (1) Control – COC ($n = 373$) were matured during 44 to 48 h without supplementation; (2) Ethanol – COC ($n = 432$) were matured during the initial 22 to 24 h with 0.1% ethanol; (3) †10,†12 CLA – COC ($n = 330$) were matured with 100 μM †10,†12 CLA during the initial 22 to 24 h; (4) Forskolin – COC ($n = 354$) were matured during the initial 22 to 24 h with 10 μM forskolin. After 44 to 48 h of maturation, COC were inseminated ($n = 908$) and cleavage rates recorded. At 22 to 24 h and 44 to 48 h, IVM samples of oocytes were assessed for meiosis progression ($n = 506$) and lipid content evaluation ($n = 75$). Immature oocytes ($n = 97$) were also evaluated.

Experiment 3. In the previous experiments, supplementation periods of forskolin showed deleterious effects on oocyte maturation and cleavage rates. A third experiment was performed to investigate whether a time reduction in forskolin supplementation up to the initial 2 h of IVM could avoid these effects, simultaneously maintaining its lipolytic effects. †10,†12 CLA and ethanol were not tested in this trial, as in previous experiments the former did not interfere with oocyte meiotic progression and the latter did interfere neither with oocyte meiotic progression nor with lipid content. Therefore, two experimental groups were performed: (1) Control – COC ($n = 291$) were matured without supplementation; (2) Forskolin – COC ($n = 299$) were matured with 10 μM forskolin during the initial 2 h culture. After 44 to 48 h of maturation, COC ($n = 238$) were inseminated and cleavage rates recorded. At 22 to 24 h and 44 to 48 h, IVM samples of oocytes were assessed for meiosis progression ($n = 307$) and lipid evaluation ($n = 45$). Immature oocytes ($n = 103$) were also evaluated.

Statistical analysis

All results are expressed as least square means (\pm standard error of means in text). In each experiment, data representing three replicates of cleavage and oocyte nuclear configuration, as well as of oocyte lipid content evaluation, were analysed using the MIXED procedure of Statistical Analysis System (SAS Institute, Cary, NC, USA). The mixed linear model included treatment or treatment and IVM time periods (0 h, 22 to 24 h and 44 to 48 h) as fixed effects for cleavage and oocyte nuclear configuration evaluation, respectively, and replicates as random effect. For lipid content evaluation, treatment and IVM time periods were also considered fixed effects, but the oocyte nested in treatment was the random effect. LD were treated as repeated measures within oocyte. When significant effects were identified, differences between individual treatments were determined by the PDIFF multiple comparison procedure.

Data from Nomarski records were analysed using a multiple regression analysis (Proc Path of SAS), testing whether the fat

Table 1 Effect of τ 10, τ 12 CLA and forskolin supplementation during 44 to 48 h of *in vitro* maturation on porcine oocyte meiosis progression, oocyte and LD areas, in three independent replicates

Treatments	Oocyte meiosis progression				Oocyte areas				LD		
	<i>n</i>	GV (%)	GVBD-MI (%)	Anaph.I-MII (%)	<i>n</i>	Total (μm^2)	Cytoplasmic (μm^2)	Fat area (μm^2)	Fat index (%)	<i>n</i>	LD area (μm^2)
Immature											
0 h	87	97.7 ^a	2.3 ^d	0.0 ^b	10	18 099 ^{de}	12 445 ^{bc}	9997 ^b	54.8	800	12.4
Control											
22 to 24 h	61	23.1 ^{cd}	76.9 ^a	0.0 ^b	10	19 414 ^{abcd}	14 173 ^{ab}	12 003 ^a	61.3	800	10.8
44 to 48 h	58	0.0 ^d	28.0 ^{bc}	72.6 ^a	10	19 963 ^{ab}	14 040 ^{ab}	10 814 ^{ab}	56.5	800	11.5
Ethanol											
22 to 24 h	61	29.2 ^{bc}	69.6 ^a	1.3 ^b	9	18 806 ^{bcd}	13 937 ^{ab}	11 416 ^{ab}	58.2	800	10.6
44 to 48 h	61	18.1 ^{cd}	17.8 ^{bcd}	65.1 ^a	11	20 503 ^{ab}	15 533 ^a	12 741 ^a	60.0	800	11.0
τ 10, τ 12 CLA											
22 to 24 h	51	23.9 ^c	72.5 ^a	37.4 ^b	9	19 791 ^{abc}	14 092 ^{ab}	11 401 ^{ab}	57.1	800	8.7
44 to 48 h	68	19.3 ^{cd}	20.7 ^{bcd}	61.0 ^a	9	20 736 ^a	14 373 ^a	11 332 ^{ab}	55.2	800	11.4
Forskolin											
22 to 24 h	71	93.4 ^a	6.6 ^{cd}	0.0 ^b	9	17 362 ^e	11 784 ^c	9759 ^b	55.8	800	10.3
44 to 48 h	62	55.6 ^b	40.0 ^b	5.1 ^b	9	17 694 ^{de}	12 063 ^c	9886 ^b	54.9	800	10.0
r.s.d.		34.2	37.7	28.4		2014	2006	2012	7.0		7.1
<i>P</i>		<0.001	<0.001	<0.001		0.002	0.001	0.015	0.311		0.129

τ 10, τ 12 CLA = *trans*-10, *cis*-12 conjugated linoleic acid; LD = lipid droplets; GV = germinal vesicle; GVBD-MI = germinal vesicle breakdown up to metaphase I; Anaph.I-MII = anaphase I up to metaphase II; r.s.d. = residual standard deviation.

Oocyte areas: total (cytoplasmic plus *zona pellucida*); cytoplasmic (cytoplasmic, without *zona pellucida*); fat area (total area occupied by LD); fat index (fat area/oocyte total area \times 100).

LD area: lipid droplets area within oocytes.

Significant differences were identified on oocyte meiosis progression and oocyte areas.

Data were analysed using the MIXED procedure of Statistical Analysis System considering treatment and *in vitro* maturation time periods as fixed effects.

^{a,b,c,d,e}Values (least square means) with different superscripts letters in the same column are significantly different ($P \leq 0.05$).

gray tones were correlated to other individual or associated variables. An appropriate multicollinearity test (Proc Reg) was performed. The χ^2 test (Proc Freq) was used to compare the distribution of oocytes into categories (dark and light gray) of fat gray value. Differences were considered significant when $P \leq 0.05$.

Results

Fat gray value was independent from oocyte total ($r = 0.07$, $P = 0.48$), fat ($r = -0.02$, $P = 0.84$) and LD ($r = -0.04$, $P = 0.58$) areas and fat oocyte index ($r = -0.14$, $P = 0.10$). Instead, fat area and fat oocyte index were highly correlated ($r = 0.81$, $P < 0.0001$). This association explained 4% of fat gray value (tolerance value = 0.35, variance inflation = 2.8). The path coefficient from fat area to fat oocyte index was +1.50.

Experiment 1

Data from meiosis progression of porcine oocytes supplemented with τ 10, τ 12 CLA and forskolin during 44 to 48 h of IVM are represented in Table 1. As expected, the GV configuration was predominant in immature oocytes. At 22 to 24 h of IVM, $93.4 \pm 7.9\%$ ($P < 0.0001$) of oocytes treated with forskolin remained at GV, whereas oocytes from control, ethanol and τ 10, τ 12 CLA treatments mainly reached GVBD-MI ($P < 0.0001$). Moreover, at the end of IVM, only

$5.1 \pm 5.1\%$ of forskolin-treated oocytes reached Anaph.-MII compared with $72.6 \pm 5.0\%$, $65.1 \pm 5.1\%$ and $61.0 \pm 5.0\%$ from control, ethanol and τ 10, τ 12 CLA treatments ($P < 0.0001$), respectively. At each time period, no differences ($P > 0.05$) were identified in nuclear configuration among oocytes from control, ethanol and τ 10, τ 12 CLA treatments.

The cleavage rate of forskolin-matured oocytes was 0.0%, different ($P \leq 0.0001$) from those of control ($55.4 \pm 4.1\%$), ethanol ($58.9 \pm 4.2\%$) and τ 10, τ 12 CLA ($45.6 \pm 4.2\%$) treatments.

At the end of IVM, total areas of oocytes from control ($P = 0.04$), ethanol ($P = 0.008$) and τ 10, τ 12 CLA ($P = 0.006$) treatments were larger than those of immature oocytes (Table 1), whereas areas of forskolin-treated oocytes were similar to those of immature oocytes. Cytoplasmic areas of forskolin-treated oocytes were the smallest at 22 to 24 h ($P \leq 0.03$) and also at 44 to 48 h ($P \leq 0.05$) of IVM. At the end of maturation, cytoplasmic areas of τ 10, τ 12 CLA and ethanol-treated oocytes were larger ($P \leq 0.04$) than those of immature but similar to control. Fat areas were smaller in immature and forskolin-treated oocytes (22 to 24 h and 44 to 48 h) than in control at 22 to 24 h ($P \leq 0.03$) and ethanol at 44 to 48 h ($P \leq 0.0002$) of IVM. No differences were identified among groups in fat oocyte index and LD areas. However, after 44 to 48 h of culture, τ 10, τ 12 CLA oocytes were lighter than those of control ($P = 0.03$) and forskolin ($P = 0.04$) treatments (Figure 2).

Experiment 2

Results from meiosis progression of porcine oocytes supplemented with t10,c12 CLA and forskolin during the initial 22 to 24 h of IVM are shown in Table 2. In immature oocytes, GV configuration was predominant. Forskolin-treated oocytes remained mainly at GV stage at 22 to 24 h of IVM ($91.0 \pm 7.3\%$, $P < 0.0001$). On the contrary, the majority of oocytes from control, ethanol and t10,c12 CLA treatments were at GVBD-MI ($P < 0.0001$). At 44 to 48 h of IVM,

$24.3 \pm 5.4\%$ of oocytes treated with forskolin during 22 to 24 h reached Anaph.-MII. This rate continues lower ($P < 0.0001$) than those from control, ethanol and t10,c12 CLA treatments. At each time period, no differences ($P > 0.05$) were identified in nuclear configuration among oocytes from control, ethanol and t10,c12 CLA treatments.

The cleavage rate of oocytes matured with forskolin ($8.3 \pm 3.3\%$) was lower ($P \leq 0.0001$) than those of control ($54.5 \pm 3.0\%$), ethanol ($44.9 \pm 3.0\%$) and t10,c12 CLA ($50.5 \pm 4.2\%$) treatments.

After 44 to 48 h of maturation, total areas of oocytes from control and t10,c12 CLA treatments were larger ($P \leq 0.05$) than those of immature and forskolin at 22 to 24 h of maturation (Table 2). These latter oocytes presented also the smallest cytoplasmic areas ($P \leq 0.02$). After 44 to 48 h of maturation with forskolin supplementation during 22 to 24 h, oocyte areas (total and cytoplasmic) were not different from those of other groups at the same time period. At 22 to 24 h of IVM, oocytes cultured with forskolin presented smaller fat areas than control ($P = 0.006$) and t10,c12 CLA ($P = 0.05$). However, the forskolin group showed a larger fat area than control ($P = 0.04$) at 44 to 48 h of IVM. Oocyte fat areas of immature (0 h) and forskolin groups at 22 to 24 h were also smaller ($P \leq 0.03$) than those of forskolin and ethanol oocytes at 44 to 48 h of IVM (Table 2). Fat oocyte index of forskolin treatment was lower ($P = 0.02$) than control at 22 to 24 h. At 44 to 48 h of IVM, the former treatment results became superior to control ($P = 0.02$)




Figure 2 Effect of *trans*-10, *cis*-12 conjugated linoleic acid (t10,c12 CLA) and forskolin, during 44 to 48 h of *in vitro* maturation supplementation, on porcine oocytes lipid content. Oocytes were classified into two categories: dark (<112 pixels/ μm^2) or light (>112 pixels/ μm^2) gray according to the gray mean value within fat area. $a \neq b$, significant differences ($P \leq 0.04$) were observed in gray mean values within fat area of t10,c12 CLA oocytes compared to control or forskolin ones.

Table 2 Effect of t10,c12 CLA and forskolin supplementation during 22 to 24 h of *in vitro* maturation on porcine oocyte meiosis progression, oocyte and LD areas, in three independent replicates

Treatments	Oocyte meiosis progression				Oocyte areas				LD		
	n	GV (%)	GVBD-MI (%)	Anaph.I-MII (%)	n	Total (μm^2)	Cytoplasmic (μm^2)	Fat area (μm^2)	Fat index (%)	n	LD area (μm^2)
Immature											
0 h	87	97.9 ^a	2.2 ^e	0.0 ^c	10	18280 ^{cd}	12499 ^b	10036 ^{cd}	54.8 ^c	800	11.9
Control											
22 to 24 h	61	25.0 ^{bc}	75.8 ^a	0.0 ^c	10	19544 ^{abc}	14206 ^{ab}	12147 ^{ab}	61.9 ^{ab}	800	10.4
44 to 48 h	58	0.0 ^d	28.1 ^c	71.6 ^a	10	20009 ^{ab}	14054 ^{ab}	10903 ^{bcd}	57.0 ^c	800	11.1
Ethanol											
22 to 24 h	61	31.6 ^b	68.8 ^{ab}	1.3 ^c	9	18863 ^{bcd}	13961 ^{ab}	11474 ^{abcd}	58.5 ^{abc}	800	10.1
44 to 48 h	65	6.0 ^{cd}	17.7 ^{cde}	76.7 ^a	8	19747 ^{abc}	14963 ^a	12141 ^{ab}	59.4 ^{abc}	800	12.7
t10,c12 CLA											
22 to 24 h	51	27.6 ^{bc}	69.5 ^{ab}	3.8 ^c	9	19942 ^{abc}	14137 ^{ab}	11521 ^{abc}	57.5 ^{bc}	800	8.3
44 to 48 h	68	6.3 ^{cd}	25.0 ^{cd}	68.3 ^a	10	20995 ^a	14628 ^a	11745 ^{ab}	55.7 ^c	800	9.8
Forskolin											
22 to 24 h	71	91.0 ^a	7.6 ^{de}	0.0 ^c	9	17456 ^d	11805 ^c	9879 ^d	56.3 ^c	800	9.9
44 to 48 h	71	24.8 ^{bc}	53.0 ^b	24.3 ^b	10	19959 ^{abc}	14116 ^{ab}	12573 ^a	62.7 ^a	800	10.3
r.s.d.		30.2	39.0	30.2		1971	1982	1756	5.1		7.2
P		<0.001	<0.001	<0.001		0.010	0.024	0.011	0.011		0.152

t10,c12 CLA = *trans*-10, *cis*-12 conjugated linoleic acid; LD = lipid droplets; GV = germinal vesicle; GVBD-MI = germinal vesicle breakdown up to metaphase I; Anaph.I-MII = anaphase I up to metaphase II; r.s.d. = residual standard deviation.

Oocyte areas: total (cytoplasmic plus *zona pellucida*); cytoplasmic (cytoplasmic, without *zona pellucida*); fat area (total area occupied by LD); fat index (fat area/oocyte total area $\times 100$).

LD area: lipid droplets area within oocytes.

Significant differences were identified on oocyte meiosis progression and oocyte areas.

Data were analysed using the MIXED procedure of Statistical Analysis System considering treatment and *in vitro* maturation time periods as fixed effects.

^{a,b,c,d,e} Values (least square means) with different superscripts letters in the same column are significantly different ($P \leq 0.05$).

Table 3 Effect of forskolin supplementation during the initial 2 h of *in vitro* maturation on porcine oocyte meiosis progression, oocyte and LD areas, in three independent replicates.

Treatments	Oocyte meiosis progression				Oocyte areas				LD		
	<i>n</i>	GV (%)	GVBD-MI (%)	Anaph.I-MII (%)	<i>n</i>	Total (μm^2)	Cytoplasmic (μm^2)	Fat area (μm^2)	Fat index (%)	<i>n</i>	LD area (μm^2)
Immature											
0 h	91	100.0 ^a	0.0 ^c	0.0 ^b	12	17 574 ^b	12 280 ^b	10 161 ^b	57.6 ^{bc}	1000	13.3 ^a
Control											
22 to 24 h	77	19.0 ^{bc}	81.0 ^a	0.0 ^b	11	21 336 ^a	16 294 ^a	14 148 ^a	65.6 ^a	1000	11.5 ^{ab}
44 to 48 h	75	1.1 ^c	29.1 ^{bc}	71.5 ^a	11	20 332 ^a	14 977 ^a	11 049 ^b	54.1 ^c	1000	10.7 ^{bc}
Forskolin											
22 to 24 h	84	43.9 ^b	56.2 ^{ab}	0.0 ^b	11	19 740 ^a	14 132 ^{ab}	11 896 ^b	59.8 ^{ab}	1000	8.0 ^c
44 to 48 h	71	10.4 ^{bc}	32.8 ^{bc}	56.2 ^a	12	20 199 ^a	14 578 ^a	11 054 ^b	55.0 ^{bc}	1000	11.4 ^{ab}
r.s.d.		26.8	34.3	27.0		2402	2609	2375	6.4		7.4
<i>P</i>		0.002	0.04	<0.001		0.007	0.011	0.003	0.0006		0.006

LD = lipid droplets; GV = germinal vesicle; GVBD-MI = germinal vesicle breakdown up to metaphase I; Anaph.I-MII = anaphase I up to metaphase II; r.s.d. = residual standard deviation.

Oocyte areas: total (cytoplasmic plus *zona pellucida*); cytoplasmic (cytoplasmic, without *zona pellucida*); fat area (total area occupied by LD); fat index (fat area/oocyte total area \times 100).

LD area : lipid droplets area within oocytes.

Significant differences were identified on oocyte meiosis progression and oocyte and LD areas.

Data were analysed using the MIXED procedure of Statistical Analysis System considering treatment and *in vitro* maturation time periods as fixed effects.

^{a,b,c}Values (least square means) with different superscripts letters in the same column are significantly different ($P \leq 0.05$).

and also to forskolin ($P = 0.009$) oocytes at 22 to 24 h. No differences were identified between oocyte categories (dark and light gray) or LD areas among groups.

Experiment 3

By reducing the time of forskolin exposure to 2 h, oocyte meiotic progression at 22 to 24 h and 44 to 48 h was similar ($P > 0.05$) to those of control (Table 3). No differences were identified in cleavage rates between treatments.

At the end of IVM, cytoplasmic and total areas of oocytes from both groups were similar among themselves and larger ($P \leq 0.01$) than those of immature oocytes (Table 3). Exception made for cytoplasmic areas of forskolin oocytes, these measurements at 22 to 24 h of IVM were also larger ($P \leq 0.04$) than those of immature oocytes. Control oocytes matured during 22 to 24 h presented the largest ($P \leq 0.03$) fat area (Table 3). Exception made for forskolin oocytes at 22 to 24 h, similar results ($P \leq 0.03$) were obtained for fat oocyte index. This index presented no differences between treatments at the end of IVM. In addition, no differences were identified between oocyte categories (dark and light gray). Although reducing the time exposure to forskolin (2 h), at 22 to 24 h of IVM, these oocytes had smaller LD areas than those of control ($P = 0.03$). These LD were also smaller than those of immature ($P = 0.002$) and forskolin oocytes at the end of IVM ($P = 0.03$). LD areas of immature oocytes were larger ($P = 0.02$) than control ones at 44 to 48 h (Table 3).

Discussion

The present study demonstrated that the supplementation of 100 μM $\text{t}10,\text{c}12$ CLA or 10 μM forskolin during different

time periods distinctively interfere on maturation progress and lipid content of porcine oocytes. Data presented here suggest for the first time that $\text{t}10,\text{c}12$ CLA did not affect meiosis progression or competence for cleavage of porcine oocytes, although matured oocytes were lighter when this CLA isomer was supplemented during the entire maturation period. In opposition, forskolin presence during 44 to 48 h or 22 to 24 h impaired oocyte maturation and their competence for cleavage. However, by decreasing forskolin supplementation interval to the initial 2 h of IVM, these deleterious effects disappeared though morphological changes in LD and oocyte fat areas were identified during maturation progression.

Previous reports of continuing exposure of porcine oocytes to forskolin along maturation have demonstrated an increase of intracellular level of cyclic adenosine monophosphate (cAMP) thus inhibiting GVBD (Racowsky, 1985; Xia *et al.*, 2000). In accordance, since at 44 to 48 h of culture, GV configuration continued to be predominant in forskolin-treated oocytes, remaining their areas similar to immature ones, these oocytes were incapable of normal fertilization and cleavage (0%). In contrast, Fu *et al.* (2011) achieved a cleavage rate of $64.5 \pm 3.6\%$ in parthenogenetically activated porcine oocytes cultured with 10 μM forskolin during the entire IVM period. An increase in cytoplasmic calcium during parthenogenic activation was referred to be sufficient to induce low to moderate extents of oocyte activation events such as cell cycle resumption (Gardner and Evans, 2006). Differences between these two mechanisms, parthenogenesis and cleavage, might explain the above discrepancies.

The inhibitory effect of forskolin during maturation seems to be transiently depending on the supplementation time and dose. However, at higher doses above 10 μM , forskolin presents a toxic effect on COC metabolism (Racowsky, 1985;

Xia *et al.*, 2000). Our findings showed that by reducing forskolin exposure to the initial 22 to 24 h of maturation, GVBD-MI nuclear status became predominant at the end of IVM. Despite of a maturation rate of $24.3 \pm 5.4\%$, only $8.3 \pm 3.3\%$ of inseminated oocytes cleaved. Moreover, forskolin affected not only nuclear progression, but also oocyte cytoplasmic and total areas. Therefore, the transient delay in nuclear meiosis progress of porcine COC matured in the presence of $10 \mu\text{M}$ forskolin identified initially by Racowsky (1985) might be extended to cytoplasmic maturation. However, when this supplementation was reduced to the first 2 h, meiosis progression, oocyte areas and cleavage rates were similar to those of control group. Nonetheless, several authors suggest that forskolin might be applied during oocytes IVM to improve synchronization of nuclear and cytoplasmic maturation, simultaneously increasing the homogeneity and developmental potential of matured oocytes (Racowsky, 1985; Thomas *et al.*, 2004). Further studies are needed to clarify forskolin positive action on oocyte developmental competence for embryo production and quality.

Porcine oocytes present variations in both morphology and amount of cytoplasmic LD during *in vivo* or *in vitro* maturation (Kikuchi *et al.*, 2002). In the present study, we investigated the extent of these variations by measuring LD areas and total area occupied by LD, the fat areas, as well as the relationship of fat area with total oocyte area through the fat oocyte index, before and during porcine oocyte maturation. Moreover, oocytes exhibit distinct colour tones of cytoplasm due to LD content: dark colour and opaqueness in lipid rich ones contrasting with the brightness and transparency of leaner oocytes (Fujihira *et al.*, 2004). Furthermore, Isachenko *et al.* (2003) identified two kinds of LD in porcine oocytes: homogenous, dark looking vesicles and 'gray' looking droplets with electron-lucent streaks. Silva *et al.* (2011) suggested that dark LD may change to gray after lipid utilization. By measuring the fat gray value of porcine oocytes using a Nomarski microscope and ImageJ software, it was possible to distinguish different colour tones of fat areas reflecting alterations in lipid content. Differences in the gray tones might suggest not only alterations of the composition, amount and morphology of LD but also its aggregation or dispersion in the ooplasm. Several authors confirmed that as maturation progresses, the ooplasm can have different shades of gray caused by uneven distribution of organelles, specifically mitochondria and LD, which reside in close proximity (Kikuchi *et al.*, 2002; Ambruosi *et al.*, 2009). Here it was shown that after 44 to 48 h of culture, †10,†12 CLA oocytes were lighter (dark gray = 55.6% and light gray = 44.4%) than those of control and forskolin treatments (dark gray = 100%). Reports confirm the reduction of lipid accumulation in *in vitro* embryos (Pereira *et al.*, 2007 and 2008) or the decrease in lipogenesis of porcine adipose explants (José *et al.*, 2008) after culture with this CLA isomer. Moreover, the presence of †10,†12 CLA during maturation improved bovine oocyte developmental competence (Lapa *et al.*, 2011). Although these latter results at present have not been confirmed for porcine oocytes,

†10,†12 CLA during maturation clearly interfered with LD organization and probably with their composition reflected by a lighter colour tone of their cytoplasm. The mechanism of action of †10,†12 CLA is not clearly known, demanding additional investigation.

Even though forskolin stimulates lipolysis (Men *et al.*, 2006), the fat gray value of forskolin-treated oocytes were not different from control independently of time exposure or meiosis progression. On the contrary, Fu *et al.* (2011) showed that $10 \mu\text{M}$ forskolin effectively reduced the fluorescence intensity of porcine oocytes stained with Nile Red as well as the number of LD during IVM. Data presented here demonstrated no differences in LD areas when this lipolytic agent was present during the entire maturation period or till 22 to 24 h, but fat areas were reduced (exp. 1: control = $12003 \mu\text{m}^2$ v. forskolin = $9769 \mu\text{m}^2$, $P \leq 0.03$; exp. 2: control = $12147 \mu\text{m}^2$ v. forskolin = $9879 \mu\text{m}^2$, $P = 0.006$). It seems that by blocking meiosis progression through the cAMP intracellular level increase (Racowsky, 1985; Xia *et al.*, 2000), forskolin interferes also in the characteristic distribution of LD in ooplasm during the maturation process, resembling the immature oocyte organization. Similar to what happened to the total and cytoplasmic oocyte areas, the fat areas did not grow. Therefore, the oocyte fat indexes were similar to control. The mechanism by which forskolin interfered with LD organization is not well understood but its lipolytic activity in porcine oocytes as well as in embryos has already been identified (Men *et al.*, 2006; Fu *et al.*, 2011). Herein, even when forskolin exposure was only at the initial 2 h of culture, these oocytes at 22 to 24 h had smaller LD and fat areas compared with the control group, although at 44 to 48 h of meiosis these measurements were similar to those of control.

In conclusion, the present study demonstrates that the distribution and morphology of cytoplasmic LD during porcine oocyte maturation can be modified by †10,†12 CLA or forskolin reducing its lipid content in a time-dependent way. However, at the doses tested, whereas †10,†12 CLA did not seem to interfere with oocyte meiosis progression or developmental competence, forskolin should be supplemented for only 2 h to prevent the negative effects on these processes. On the other hand, fat gray value seems a potential tool to evaluate the lipid content of a single porcine oocyte.

Acknowledgements

The 'Fundação para a Ciência e Tecnologia' is greatly acknowledged for funding a grant for E.G. Prates (SFRH/BD/42359/2007).

References

- Ambruosi B, Lacalandra GM, Iorga AI, De Santis T, Mugnier S, Matarrese R, Goudet G and Dell'Aquila ME 2009. Cytoplasmic lipid droplets and mitochondrial distribution in equine oocytes: implications on oocyte maturation, fertilization and developmental competence after ICSI. *Theriogenology* 71, 1093–1104.
- Corino C, Pastorelli G, Douard V, Rossi R, Musella M and Mourot J 2006. L'acid linoléique conjugué en nutrition porcine. *INRA Productions Animales* 19, 39–46.

- Dobrinsky JR, Nagashima H, Pursel VG, Long CR and Johnson LA 1999. Cryopreservation of swine embryos with reduced lipid content. *Theriogenology* 51, 164.
- Fu X, Wu G, Li G, Hou Y, Zhou G, Wang Y and Zhu S 2011. Positive effects of Forskolin (stimulator of lipolysis) treatment on cryosurvival of *in vitro* matured porcine oocytes. *Theriogenology* 75, 268–275.
- Fujihira T, Kinoshita M, Sasaki M, Ohnishi M, Ishikawa H, Ohsumi S and Fukui Y 2004. Comparative studies on lipid analysis and ultrastructure in porcine and southern minke whale (*Balaenoptera bonaerensis*) oocytes. *Journal of Reproduction and Development* 50, 525–532.
- Gardner AJ and Evans JP 2006. Mammalian membrane block to polyspermy: new insights into how mammalian eggs prevent fertilisation by multiple sperm. *Reproduction, Fertility and Development* 18, 53–61.
- Genicot G, Leroy JLMR, Van Soom A and Donnay I 2005. The use of a fluorescent dye, Nile red, to evaluate the lipid content of single mammalian oocytes. *Theriogenology* 63, 1181–1194.
- Gesink LDC, Maclehoose RF and Longnecker MP 2007. Obesity and time to pregnancy. *Human Reproduction* 22, 414–420.
- Gil MA, Abeydeera LR, Day BN, Vazquez JM, Roca J and Martinez EA 2003. Effect of the volume of medium and number of oocytes during *in vitro* fertilization on embryo development in pigs. *Theriogenology* 60, 767–776.
- Isachenko V, Isachenko E, Rahimi G, Mallmann P, Krivokhrchenko A and Nawroth F 2003. Ultra-structure of intracellular lipid vesicles of porcine GV-oocytes after *in vitro* fertilization and parthenogenetic activation. *Anatomia, Histologia, Embryologia* 32, 126–128.
- Kikuchi K, Ekwall H, Tienthai P, Kawai Y, Noguchi J, Kaneko H and Rodriguez-Martinez H 2002. Morphological features of lipid droplet transition during porcine oocyte fertilisation and early embryonic development to blastocyst *in vivo* and *in vitro*. *Zygote* 10, 355–366.
- José AAFBV, Gama MAS and Lanna DDP 2008. Effects of trans-10, cis-12 conjugated linoleic acid on gene expression and lipid metabolism of adipose tissue of growing pigs. *Genetics and Molecular Research* 78, 284–294.
- Laforest MF, Pouliot É, Guéguen L and Richard FJ 2005. Fundamental significance of specific phosphodiesterases in the control of spontaneous meiotic resumption in porcine oocytes. *Molecular Reproduction and Development* 70, 361–372.
- Lapa M, Marques CC, Alves SP, Vasques MI, Baptista MC, Carvalhais I, Silva Pereira M, Horta AEM, Bessa RJB and Pereira RM 2011. Effect of trans-10, cis-12 conjugated linoleic acid on bovine oocyte competence and fatty acid composition. *Reproduction Domestic Animals* 46, 904–910.
- McEvoy TG, Coull GD, Broadbent PJ, Hutchinson JSM and Speake BK 2000. Fatty acid composition of lipids in immature cattle, pig and sheep oocytes with intact zona pellucida. *Journal of Reproduction and Fertility* 118, 163–170.
- Men H, Agca Y, Riley LK and Critser JK 2006. Improved survival of vitrified porcine embryos after partial delipidation through chemically stimulated lipolysis and inhibition of apoptosis. *Theriogenology* 66, 2008–2016.
- Pereira RM and Marques CC 2008. Animal oocyte and embryo cryopreservation. *Cell Tissue Banking* 9, 267–277.
- Pereira RM, Baptista MC, Vasques MI, Horta AEM, Portugal PV, Bessa RJB, Silva JC, Pereira MS and Marques CC 2007. Cryosurvival of bovine blastocysts is enhanced by culture with trans-10 cis-12 conjugated linoleic acid (10t,12c CLA). *Animal Reproduction Science* 98, 293–301.
- Pereira RM, Carvalhais I, Pimenta J, Baptista MC, Vasques MI, Horta AEM, Santos IC, Marques MR, Reis A, Pereira MS and Marques CC 2008. Biopsied and vitrified bovine embryos viability is improved by trans-10 cis-12 conjugated linoleic acid supplementation during *in vitro* embryo culture. *Animal Reproduction Science* 106, 322–332.
- Petters RM and Wells KD 1993. Culture of pig embryos. *Journal of Reproduction and Fertility* 48, 61–73.
- Racowsky C 1985. Effect of forskolin on maintenance of meiotic arrest and stimulation of cumulus expansion, progesterone and cyclic AMP production by pig oocyte-cumulus complexes. *Journal of Reproduction and Fertility* 74, 9–21.
- Silva RC, Bão SN, Jivago JL and Lucci CM 2011. Ultrastructural characterization of porcine oocytes and adjacent follicular cells during follicle development: lipid component evolution. *Theriogenology* 76, 1647–1657.
- Thomas RE, Thompson JG, Armstrong DT and Gilchrist RB 2004. Effect of specific phosphodiesterase isoenzyme inhibitors during *in vitro* maturation of bovine oocytes on meiotic and developmental capacity. *Biology of Reproduction* 71, 1142–1149.
- Walther TC and Farese RV 2009. The life of lipid droplets. *Biochimica et Biophysica Acta* 1791, 459–466.
- Wang H, Rodriguez-Osorio N, Jean Feugang JM, Jung S, Garrison K, Wolgemuth C, Greer L, Crenshaw M and Memili E 2009. Effects of culture media and inhibitors on biology of porcine early embryonic development *in vitro*. *Livestock Science* 121, 102–107.
- Xia GL, Kikuchi K, Noguchi J and Izaiké Y 2000. Short time priming of cumulus–oocyte complexes with FSH and forskolin in the presence of hypoxanthine stimulates cumulus cells to secrete a meiosis-activating substance. *Theriogenology* 53, 1807–1815.
- Zehmer JK, Huang Y, Peng G, Pu J, Anderson RGW and Liu P 2009. A role for lipid droplets in inter-membrane lipid traffic. *Proteomics* 9, 914–921.