

ACKNOWLEDGMENTS

The authors kindly thank Ms. Inácia Ferreira for technical assistance. We would like to thank Mr. José Carlos Garcia dos Santos for help in the location of soil samples on maps, using ArGIS. In the Czech Republic the research was supported by the grant ME 10078 of the Ministry of Education, Youth and Sports.

This communication is a portion of the PhD dissertation of the first author, who is supported by a doctoral scholarship from Fundação para a Ciência e a Tecnologia (SFHR\BD\22086\2005).

LITERATURE CITED

- Ahmad M., Arif M.I. and Denholm I., 2003. High resistance of field populations of the cotton aphid *Aphis gossypii* Glover (Homoptera: Aphididae) to pyrethroid insecticides in Pakistan. *Journal of Economic Entomology*, 96: 875-878.
- Altschul S.F., Gish W., Miller W., Myers E.W. and Lipman D.J., 1990. Basic local alignment search tool. *Journal of Molecular Biology*, 215: 403-410.
- Bedding R.A., 1998. Future possibilities for using entomopathogenic nematodes. *Japanese Journal of Nematology*, 28: 46-60.
- Bedding R.A. and Akhurst R.J., 1975. A simple technique for the detection of insect parasitic nematodes in soil. *Nematologica*, 21: 109-110.
- Chen S.L., Han X. and Moens M., 2003. Effect of chlorpyrifos on infectivity and survival of *Steinernema feltiae*. *Russian Journal of Nematology*, 11: 1-6.
- Felsenstein J., 1985. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 39: 783-791.
- García del Pino F., 2005. Natural occurrence of EPN in Spain. COST 850 WGM on natural occurrence and evolution of EPNs. _eské Bud_jovice, January 14-17, *Abstract*.
- Gaugler R. and Han R., 2002. Production technology. Pp. 289-310. *In: Entomopathogenic Nematology* (Gaugler R., ed.). CABI Publishing, Wallingford, UK.
- Gaugler R. and Han R., 2002. Production technology. Pp. 289-310. *In: Entomopathogenic Nematology* (Gaugler R., ed.). CABI Publishing, Wallingford, UK.
- Georgis R. and Kaya H.K., 1998. Advances in entomopathogenic nematode formulation. Pp. 289-308. *In: Formulation of Microbial Biopesticides, Beneficial Microorganisms, Nematodes and Seed Treatments* (Burgess H.D., ed.) Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Hall T.A., 1999. BioEdit: a user-friendly biological sequences alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium*, 41: 95-98.
- Hominick W.M., Reid A.P., Bohan, D.A. and Briscoe B.R., 1996. Entomopathogenic nematodes: biodiversity, geographical distribution and the Convention on Biological Diversity. *Biocontrol Science and Technology*, 6: 317-331.
- Hunt D., 2007. Overview of taxonomy and systematics. Pp. 27-47. *In: Entomopathogenic Nematodes: Systematics, Phylogeny and Bacterial Symbionts* (Nguyen K.B. and Hunt D.J., eds), Nematology Monographs and Perspectives No. 5, Hunt D.J. and Perry R.N. series editors. Brill, Leiden, The Netherlands.
- Joyce S.A., Burnell A.M. and Powers T.O., 1994. Characterisation of *Heterorhabditis* isolates by PCR amplification of segments of mtDNA and rDNA gene. *Journal of Nematology*, 26: 260-270.
- Kaya H.K. and Koppenhöfer A.M., 1999. Biology and ecology of insecticidal nematodes. Pp. 1-8. *In: Optimal Use of Insecticidal Nematodes in Pest Management* (S. Polavarapu S., ed.). Rutgers University, Rutgers, USA.
- Kaya H.K. and Stock S.P., 1997. Techniques in insect nematology. Pp. 281-324. *In: Manual of Techniques in Insect Pathology* (Lacey L.A., ed.). Academic Press, London, UK.
- Kaya H.K. and Gaugler R., 1993. Entomopathogenic nematodes. *Annual Review of Entomology*, 38: 181-206.
- Khatri-Chhetri H.B., Waeyenberge L., Spiridonov S., Manandhar H.K. and Moens M., 2011. Two new species of *Steinernema* Travassos, 1927 with short infective juveniles from Nepal. *Russian Journal of Nematology*, 19: 53-74.
- Lee M.M., Sicard M., Skeie M. and Stock S.P., 2009. *Steinernema boemarei* n. sp. (Nematoda: Steinernematidae), a new entomopathogenic nematode from southern France. *Systematic Parasitology*, 72: 127-141.
- Liu J., Berry R.E. and Blouin M.S., 1999. Molecular differentiation and phylogeny of entomopathogenic nematodes (Rhabditida: Heterorhabditidae) based on ND4 gene sequences of Mitochondrial DNA. *Journal of Parasitology*, 85: 709-715.
- Liu J. and Berry R.E., 1996. *Steinernema oregonensis* n. sp. (Rhabditida: Steinernematidae) from Oregon, USA. *Fundamental and Applied Nematology*, 19: 375-380.
- Mráček Z. and Webster J., 1993. Survey of Heterorhabditidae and Steinernematidae (Rhabditida, Nematoda) in Western Canada. *Journal of Nematology*, 25: 710-717.
- Mráček Z., Bečvář S., Kindlmann P. and Jersáková J., 2005. Habitat preference for entomopathogenic nematodes, their insect hosts and new faunistic records for the Czech Republic. *Biological Control*, 34: 27-37.
- Mráček Z., Bečvář S. and Kindlmann P., 1999. Survey of entomopathogenic nematodes from the families Steinernematidae and Heterorhabditidae (Nematoda: Rhabditida) in the Czech republic. *Folia Parasitologica*, 46: 145-148.
- Nadler S.A., 2002. Species delimitation and nematode biodiversity: phylogenies rule. *Nematology*, 4: 615-625.
- Nadler S.A., Bolotin E. and Stock S.P., 2006. Phylogenetic relationships of *Steinernema* Travassos, 1927 (Nematoda: Cephalobina: Steinernematidae) based on nuclear, mitochondrial and morphological data. *Systematic Parasitology*, 63: 161-181.
- Nei M. and Kumar S., 2000. *Molecular Evolution and Phylogenetics*. Oxford University Press, New York, USA, 333 pp.
- Nguyen K.B., Hunt D.J. and Mráček Z., 2007. Steinernematidae: species description. Pp. 121-609. *In: Entomopathogenic nematodes: Systematics, Phylogeny and Bacterial symbionts* (Nguyen K.B. and Hunt D.J., eds), Nematology Monographs and Perspectives No. 5, Hunt D.J. and Perry R.N. series editors. Brill, Leiden, The Netherlands.
- Nguyen K.B. and Smart Jr. G.C., 1995. Scanning electron microscope studies of *Steinernema glaseri* (Nematoda: Steinernematidae). *Nematologica*, 41: 183-190.