

- Qui L., Fang Y., Zhou Y., Pang Y. and Nguyen K.B., 2004. *Steinernema guangdongense* sp. n. (Nematoda: Steinernematidae), a new entomopathogenic nematode from southern China with a note on *S. serratum* (*nomen nudum*). *Zootaxa*, 704: 1-20.
- Rosa J.S. and Simões N., 2004. Evaluation of twenty-eight strains of *Heterorhabditis bacteriophora* isolated in Azores for biocontrol of the armyworm, *Pseudaletia unipuncta* (Lepidoptera: Noctuidae). *Biological Control*, 29: 409-417.
- Rosa J.S., Bonifassi E., Amaral J., Lacey L.A., Simões N. and Laumond C., 2000. Natural occurrence of entomopathogenic nematodes (Rhabditida: *Steinernema*, *Heterorhabditis*) in Azores. *Journal of Nematology*, 32: 215-222.
- Seinhorst J.W., 1959. A rapid method for the transfer of nematodes from fixative to anhydrous glycerin. *Nematologica*, 4: 67-69.
- Shapiro-Ilan D.I., Gouge D.H. and Koppenhofer A.M., 2002. Factors affecting commercial success: case studies in cotton, turf and citrus. Pp. 333-356. In: Entomopathogenic Nematology (Gaugler R., ed.). CABI Publishing, Wallingford, UK.
- Simões N., Laumond C. and Bonifassi E., 1994. Effectiveness of *Steinernema* spp. and *Heterorhabditis bacteriophora* against *Popillia japonica* in Azores. *Journal of Nematology*, 25: 480-485.
- Southey J.F., 1970. *Laboratory methods for work with plant and soil nematodes*. Ministry of Agriculture, Fisheries and Food, Her Majesty's Stationery Office, London, UK.
- Spiridonov S., Krasomil-Osterfeld K. and Moens M., 2004. *Steinernema jolietii* sp. n. (Rhabditida: Steinernematidae), a new entomopathogenic nematode from the American Midwest. *Russian Journal of Nematology*, 12: 85-95.
- Steiner W.A., 1996. Distribution of entomopathogenic nematodes in the Swiss Alps. *Révue Suisse de Zoologie*, 103: 439-452.
- Stock S.P., Campbell J.F. and Nadler S.A., 2001. Phylogeny of *Steinernema* Travassos, 1927 (Cephalobina: Steinernematidae) inferred from ribosomal DNA sequences and morphological characters. *Journal of Parasitology*, 87: 877-889.
- Stock P., Mráček Z. and Webster J.M., 2000. Morphological variation among allopatric populations of *Steinernema kraussei* (Steiner, 1923) (Rhabditida: Steinernematidae). *Nematology*, 2: 143-152.
- Sturhan D., Spiridonov S.E. and Mráček Z., 2005. *Steinernema silvaticum* sp. n. (Rhabditida: Steinernematidae), a new entomopathogenic nematode from Europe. *Nematology*, 7: 227-241.
- Sturhan D., 1999. Prevalence and habitat specificity of entomopathogenic nematodes in Germany. Pp. 123-132. In: COST 819, Entomopathogenic nematodes: application and persistence of entomopathogenic nematodes (Gwynn R.L., Smith P.H., Griffin C., Ehlers R.U., Boemare N. and Masson J.P., eds). European Commission, Luxembourg.
- Tamura K., Dudley J., Nei M. and Kumar S., 2007. MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Molecular Biology and Evolution*, 24: 1596-1599.
- Tarasco E., Clausi M., Rappazzo G., Vinciguerra M., Longo A. and Triggiani O., 2009. Could be Italy considered a favorite place in Europe for EPN biodiversity? IOBC/wprs Bulletin Vol. 45: 387-389. Working Group "Insect Pathogens and Insect Parasitic Nematodes". Proceedings of the 12th European Meeting at Pamplona, Spain, June 22-26, 2009 (Ehlers R.-U., Crickmore N., Enkerli J., Glazer I., Lopez Ferber M. and C. Tkaczuk C., eds). Bern, Switzerland.
- Valadas V., Vieira P., Oliveira S. and Mota M., 2009. First report of the genus *Heterorhabditis* (Nematoda: Heterorhabditidae) from continental Portugal. *Helminthology*, 46: 45-48.
- Valadas V., Boyle S., Vieira P., Kakouli-Duarte T. and Mota M., 2007. First report of an entomopathogenic nematode from continental Portugal. *Helminthology*, 44: 226-229.
- Weiser J., 1955. *Neoaplectana carpocapsae* n. sp. (Anguillulata, Steinernematidae) nový cizopasník housenek obalece jablecneho, *Carpocapsa pomonella* L. *Vestník Československé Zoologické Společnosti*, 19: 44-52.
- White G.F., 1929. A method for obtaining infective juveniles from cultures. *Science*, 66: 302-303.