

# ***Bactrocera oleae* (Diptera: Tephritidae) in Iran: An invasion from the Middle West**

Sadrollah Ramezani 1, Imen Blibech 2, Fernando Trindade Rei 3, Barbara van Asch 4 and Luís Teixeira da Costa 3

1 Department of Horticulture Science, College of Agriculture, Tarbiat Modares University, Tehran, Iran; e-mail: sramezani@gmail.com

2 Laboratory of Genetic Resources and Protection of the Olive Tree, Olive Tree Institute, Sousse Station, Sousse, Tunisia; e-mail: imenblibech@gmail.com

3 Instituto de Ciências Agrárias e Ambientais Mediterrânicas (ICAAM), Universidade de Évora, Núcleo da Mitra, 7002-774 Évora,

Portugal, e-mails: frei@uevora.pt; luisteixeiracosta@gmail.com

4 Department of Genetics, Faculty of Agrisciences, Stellenbosch University, 7602 Stellenbosch, South Africa; e-mail: barbaravanasch@gmail.com

**Key words.** Diptera, Tephritidae, *Bactrocera oleae*, invasion, Iran, genetics, mitochondrial DNA, haplotype, olive, olive fly

## **Abstract.**

Despite an age-old tradition of olive growing and its geographical location, Iran was apparently free of the olive fly, *Bactrocera oleae* (Rossi, 1790) (Diptera: Tephritidae), the major worldwide olive tree pest, until the last decade. However, this situation has changed radically since *B. oleae* was first reported in 2004, and the olive fly is now Iran's most important pest of olive trees, and a very serious threat to olive production in the Northern and more humid parts of the country. Genetic analyses of mitochondrial markers of *B. oleae* collected in the traditional olive growing area in Northwestern Iran were used to determine the origin of this pest. The results indicate it was most likely imported on olive trees from the Central Mediterranean area.