

Rhizobia of chickpea from southern Portugal: symbiotic efficiency and genetic diversity



Print E-mail Add to Marked List Save to EndNote Web
Save to EndNote, RefMan, ProCite more options

Author(s): Laranjo M, Rodrigues R, Alho L, Oliveira S

Source: JOURNAL OF APPLIED MICROBIOLOGY **Volume:** 90 **Issue:** 4 **Pages:** 662-667 **Published:** APR 2001

Times Cited: 10 **References:** 21 Citation Map

Abstract: Aims: In order to evaluate differences between chickpea rhizobial populations from three geographical areas in southern Portugal (Beja, Elvas and Evora), isolates from the three regions were obtained and analysed.

Methods and Results: The genetic characterization of the isolates was done by plasmid profiles and restriction analysis of the nif H gene. Symbiotic efficiency of the isolates was also determined. Relationships between geographical origin, symbiotic efficiency and molecular characteristics were established. Beja soil revealed a larger rhizobia population as well as the presence of some of the isolates with higher symbiotic efficiency values. Isolates with a single plasmid showed a significantly higher symbiotic efficiency.

Conclusions: Genetic and phenotypic differences were detected between the natural rhizobial populations from the three locations.

Significance and Impact of the Study: The different yield potential with cultivars of chickpea usually obtained in the three regions of southern Portugal could be due to their different natural rhizobial populations.

Document Type: Article

Language: English

KeyWords Plus: NITROGEN-FIXATION; NIFH GENES; PLASMID; TRIFOLIUM; MELILOTI; STRAINS

Reprint Address: Oliveira, S (reprint author), Univ Evora, Nucleo Mitra, Dept Biol, Apartado 94, P-7002554 Evora, Portugal

Addresses:

1. Univ Evora, Nucleo Mitra, Dept Biol, P-7002554 Evora, Portugal
2. Univ Evora, ICAM, Lab Microbiol Solo, Evora, Portugal

Publisher: BLACKWELL SCIENCE LTD, P O BOX 88, OSNEY MEAD, OXFORD OX2 0NE, OXON, ENGLAND