

# EXISTENCE OF SOLUTION FOR FUNCTIONAL COUPLED SYSTEMS WITH FULL NONLINEAR TERMS AND APPLICATIONS TO A COUPLED MASS-SPRING MODEL

FELIZ MINHÓS AND ROBERT DE SOUSA

*(Communicated by Rodrigo L. Pouso)*

*Abstract.* In this paper we consider some boundary value problems composed by coupled systems of second order differential equations with full nonlinearities and general functional boundary conditions verifying some monotone assumptions.

The arguments apply lower and upper solutions method and fixed point theory. Due to an adequate auxiliary problem, including a convenient truncature, there is no need of sign, bound, monotonicity or other growth assumptions on the nonlinearities, besides the Nagumo condition.

An application to a coupled mass-spring system with functional behavior at the final instant is shown.

## 1. Introduction

In this paper we consider the boundary value problem composed by the coupled system of the second order differential equations with full nonlinearities

$$\begin{cases} u''(t) = f(t, u(t), v(t), u'(t), v'(t)), & t \in [a, b], \\ v''(t) = h(t, u(t), v(t), u'(t), v'(t)) \end{cases} \quad (1.1)$$

with  $f, h : [a, b] \times \mathbb{R}^4 \rightarrow \mathbb{R}$  continuous functions, and the functional boundary conditions

$$\begin{cases} u(a) = v(a) = 0 \\ L_1(u, u(b), u'(b)) = 0 \\ L_2(v, v(b), v'(b)) = 0, \end{cases} \quad (1.2)$$

where  $L_1, L_2 : C[a, b] \times \mathbb{R}^2 \rightarrow \mathbb{R}$  are continuous functions verifying some monotone assumptions.

Ordinary differential systems have been studied by many authors, like, for instance, [1, 5, 11, 12, 13, 14, 15, 17, 18, 20, 21, 24, 28]. In particular, coupled second order ordinary differential systems can be applied to several real phenomena, such as, Lokta-Volterra models, reaction diffusion processes, prey-predator or other interaction

---

*Mathematics subject classification (2010):* 34B15, 34B27, 34L30, 74H20.

*Keywords and phrases:* Coupled systems, functional boundary conditions, Green's functions, Schauder's fixed-point theorem, coupled mass-spring system.

systems, Sturm-Liouville problems, mathematical biology, chemical systems (see, for example, [2, 3, 4, 16, 19, 26] and the references therein).

In [23] the authors study the existence of solutions for the nonlinear second order coupled system

$$\begin{cases} -u''(t) = f_1(t, v(t)) \\ -v''(t) = f_2(t, v(t)), \end{cases} \quad t \in [0, 1],$$

with  $f_1, f_2 : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$  continuous functions, together with the nonlinear boundary conditions

$$\begin{aligned} \varphi(u(0), v(0), u'(0), v'(0), u'(1), v'(1)) &= (0, 0), \\ \psi(u(0), v(0)) + (u(1), v(1)) &= (0, 0), \end{aligned}$$

where  $\varphi : \mathbb{R}^6 \rightarrow \mathbb{R}^2$  and  $\psi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  are continuous functions.

In [7] it is provided some growth conditions on the nonnegative nonlinearities of the system

$$\begin{cases} -x''(t) = f_1(t, x(t), y(t)) \\ -y''(t) = f_2(t, x(t), y(t)), t \in (0, 1), \\ x(0) = y(0) = 0, \\ x(1) = \alpha[y], \\ y(1) = \beta[x], \end{cases}$$

where  $f_1, f_2 : (0, 1) \times [0, +\infty)^2 \rightarrow [0, +\infty)$  are continuous and may be singular at  $t = 0, 1$ , and  $\alpha[x], \beta[x]$  are bounded linear functionals on  $C[0, 1]$  given by

$$\alpha[y] = \int_0^1 y(t) dA(t), \quad \beta[x] = \int_0^1 x(t) dB(t),$$

involving Stieltjes integrals, and  $A, B$  are functions of bounded variation with positive measures.

Motivated by these works we consider the second order coupled fully differential equations (1.1) together with the functional boundary conditions (1.2). To the best of our knowledge, it is the first time where these coupled differential systems embrace functional boundary conditions. Remark that, the functional dependence includes and generalizes the classical boundary conditions such as separated, multi-point, nonlocal, integro-differential, with maximum or minimum arguments, . . . More details on such conditions and their application potentialities can be seen, for instance, in [6, 9, 10, 22, 25] and the references therein. Our main result is applied to a coupled mass-spring systems subject to a new type of global boundary data.

The arguments in this paper follow lower and upper solutions method and fixed point theory. Therefore, the main result is an existence and localization theorem, as it provides not only the existence of solution, but a strip where the solution varies, as well. Due to an adequate auxiliary problem, including a convenient truncature, there is no need of sign, bound, monotonicity or other growth assumptions on the nonlinearities, besides the Nagumo condition.

The paper is organized as it follows: Section 2 contains the space framework, lower an upper solutions definition and some *a priori* estimates on the first derivative of the unknown functions. In sections 3 and 4 we present an existence and localization result and an example to show the applicability of the main theorem. An application to a real phenomenon is shown in the last section: a coupled mass-spring system together with functional behavior at the final instant.

## 2. Definitions and preliminaries

Let  $E = C^1[a, b]$  be the Banach space equipped with the norm  $\|\cdot\|_{C^1}$ , defined by

$$\|w\|_{C^1} := \max \{ \|w\|, \|w'\| \},$$

where

$$\|y\| := \max_{t \in [a, b]} |y(t)|$$

and  $E^2 = (C^1[a, b])^2$  with the norm

$$\|(u, v)\|_{E^2} = \|u\|_{C^1} + \|v\|_{C^1}.$$

Forward in this work, we consider the following assumption

- (A) The functions  $L_1, L_2 : C[a, b] \times \mathbb{R}^2 \rightarrow \mathbb{R}$  are continuous, nonincreasing in the first variable and nondecreasing in the second one.

To apply lower and upper solutions method we consider next definition:

**DEFINITION 1.** A pair of functions  $(\alpha_1, \alpha_2) \in (C^2[a, b])^2$  is a coupled lower solution of problem (1.1), (1.2) if

$$\begin{aligned} \alpha_1''(t) &\leq f(t, \alpha_1(t), \alpha_2(t), \alpha_1'(t), \alpha_2'(t)) \\ \alpha_2''(t) &\leq h(t, \alpha_1(t), \alpha_2(t), \alpha_1'(t), \alpha_2'(t)) \\ \alpha_1(a) &\leq 0 \\ \alpha_2(a) &\leq 0 \\ L_1(\alpha_1, \alpha_1(b), \alpha_1'(b)) &\geq 0 \\ L_2(\alpha_2, \alpha_2(b), \alpha_2'(b)) &\geq 0. \end{aligned}$$

A pair of functions  $(\beta_1, \beta_2) \in (C^2[a, b])^2$  is a coupled upper solution of problem (1.1), (1.2) if it verifies the reverse inequalities.

A Nagumo-type condition is useful to obtain *a priori* bounds on the first derivatives of the unknown functions:

DEFINITION 2. Let  $\alpha_1(t), \beta_1(t), \alpha_2(t)$  and  $\beta_2(t)$  be continuous functions such that

$$\alpha_1(t) \leq \beta_1(t), \quad \alpha_2(t) \leq \beta_2(t), \quad \forall t \in [a, b].$$

The continuous functions  $f, h : [a, b] \times \mathbb{R}^4 \rightarrow \mathbb{R}$  satisfy a Nagumo type condition relative to intervals  $[\alpha_1(t), \beta_1(t)]$  and  $[\alpha_2(t), \beta_2(t)]$ , if, there are  $N_1 > r_1$ ,  $N_2 > r_2$ , with

$$r_1 := \max \left\{ \frac{\beta_1(b) - \alpha_1(a)}{b-a}, \frac{\beta_1(a) - \alpha_1(b)}{b-a} \right\}, \quad (2.1)$$

$$r_2 := \max \left\{ \frac{\beta_2(b) - \alpha_2(a)}{b-a}, \frac{\beta_2(a) - \alpha_2(b)}{b-a} \right\}, \quad (2.2)$$

and continuous positive functions  $\varphi, \psi : [0, +\infty) \rightarrow (0, +\infty)$ , such that

$$|f(t, x, y, z, w)| \leq \varphi(|z|), \quad (2.3)$$

and

$$|h(t, x, y, z, w)| \leq \psi(|w|), \quad (2.4)$$

for

$$\alpha_1(t) \leq x \leq \beta_1(t), \quad \alpha_2(t) \leq y \leq \beta_2(t), \quad \forall t \in [a, b], \quad (2.5)$$

verifying

$$\int_{r_1}^{N_1} \frac{ds}{\varphi(s)} > b-a, \quad \int_{r_2}^{N_2} \frac{ds}{\psi(s)} > b-a. \quad (2.6)$$

LEMMA 1. Let  $f, h : [a, b] \times \mathbb{R}^4 \rightarrow \mathbb{R}$  be continuous functions satisfying a Nagumo type condition relative to intervals  $[\alpha_1(t), \beta_1(t)]$  and  $[\alpha_2(t), \beta_2(t)]$ .

Then for every solution  $(u, v) \in (C^2[a, b])^2$  verifying (2.5), there are  $N_1, N_2 > 0$ , given by (2.6), such that

$$\|u'\| \leq N_1 \text{ and } \|v'\| \leq N_2. \quad (2.7)$$

*Proof.* Let  $(u(t), v(t))$  be a solution of (1.1) satisfying (2.5).

By Lagrange Theorem, there are  $t_0, t_1 \in [a, b]$  such that

$$u'(t_0) = \frac{u(b) - u(a)}{b-a} \text{ and } v'(t_1) = \frac{v(b) - v(a)}{b-a}.$$

Suppose, by contradiction, that  $|u'(t)| > r_1$ ,  $\forall t \in [a, b]$ , with  $r_1$  given by (2.1). If  $u'(t) > r_1$ ,  $\forall t \in [a, b]$ , by (2.5) we obtain the following contradiction with (2.1):

$$u'(t_0) = \frac{u(b) - u(a)}{b-a} \leq \frac{\beta_1(b) - \alpha_1(a)}{b-a} \leq r_1.$$

If  $u'(t) < -r_1$ ,  $\forall t \in [a, b]$ , the contradiction is similar.

In the case where  $|u'(t)| \leq r_1$ ,  $\forall t \in [a, b]$ , the proof will be finished.

So, assume that there are  $t_2, t_3 \in [a, b]$  such that  $t_2 < t_3$ , and

$$u'(t_2) < r_1 \text{ and } u'(t_3) > r_1.$$

By continuity, there is  $t_4 \in [t_2, t_3]$  such that

$$u'(t_4) = r_1 \text{ and } u'(t_3) > r_1, \forall t \in ]t_4, t_3].$$

So, by a convenient change of variable, by (2.3), (2.4), and (2.6), we obtain

$$\begin{aligned} \int_{u'(t_4)}^{u'(t_3)} \frac{ds}{\varphi(|s|)} &= \int_{t_4}^{t_3} \frac{u''(t)}{\varphi(|u'(t)|)} dt \leqslant \int_a^b \frac{|u''(t)|}{\varphi(|u'(t)|)} dt \\ &= \int_a^b \frac{|f(t, u(t), v(t), u'(t), v'(t))|}{\varphi(|u'(t)|)} dt \leqslant b - a < \int_{r_1}^{N_1} \frac{ds}{\varphi(|s|)}. \end{aligned}$$

Therefore  $u'(t_3) < N_1$ , and, as  $t_3$  is taken arbitrarily,  $u'(t_3) < N_1$ , for values of  $t$  where  $u'(t) > r_1$ .

If  $t_2 > t_3$ , the technique is analogous for  $t_4 \in [t_3, t_2]$ .

The same conclusion can be achieved if there are  $t_2, t_3 \in [a, b]$  such that

$$u'(t_2) > -r_1 \text{ and } u'(t_3) < -r_1.$$

Therefore  $\|u'\| \leqslant N_1$  and, by similar arguments, it can be proved that  $\|v'\| \leqslant N_2$ .  $\square$

For the reader's convenience we present Schauder's fixed point theorem:

**THEOREM 1.** ([27]) *Let  $Y$  be a nonempty, closed, bounded and convex subset of a Banach space  $X$ , and suppose that  $P : Y \rightarrow Y$  is a compact operator. Then  $P$  has at least one fixed point in  $Y$ .*

### 3. Main result

Along this work we denote  $(a, b) \leqslant (c, d)$  meaning that  $a \leqslant c$  and  $b \leqslant d$ , for  $a, b, c, d \in \mathbb{R}$ .

**THEOREM 2.** *Let  $f, h : [a, b] \times \mathbb{R}^4 \rightarrow \mathbb{R}$  be continuous functions, and assume that hypothesis (A) holds. If there are coupled lower and upper solutions of (1.1), (1.2),  $(\alpha_1, \alpha_2)$  and  $(\beta_1, \beta_2)$ , respectively, according Definition 1, such that*

$$(\alpha_1(t), \alpha_2(t)) \leqslant (\beta_1(t), \beta_2(t)), \quad (3.1)$$

and  $f$  and  $h$  verify the Nagumo conditions relative to intervals  $[\alpha_1(t), \beta_1(t)]$  and  $[\alpha_2(t), \beta_2(t)]$ , then there is at least a pair  $(u(t), v(t)) \in (C^2[a, b], \mathbb{R})^2$  solution of (1.1), (1.2) and, moreover,

$$\alpha_1(t) \leqslant u(t) \leqslant \beta_1(t), \quad \alpha_2(t) \leqslant v(t) \leqslant \beta_2(t), \quad \forall t \in [a, b].$$

and

$$\|u'\| \leq N_1 \text{ and } \|v'\| \leq N_2,$$

with  $N_1$  and  $N_2$  given by Lemma 1.

*Proof.* Consider the auxiliary functions  $F(t, x, y, z, w) := F$  defined as

- $f(t, \beta_1(t), \beta_2(t), \beta'_1(t), \beta'_2(t)) - \frac{x-\beta_1(t)}{1+|x-\beta_1(t)|} - \frac{y-\beta_2(t)}{1+|y-\beta_2(t)|}$  if  $x > \beta_1(t)$ ,  $y > \beta_2(t)$ ,
- $f(t, u(t), v(t), u'(t), v'(t)) - \frac{y-\beta_2(t)}{1+|y-\beta_2(t)|}$  if  $\alpha_1(t) \leq x \leq \beta_1(t)$ ,  $y > \beta_2(t)$ ,
- $f(t, \alpha_1(t), \alpha_2(t), \alpha'_1(t), \alpha'_2(t)) - \frac{x-\alpha_1(t)}{1+|x-\alpha_1(t)|} + \frac{y-\beta_2(t)}{1+|y-\beta_2(t)|}$  if  $x < \alpha_1(t)$ ,  $y > \beta_2(t)$ ,
- $f(t, \beta_1(t), \beta_2(t), \beta'_1(t), \beta'_2(t)) - \frac{x-\beta_1(t)}{1+|x-\beta_1(t)|}$ , if  $x > \beta_1(t)$ ,  $\alpha_2(t) \leq y \leq \beta_2(t)$ ,
- $f(t, u(t), v(t), u'(t), v'(t))$  if  $\alpha_1(t) \leq x \leq \beta_1(t)$ ,  $\alpha_2(t) \leq y \leq \beta_2(t)$ ,
- $f(t, \alpha_1(t), \alpha_2(t), \alpha'_1(t), \alpha'_2(t)) - \frac{x-\alpha_1(t)}{1+|x-\alpha_1(t)|}$  if  $x < \alpha_1(t)$ ,  $\alpha_2(t) \leq y \leq \beta_2(t)$ ,
- $f(t, \beta_1(t), \beta_2(t), \beta'_1(t), \beta'_2(t)) - \frac{x-\beta_1(t)}{1+|x-\beta_1(t)|} + \frac{y-\alpha_2(t)}{1+|y-\alpha_2(t)|}$  if  $x > \beta_1(t)$ ,  $y < \alpha_2(t)$ ,
- $f(t, u(t), v(t), u'(t), v'(t)) - \frac{y-\alpha_2(t)}{1+|y-\alpha_2(t)|}$  if  $\alpha_1(t) \leq x \leq \beta_1(t)$ ,  $y < \alpha_2(t)$ ,
- $f(t, \alpha_1(t), \alpha_2(t), \alpha'_1(t), \alpha'_2(t)) - \frac{x-\alpha_1(t)}{1+|x-\alpha_1(t)|} - \frac{y-\alpha_2(t)}{1+|y-\alpha_2(t)|}$  if  $x < \alpha_1(t)$ ,  $y < \alpha_2(t)$ ,

and  $H(t, x, y, z, w) := H$  given by

- $h(t, \beta_1(t), \beta_2(t), \beta'_1(t), \beta'_2(t)) - \frac{x-\beta_1(t)}{1+|x-\beta_1(t)|} - \frac{y-\beta_2(t)}{1+|y-\beta_2(t)|}$  if  $x > \beta_1(t)$ ,  $y > \beta_2(t)$ ,
- $h(t, u(t), v(t), u'(t), v'(t)) - \frac{y-\beta_2(t)}{1+|y-\beta_2(t)|}$  if  $\alpha_1(t) \leq x \leq \beta_1(t)$ ,  $y > \beta_2(t)$ ,
- $h(t, \alpha_1(t), \alpha_2(t), \alpha'_1(t), \alpha'_2(t)) - \frac{x-\alpha_1(t)}{1+|x-\alpha_1(t)|} + \frac{y-\beta_2(t)}{1+|y-\beta_2(t)|}$  if  $x < \alpha_1(t)$ ,  $y > \beta_2(t)$ ,
- $h(t, \beta_1(t), \beta_2(t), \beta'_1(t), \beta'_2(t)) - \frac{x-\beta_1(t)}{1+|x-\beta_1(t)|}$ , if  $x > \beta_1(t)$ ,  $\alpha_2(t) \leq y \leq \beta_2(t)$ ,
- $h(t, u(t), v(t), u'(t), v'(t))$  if  $\alpha_1(t) \leq x \leq \beta_1(t)$ ,  $\alpha_2(t) \leq y \leq \beta_2(t)$ .
- $h(t, \alpha_1(t), \alpha_2(t), \alpha'_1(t), \alpha'_2(t)) - \frac{x-\alpha_1(t)}{1+|x-\alpha_1(t)|}$  if  $x < \alpha_1(t)$ ,  $\alpha_2(t) \leq y \leq \beta_2(t)$ ,
- $h(t, \beta_1(t), \beta_2(t), \beta'_1(t), \beta'_2(t)) - \frac{x-\beta_1(t)}{1+|x-\beta_1(t)|} + \frac{y-\alpha_2(t)}{1+|y-\alpha_2(t)|}$  if  $x > \beta_1(t)$ ,  $y < \alpha_2(t)$ ,
- $h(t, u(t), v(t), u'(t), v'(t)) - \frac{y-\alpha_2(t)}{1+|y-\alpha_2(t)|}$  if  $\alpha_1(t) \leq x \leq \beta_1(t)$ ,  $y < \alpha_2(t)$ ,
- $h(t, \alpha_1(t), \alpha_2(t), \alpha'_1(t), \alpha'_2(t)) - \frac{x-\alpha_1(t)}{1+|x-\alpha_1(t)|} - \frac{y-\alpha_2(t)}{1+|y-\alpha_2(t)|}$  if  $x < \alpha_1(t)$ ,  $y < \alpha_2(t)$ ,

and the auxiliary problem

$$\begin{cases} u''(t) = F(t, u(t), v(t), u'(t), v'(t)) \\ v''(t) = H(t, u(t), v(t), u'(t), v'(t)) \\ u(a) = v(a) = 0 \\ u(b) = \delta_1(b, u(b) + L_1(u, u(b), u'(b))) \\ v(b) = \delta_2(b, v(b) + L_2(v, v(b), v'(b))), \end{cases} \quad (3.2)$$

where, for each  $i = 1, 2$ ,

$$\delta_i(t, w) = \begin{cases} \beta_i(t), & w > \beta_i(t) \\ w, & \alpha_i(t) \leq w \leq \beta_i(t) \\ \alpha_i(t), & w < \alpha_i(t). \end{cases} \quad (3.3)$$

*Claim 1:* Solutions of problem (3.2) can be written as

$$\begin{aligned} u(t) &= \frac{t-a}{b-a} \delta_1(b, u(b) + L_1(u, u(b), u'(b))) \\ &\quad + \int_a^b G(t, s) F(s, u(s), v(s), u'(s), v'(s)) ds \\ v(t) &= \frac{t-a}{b-a} \delta_2(b, v(b) + L_2(v, v(b), v'(b))) \\ &\quad + \int_a^b G(t, s) H(s, u(s), v(s), u'(s), v'(s)) ds, \end{aligned}$$

where

$$G(t, s) = \frac{1}{b-a} \begin{cases} (a-s)(b-t), & a \leq t \leq s \leq b \\ (a-t)(b-s), & a \leq s \leq t \leq b. \end{cases} \quad (3.4)$$

In fact, for the equation  $u''(t) = F(t)$ , the solution is,

$$u(t) = At + B + \int_a^t (t-s) F(s) ds, \quad (3.5)$$

for some  $A, B \in \mathbb{R}$ .

By the boundary conditions, it follows that,

$$\begin{aligned} A &= \frac{1}{b-a} \delta_1(b, u(b) + L_1(u, u(b), u'(b))) - \frac{1}{b-a} \int_a^b (b-s) F(s) ds \\ B &= -\frac{a}{b-a} \delta_1(b, u(b) + L_1(u, u(b), u'(b))) + \frac{a}{b-a} \int_a^b (b-s) F(s) ds. \end{aligned} \quad (3.6)$$

By (3.5) and (3.6), then

$$\begin{aligned}
u(t) &= \frac{t-a}{b-a} \delta_1(b, u(b) + L_1(u, u(b), u'(b))) \\
&\quad - \frac{t-a}{b-a} \int_a^b (b-s) F(s) ds + \int_a^t (t-s) F(s) ds \\
&= \frac{t-a}{b-a} \delta_1(b, u(b) + L_1(u, u(b), u'(b))) \\
&\quad + \int_a^t \left( \frac{(a-t)(b-s)}{b-a} + t-s \right) F(s) ds + \int_t^b \frac{(a-t)(b-s)}{b-a} F(s) ds \\
&= \frac{t-a}{b-a} \delta_1(b, u(b) + L_1(u, u(b), u'(b))) \\
&\quad + \int_a^t \left( \frac{(a-s)(b-t)}{b-a} + t-s \right) F(s) ds + \int_t^b \frac{(a-t)(b-s)}{b-a} F(s) ds \\
&= \frac{t-a}{b-a} \delta_1(b, u(b) + L_1(u, u(b), u'(b))) + \int_a^t G(t, s) F(s) ds,
\end{aligned}$$

with  $G(t, s)$  given by (3.4).

The integral form of  $v(t)$  can be achieved by the same arguments.

*Claim 2:* Every solution  $(u, v)$  of (3.2) satisfies

$$\|u'\| \leq N_1 \text{ and } \|v'\| \leq N_2,$$

with  $N_1$  and  $N_2$  given by Lemma 1.

This claim is a direct consequence of Lemma 1, as  $F(t, x, y, z, w)$  and  $H(t, x, y, z, w)$  are defined on bounded arguments of  $x$  and  $y$ .

Define the operators  $T_1 : (C^1[a, b])^2 \rightarrow C^1[a, b]$  and  $T_2 : (C^1[a, b])^2 \rightarrow C^1[a, b]$  such that

$$\begin{cases} T_1(u, v)(t) = \frac{t-a}{b-a} \delta_1(b, u(b) + L_1(u, u(b), u'(b))) \\ \quad + \int_a^b G(t, s) F(s, u(s), v(s), u'(s), v'(s)) ds \\ T_2(u, v)(t) = \frac{t-a}{b-a} \delta_2(b, v(b) + L_2(v, v(b), v'(b))) \\ \quad + \int_a^b G(t, s) H(s, u(s), v(s), u'(s), v'(s)) ds, \end{cases} \quad (3.7)$$

where  $G(t, s)$  is given by (3.4), and  $T : (C^1[a, b])^2 \rightarrow (C^1[a, b])^2$  by

$$T(u, v)(t) = (T_1(u, v)(t), T_2(u, v)(t)). \quad (3.8)$$

By Claim 1, fixed points of the operator  $T := (T_1, T_2)$  are solutions of problem (3.2).

*Claim 3:* The operator  $T$ , given by (3.8) has a fixed point  $(u_0, v_0)$ .

In order to apply Theorem 1, we will prove the following steps for operator  $T_1(u, v)$ . The proof for the operator  $T_2(u, v)$  is analogous.

(i)  $T_1 : (C^1[a, b])^2 \rightarrow C^1[a, b]$  is well defined.

The function  $F$  is bounded and the Green function  $G(t, s)$  is continuous in  $[a, b]^2$ , then the operator  $T_1(u, v)$  is continuous. Moreover, as  $\frac{\partial G}{\partial t}(t, s)$  is bounded in  $[a, b]^2$  and

$$\begin{aligned} (T_1(u, v))'(t) &= \frac{1}{b-a} \delta_1(b, u(b) + L_1(u, u(b), u'(b))) \\ &\quad + \int_a^b \frac{\partial G}{\partial t}(t, s) F(s, u(s), v(s), u'(s), v'(s)) ds, \end{aligned}$$

with

$$\frac{\partial G}{\partial t}(t, s) = \frac{1}{b-a} \begin{cases} s-a, & a \leq t \leq s \leq b \\ s-b, & a \leq s \leq t \leq b, \end{cases}$$

verifying

$$\left| \frac{\partial G}{\partial t}(t, s) \right| \leq 1, \quad \forall (t, s) \in [a, b]^2,$$

therefore,  $(T_1(u, v))'$  is continuous on  $[a, b]$ . So,  $T_1 \in C^1[a, b]$ .

(ii)  *$TB$  is uniformly bounded, for  $B$  a bounded set in  $(C^1[a, b])^2$ .*

Let  $B$  be a bounded set of  $(C^1[a, b])^2$ . Then there exists  $K > 0$  such that

$$\|(u, v)\|_{E^2} = \|u\|_{C^1} + \|v\|_{C^1} \leq K, \quad \forall (u, v) \in B.$$

By (3.3), and taking into account that  $F$  and  $H$  are bounded, then there are  $M_1, M_2, M_3 > 0$  such that

$$\begin{aligned} \delta_1 &\leq \max \{\|\alpha_1\|, \|\beta_1\|\} := M_1, \\ \int_a^b \max_{t \in [a, b]} |G(t, s)| |F(s, u(s), v(s), u'(s), v'(s))| ds &\leq M_2, \\ \int_a^b \max_{t \in [a, b]} \left| \frac{\partial G}{\partial t}(t, s) \right| |F(s, u(s), v(s), u'(s), v'(s))| ds &\leq M_3. \end{aligned}$$

Moreover,

$$\begin{aligned} \|T_1(u, v)(t)\| &= \max_{t \in [a, b]} \left| \frac{t-a}{b-a} \delta_1(b, u(b) + L_1(u, u(b), u'(b))) \right. \\ &\quad \left. + \int_a^b G(t, s) F(s, u(s), v(s), u'(s), v'(s)) ds \right| \\ &\leq \max_{t \in [a, b]} \left| \frac{t-a}{b-a} \right| |\delta_1(b, u(b) + L_1(u, u(b), u'(b)))| \\ &\quad + \int_a^b \max_{t \in [a, b]} |G(t, s)| |F(s, u(s), v(s), u'(s), v'(s))| ds \\ &\leq M_1 + M_2 < +\infty, \quad \forall (u, v) \in B, \end{aligned}$$

and

$$\begin{aligned}
\| (T_1(u, v))'(t) \| &= \max_{t \in [a, b]} \left| \frac{1}{b-a} \delta_1(b, u(b) + L_1(u, u(b), u'(b))) \right. \\
&\quad \left. + \int_a^b \frac{\partial G}{\partial t}(t, s) F(s, u(s), v(s), u'(s), v'(s)) ds \right| \\
&\leq \frac{M_1}{b-a} + \int_a^b \max_{t \in [a, b]} \left| \frac{\partial G}{\partial t}(t, s) \right| |F(s, u(s), v(s), u'(s), v'(s))| ds \\
&\leq \frac{M_1}{b-a} + M_3 < +\infty, \forall (u, v) \in B.
\end{aligned}$$

So,  $TB$  is uniformly bounded, for  $B$  a bounded set in  $(C^1[a, b])^2$ .

(iii)  $TB$  is equicontinuous on  $(C^1[a, b])^2$ .

Let  $t_1$  and  $t_2 \in [a, b]$ . Without loss of generality suppose  $t_1 \leq t_2$ . As  $G(t, s)$  is uniformly continuous and  $F$  is bounded, then

$$\begin{aligned}
&|T_1(u, v)(t_1) - T_1(u, v)(t_2)| \\
&= \left| \frac{(t_1-a)-(t_2-a)}{b-a} \delta_1(b, u(b) + L_1(u, u(b), u'(b))) \right. \\
&\quad \left. + \int_a^b [G(t_1, s) - G(t_2, s)] F(s, u(s), v(s), u'(s), v'(s)) ds \right| \\
&\leq \left| \frac{t_1-t_2}{b-a} \right| M_1 + \left| \int_a^b [G(t_1, s) - G(t_2, s)] \right| |F(s, u(s), v(s), u'(s), v'(s))| ds \\
&\rightarrow 0, \text{ as } t_1 \rightarrow t_2,
\end{aligned}$$

and

$$\begin{aligned}
&|(T_1(u, v)(t_1))' - (T_1(u, v)(t_2))'| \\
&= \left| \int_a^{t_1} \left[ \frac{\partial G}{\partial t}(t_1, s) - \frac{\partial G}{\partial t}(t_2, s) \right] F(s, u(s), v(s), u'(s), v'(s)) ds \right| \\
&\leq \int_a^{t_1} \left| \frac{\partial G}{\partial t}(t_1, s) - \frac{\partial G}{\partial t}(t_2, s) \right| |F(s, u(s), v(s), u'(s), v'(s))| ds \\
&\quad + \int_{t_1}^{t_2} \left| \frac{\partial G}{\partial t}(t_1, s) - \frac{\partial G}{\partial t}(t_2, s) \right| |F(s, u(s), v(s), u'(s), v'(s))| ds \\
&\quad + \int_{t_2}^b \left| \frac{\partial G}{\partial t}(t_1, s) - \frac{\partial G}{\partial t}(t_2, s) \right| |F(s, u(s), v(s), u'(s), v'(s))| ds.
\end{aligned}$$

As the function  $\frac{\partial G}{\partial t}(t, s)$  has only a jump discontinuity at  $t = s$ , therefore, as previously, the first and third integrals tend to 0, as  $t_1 \rightarrow t_2$ . For the second integral, as the functions  $\frac{\partial G}{\partial t}(t_1, s)$  and  $\frac{\partial G}{\partial t}(t_2, s)$  are uniformly continuous, for  $s \in [a, t_1 \cup t_1, b]$  and  $s \in [a, t_2 \cup t_2, b]$ , respectively, and

$$\left| \frac{\partial G}{\partial t}(t_1, s) - \frac{\partial G}{\partial t}(t_2, s) \right| |F(s, u(s), v(s), u'(s), v'(s))|$$

is bounded, then

$$\int_{t_1}^{t_2} \left| \frac{\partial G}{\partial t}(t_1, s) - \frac{\partial G}{\partial t}(t_2, s) \right| |F(s, u(s), v(s), u'(s), v'(s))| ds \rightarrow 0, \text{ as } t_1 \rightarrow t_2.$$

By the Arzèla-Ascoli Theorem  $T_1(u, v)$  is compact in  $(C^1[a, b])^2$ .

Following similar arguments with  $K_1, K_2, K_3 > 0$  such that

$$\begin{aligned} \delta_2 &\leq \max \{ \| \alpha_2 \|, \| \beta_2 \| \} := K_1, \\ \int_a^b \max_{t \in [a, b]} |G(t, s)| |H(s, u(s), v(s), u'(s), v'(s))| ds &\leq K_2, \\ \int_a^b \max_{t \in [a, b]} \left| \frac{\partial G}{\partial t}(t, s) \right| |H(s, u(s), v(s), u'(s), v'(s))| ds &\leq K_3, \end{aligned}$$

it can be shown that  $T_2(u, v)$  is compact in  $(C^1[a, b])^2$ , too.

(iv)  $TD \subset D$  for some  $D \subset (C^1[a, b])^2$  a closed and bounded set.

Suppose  $D \subset (C^1[a, b])^2$  defined by

$$D = \left\{ (u, v) \in (C^1[a, b])^2 : \|(u, v)\|_{E^2} \leq 2\rho \right\},$$

where  $\rho$  is such that

$$\rho := \max \left\{ M_1 + M_2, \frac{M_1}{b-a} + M_3, K_1 + K_2, \frac{K_1}{b-a} + K_3, N_1, N_2 \right\},$$

with  $N_1, N_2$  given by ((2.7)).

Arguing as in Claim 3 (ii), it can be shown that

$$\begin{aligned} \|T_1(u, v)\| &\leq M_1 + M_2 \leq \rho, \\ \|(T_1(u, v))'\| &\leq \frac{M_1}{b-a} + M_3 \leq \rho \end{aligned}$$

and, therefore,  $\|T_1(u, v)\|_{C^1} \leq \rho$ .

Analogously  $\|T_2(u, v)\|_{C^1} \leq \rho$  and, so,

$$\begin{aligned} \|T(u, v)\|_{E^2} &= \|(T_1(u, v), T_2(u, v))\|_{E^2} \\ &= \|T_1(u, v)\|_{C^1} + \|T_2(u, v)\|_{C^1} \leq 2\rho. \end{aligned}$$

By Theorem 1, the operator  $T$ , given by (3.8) has a fixed point  $(u_0, v_0)$ .

*Claim 4:* This fixed point  $(u_0, v_0)$  is also solution of the initial problem (1.1), (1.2), if every solution of (3.2) verifies

$$\alpha_1(t) \leq u_0(t) \leq \beta_1(t), \quad \alpha_2(t) \leq v_0(t) \leq \beta_2(t), \quad \forall t \in [a, b], \quad (3.9)$$

$$\alpha_1(b) \leq u_0(b) + L_1(u_0, u_0(b), u'_0(b)) \leq \beta_1(b), \quad (3.9)$$

$$\alpha_2(b) \leq v_0(b) + L_2(v_0, v_0(b), v'_0(b)) \leq \beta_2(b). \quad (3.10)$$

Let  $(u_0, v_0)$  be a fixed point of  $T$ , that is  $(u_0, v_0)$  is a fixed point of  $T_1$  and  $T_2$ .

By Claim 1,  $(u_0, v_0)$  is solution of problem (3.2).

In the following we will prove the estimations for  $u_0$ , as for  $v_0$  the procedure is analogous.

Suppose, by contradiction, that the first inequality is not true. So, there exists  $t \in [a, b]$  such that  $\alpha_1(t) > u_0(t)$  and it can be defined

$$\max_{t \in [a, b]} (\alpha_1(t) - u_0(t)) := \alpha_1(t_0) - u_0(t_0) > 0. \quad (3.11)$$

Remark that, by (3.2), Definition 1 and (3.3),  $t_0 \neq a$ , as  $\alpha_1(a) - u_0(a) \leq 0$ , and  $t_0 \neq b$ , because

$$\alpha_1(b) - u_0(b) = \alpha_1(b) - \delta_1(b, u_0(b) + L_1(u_0, u_0(b), u'_0(b))) \leq 0.$$

Then  $t_0 \in ]a, b[$ ,

$$\alpha'_1(t_0) - u'_0(t_0) = 0 \text{ and } \alpha''_1(t_0) - u''_0(t_0) \leq 0. \quad (3.12)$$

There are three possibilities for the value of  $v_0(t_0)$ :

- If  $v_0(t_0) > \beta_2(t_0)$ , then, by (3.2) and Definition 1, the following contradiction with (3.12) is obtained

$$\begin{aligned} u''_0(t_0) &= F(t_0, u_0(t_0), v_0(t_0), u'_0(t_0), v'_0(t_0)) \\ &= f(t_0, \alpha_1(t_0), \alpha_2(t_0), \alpha'_1(t_0), \alpha'_2(t_0)) - \frac{u_0(t_0) - \alpha_1(t_0)}{1 + |u_0(t_0) - \alpha_1(t_0)|} \\ &\quad + \frac{v_0(t_0) - \beta_2(t_0)}{1 + |v_0(t_0) - \beta_2(t_0)|} \\ &> f(t_0, \alpha_1(t_0), \alpha_2(t_0), \alpha'_1(t_0), \alpha'_2(t_0)) \geq \alpha''_1(t_0). \end{aligned}$$

- If  $\alpha_2(t_0) \leq v_0(t_0) \leq \beta_2(t_0)$ , then

$$\begin{aligned} u''_0(t_0) &= F(t_0, u_0(t_0), v_0(t_0), u'_0(t_0), v'_0(t_0)) \\ &= f(t_0, \alpha_1(t_0), \alpha_2(t_0), \alpha'_1(t_0), \alpha'_2(t_0)) - \frac{u_0(t_0) - \alpha_1(t_0)}{1 + |u_0(t_0) - \alpha_1(t_0)|} \\ &> f(t_0, \alpha_1(t_0), \alpha_2(t_0), \alpha'_1(t_0), \alpha'_2(t_0)) \geq \alpha''_1(t_0). \end{aligned}$$

- If  $\alpha_2(t_0) < v_0(t_0)$ , the contradiction is

$$\begin{aligned} u''_0(t_0) &= F(t_0, u_0(t_0), v_0(t_0), u'_0(t_0), v'_0(t_0)) \\ &= f(t_0, \alpha_1(t_0), \alpha_2(t_0), \alpha'_1(t_0), \alpha'_2(t_0)) - \frac{u_0(t_0) - \alpha_1(t_0)}{1 + |u_0(t_0) - \alpha_1(t_0)|} \\ &\quad + \frac{v_0(t_0) - \alpha_2(t_0)}{1 + |v_0(t_0) - \alpha_2(t_0)|} \\ &> f(t_0, \alpha_1(t_0), \alpha_2(t_0), \alpha'_1(t_0), \alpha'_2(t_0)) \geq \alpha''_1(t_0). \end{aligned}$$

Therefore  $\alpha_1(t) \leq u_0(t)$ ,  $\forall t \in [a, b]$ .

By a similar technique it can be shown that  $u_0(t) \leq \beta_1(t)$ ,  $\forall t \in [a, b]$ , and so,

$$\alpha_1(t) \leq u_0(t) \leq \beta_1(t), \forall t \in [a, b]. \quad (3.13)$$

Assume now that, to prove the first inequality of (3.9),

$$\alpha_1(b) > u_0(b) + L_1(u_0, u_0(b), u'_0(b)). \quad (3.14)$$

Then, by (3.2) and (3.3),

$$u_0(b) = \delta_1(b, u_0(b) + L_1(u_0, u_0(b), u'_0(b))) = \alpha_1(b),$$

and

$$u'_0(b) \leq \alpha'_1(b).$$

By (3.14), (A) and Definition 1, we have the contradiction

$$\begin{aligned} 0 &> L_1(u_0, u_0(b), u'_0(b)) + u_0(b) - \alpha_1(b) \\ &\geq L_1(u_0, u_0(b), u'_0(b)) \geq L_1(\alpha_1, \alpha_1(b), \alpha'_1(b)) \geq 0. \end{aligned}$$

Then  $\alpha_1(b) \leq u_0(b) + L_1(u_0, u_0(b), u'_0(b))$ .

To prove the second inequality of (3.9), assume that

$$u_0(b) + L_1(u_0, u_0(b), u'_0(b)) > \beta_1(b). \quad (3.15)$$

Then, by (3.2) and (3.3),

$$u_0(b) = \delta_1(b, u_0(b) + L_1(u_0, u_0(b), u'_0(b))) = \beta_1(b), \quad (3.16)$$

and

$$u'_0(b) \geq \beta'_1(b).$$

By (3.14), (3.16), (A) and Definition 1, we have the contradiction

So,  $u_0(b) + L_1(u_0, u_0(b), u'_0(b)) \leq \beta_1(b)$ , and, therefore, (3.9) holds.

To prove (3.10) the technique is analogous.

So, the fixed point  $(u_0, v_0)$  of  $T$ , solution of problem (3.2), is a solution of problem (1.1), (1.2), too.  $\square$

#### 4. Example

Consider the boundary value problem composed by the coupled system of the second order differential equations with full nonlinearities

$$\begin{cases} u''(t) = -u'(t)v(t) + \arctan(u(t)v'(t)) + t, \\ v''(t) = t^2 \left[ -e^{-|u'(t)|}v(t) + u(t)(v'(t) - 2) \right] \end{cases} \quad (4.1)$$

with  $t \in [0, 1]$ , and the functional boundary conditions

$$\begin{cases} u(0) = v(0) = 0 \\ u(1) = \max_{t \in [0,1]} u(t) - (u'(1))^2 \\ v(1) = 2 \int_0^1 v(s) ds - \frac{(v'(1))^3}{8}. \end{cases} \quad (4.2)$$

This problem is a particular case of system (1.1), (1.2) with

$$\begin{aligned} f(t, x, y, z, w) &= -z y + \arctan(x w) + t, \\ h(t, x, y, z, w) &= t^2 [-e^{-|z|} y + x(w - 2)], \end{aligned} \quad (4.3)$$

continuous functions,  $t \in [0, 1]$ ,  $a = 0$ ,  $b = 1$ , and

$$\begin{aligned} L_1(w, x, y) &= x - \max_{t \in [0,1]} w(t) + y^2, \\ L_2(w, x, y) &= x - 2 \int_0^1 w(s) ds + \frac{y^3}{8}. \end{aligned} \quad (4.4)$$

Remark that,  $L_1$  and  $L_2$  are continuous functions, verifying (A).

The functions given by

$$(\alpha_1(t), \alpha_2(t)) = (-1, -1) \text{ and } (\beta_1(t), \beta_2(t)) = (3+t, 2t+2)$$

are, respectively, lower and upper solutions of problem (4.1)–(4.2), satisfying (3.1), because, by Definition (1), we have

$$\begin{aligned} \alpha_1''(t) &= 0 \leqslant f(t, -1, -1, 0, 0) = t, \quad t \in [0, 1] \\ \alpha_2''(t) &= 0 \leqslant h(t, -1, -1, 0, 0) = 3t^2, \quad t \in [0, 1] \\ L_1(\alpha_1, \alpha_1(1), \alpha_1'(1)) &= 0 \geqslant 0 \\ L_2(\alpha_2, \alpha_2(b), \alpha_2'(b)) &= \frac{7}{8} \geqslant 0, \end{aligned}$$

and

$$\begin{aligned} \beta_1(t)''(t) &= 0 \geqslant f(t, 3+t, 2t+2, 1, 2) \\ &= -(2t+2) + \arctan(6t+2) + t, \quad t \in [0, 1] \\ \beta_2''(t) &= 0 \geqslant h(t, 3+t, 2t+2, 1, 2) = -\frac{2t^3 + 2t^2}{e}, \quad t \in [0, 1] \\ L_1(\beta_1, \beta_1(1), \beta_1'(1)) &= 0 \leqslant 0 \\ L_2(\beta_2, \beta_2(b), \beta_2'(b)) &= -1 \leqslant 0. \end{aligned}$$

Furthermore, the functions  $f$  and  $h$ , given by (4.3), satisfy a Nagumo condition relative to the intervals  $[-1, 2+t]$  and  $[-1, 2t+1]$ , for  $t \in [0, 1]$ , with  $r_1 = r_2 = 4$ ,

$$|f(t, x, y, z, w)| \leqslant 3|z| + \frac{\pi}{2} + 1 := \varphi(|z|),$$

$$|h(t, x, y, z, w)| \leq 3(1 + |w - 2|) := \psi(|w|),$$

and

$$\int_4^{N_1} \frac{ds}{\varphi(|s|)} = \int_4^{N_1} \frac{1}{3s + \frac{\pi}{2} + 1} ds > 1,$$

for  $N_1 \geq 100$  and

$$\int_4^{N_2} \frac{ds}{\psi(|s|)} = \int_4^{N_2} \frac{1}{3(1 + |s - 2|)} ds > 1,$$

for  $N_2 \geq 80$ .

Then, by Theorem (2), there is at least a pair  $(u(t), v(t)) \in (C^2[0, 1], \mathbb{R})^2$  solution of (4.1), (4.2) and, moreover,

$$\begin{aligned} -1 \leq u(t) \leq 2 + t, \quad -1 \leq v(t) \leq 2t + 1, \quad \forall t \in [0, 1], \\ \|u'\| \leq N_1 \text{ and } \|v'\| \leq N_2. \end{aligned}$$

## 5. Coupled mass-spring system

Consider the mass-spring system composed by two springs with constants of proportionality  $k_1$  and  $k_2$ , and two weights of mass  $m_1$  and  $m_2$ . The mass  $m_1$  is suspended vertically from a fixed support by a spring with constant  $k_1$  and the mass  $m_2$  is attached to the first weight by a spring with constant  $k_2$ . The system described is illustrated in the Figure 1.




Figure 1: The coupled springs.

Let us call  $u(t)$  and  $v(t)$  the displacements of the weights of mass  $m_1$  and  $m_2$ , respectively, in relation to their respective equilibrium positions. Thus, at time  $t$ , the

position of the displacement of the mass  $m_1$  is  $u(t)$  and the displacement of mass  $m_2$  is  $v(t)$ .

For simplicity we consider  $t \in [0, 1]$ , and, therefore,  $u(0)$  and  $u(1)$  are the initial and final displacement of mass  $m_1$ , and  $v(0)$  and  $v(1)$  are the similar displacements of mass  $m_2$ .

As it can be seen in [8], the above system is modelled by the second order nonlinear system of differential equations forced and with friction

$$\begin{cases} m_1 u''(t) = -\delta_1 u'(t) - \kappa_1 u(t) + \mu_1(u(t))^3 - \kappa_2(u(t) - v(t)) \\ \quad + \mu_2(u(t) - v(t))^3 + F_1 \cos(\omega_1 t), \\ m_2 v''(t) = -\delta_2 v'(t) - \kappa_2(v(t) - u(t)) + \mu_2(v(t) - u(t))^3 \\ \quad + F_2 \cos(\omega_2 t) \end{cases} \quad (5.1)$$

where  $t \in [0, 1]$ ,

- $\delta_1, \delta_2$  are the damping coefficients;
- $\mu_1, \mu_2$  are the coefficients of the nonlinear terms of each system equation;
- $\kappa_2(u(t) - v(t)) + \mu_2(u(t) - v(t))^3$  and  $\kappa_2(v(t) - u(t)) + \mu_2(v(t) - u(t))^3$  are the nonlinear restoring forces;
- $F_1, F_2$  are the forcing amplitudes of the sinusoidal forces  $F_1 \cos(\omega_1 t)$  and  $F_2 \cos(\omega_2 t)$ , where  $\omega_1, \omega_2$  are the forcing frequencies.

In this work we add to the system the functional boundary conditions

$$\begin{cases} u(0) = v(0) = 0 \\ u(1) = \max_{t \in [0, 1]} u(t) + 2u'(1) \\ v(1) = \max_{t \in [0, 1]} v(t) + 2(v'(1))^3. \end{cases} \quad (5.2)$$

The functional conditions (5.2) can have a physical meaning such as, for example, the first one can be seen as the displacement of mass 1 at the final moment given by the sum of the maximum displacement in this period of time, with the double of the velocity of the displacement at the end point.

Clearly, the above model (5.1), (5.2) is a particular case of system (1.1), (1.2) with

$$f(t, x, y, z, w) = \frac{1}{m_1} \left[ -\delta_1 z - \kappa_1 x + \mu_1 x^3 - \kappa_2(x - y) + \mu_2(x - y)^3 \right. \\ \left. + F_1 \cos(\omega_1 t) \right], \quad (5.3)$$

$$h(t, x, y, z, w) = \frac{1}{m_2} \left[ -\delta_2 w - \kappa_2(y - x) + \mu_2(y - x)^3 + F_2 \cos(\omega_2 t) \right].$$

These functions are continuous in  $[0, 1] \times \mathbb{R}^4$ , and

$$\begin{aligned} L_1(w, x, y) &= x - \max_{t \in [0, 1]} w(t) - 2y, \\ L_2(w, x, y) &= x - \max_{t \in [0, 1]} w(t) - 2y^3 \end{aligned} \quad (5.4)$$

verify (A).

The functions given by

$$(\alpha_1(t), \alpha_2(t)) = (-t, -t) \text{ and } (\beta_1(t), \beta_2(t)) = (t, t)$$

are, respectively, lower and upper solutions of problem (5.1), (5.2), satisfying (3.1), for every positive  $m_1, m_2$ , non negative  $\delta_1, \delta_2, F_1, F_2, \omega_1, \omega_2$ , and any real  $\kappa_1, \kappa_2, \mu_1, \mu_2$ , such that

$$\begin{aligned} F_1 &\leq \delta_1, \\ F_2 &\leq \delta_2, \\ \mu_1 &\leq 0. \end{aligned} \tag{5.5}$$

Indeed, Definition 1 holds because,

$$\begin{aligned} \alpha_1''(t) = 0 &\leq \frac{1}{m_1} (\delta_1 - F_1) \\ &\leq \frac{1}{m_1} [\delta_1 - \mu_1 t^3 + F_1 \cos(\omega_1 t)], \forall t \in [0, 1] \\ &\leq \frac{1}{m_1} [\delta_1 + \kappa_1 t - \mu_1 t^3 + F_1 \cos(\omega_1 t)], \forall t \in [0, 1] \\ \alpha_2''(t) = 0 &\leq \frac{1}{m_2} (\delta_2 - F_2) \leq \frac{1}{m_2} [\delta_2 + F_2 \cos(\omega_2 t)], \forall t \in [0, 1] \\ L_1(\alpha_1, \alpha_1(1), \alpha_1'(1)) &= \alpha_1(1) - \max_{t \in [0, 1]} \alpha_1(t) - 2\alpha_1'(1) = 1 \geq 0 \\ L_2(\alpha_2, \alpha_2(b), \alpha_2'(b)) &= \alpha_2(1) - \max_{t \in [0, 1]} \alpha_2(t) - 2(\alpha_2'(1))^3 = 1 \geq 0, \end{aligned}$$

and

$$\begin{aligned} \beta_1''(t) = 0 &\geq \frac{1}{m_1} (-\delta_1 + F_1) \\ &\geq \frac{1}{m_1} [-\delta_1 - \kappa_1 t + \mu_1 t^3 + F_1 \cos(\omega_1 t)], \forall t \in [0, 1] \\ \beta_2''(t) = 0 &\geq \frac{1}{m_2} (-\delta_2 + F_2) \geq \frac{1}{m_2} [-\delta_2 + F_2 \cos(\omega_2 t)], \forall t \in [0, 1] \\ L_1(\beta_1, \beta_1(1), \beta_1'(1)) &= \beta_1(1) - \max_{t \in [0, 1]} \beta_1(t) - 2\beta_1'(1) = -2 \leq 0 \\ L_2(\beta_2, \beta_2(b), \beta_2'(b)) &= \beta_2(1) - \max_{t \in [0, 1]} \beta_2(t) - 2(\beta_2'(1))^3 = -2 \leq 0. \end{aligned}$$

Furthermore,  $f$  and  $h$ , given by (5.3), verify a Nagumo condition relative to the interval  $[-t, t]$ , for  $t \in [0, 1]$ , with  $r_1 = r_2 = 1$ , for

$$-t \leq x \leq t, -t \leq y \leq t, \forall t \in [0, 1],$$

$$\begin{aligned} |f(t, x, y, z, w)| &\leq \frac{1}{m_1} (\delta_1 |z| + \kappa_1 + |\mu_1| + 2\kappa_2 + 8|\mu_2| + F_1) := \varphi(|z|), \\ |h(t, x, y, z, w)| &\leq \frac{1}{m_2} (\delta_2 |w| + 2\kappa_2 + 8|\mu_2| + F_2) := \psi(|w|), \end{aligned}$$

and for  $N_1$  and  $N_2$  positive, and large enough such that

$$\int_1^{N_1} \frac{ds}{\varphi(s)} = \int_1^{N_1} \left( \frac{m_1}{\delta_1 s + \kappa_1 + |\mu_1| + 2\kappa_2 + 8|\mu_2| + F_1} \right) ds > 1,$$

and

$$\int_1^{N_2} \frac{ds}{\psi(s)} = \int_1^{N_2} \left( \frac{m_2}{\delta_2 s + 2\kappa_2 + 8|\mu_2| + F_2} \right) ds > 1.$$

So, by Theorem 2, there is a solution  $(u(t), v(t))$  of the mass-spring system (5.1), (5.2), for the values of coefficients verifying (5.5), such that

$$\begin{aligned} -t &\leq u(t) \leq t, \\ -t &\leq v(t) \leq t, \forall t \in [0, 1], \end{aligned}$$

that is, both with values in the strip and

$$\|u'\| \leq N_1 \text{ and } \|v'\| \leq N_2. \quad \square$$




Figure 2:  $u(t), v(t)$  in the strip

## REFERENCES

- [1] Y. AN, *Nonlinear perturbations of a coupled system of steady state suspension bridge equations*, Nonlinear Analysis **51** (2002), 1285–1292.
- [2] H. AMANN, *Parabolic evolution and nonlinear boundary conditions*, J. Differential Equations **72** (1988), 201–269.
- [3] R. ARIS, *Introduction to the analysis of chemical reactors*, Prentice-Hall, Englewood Cliffs, NJ, (1965).
- [4] D. G. ARONSON, *A comparison method for stability analysis of nonlinear parabolic problems*, SIAM Rev. **20**, (1978).
- [5] N. A. ASIF, P. W. ELOE, R. A. KHAN, *Positive solutions for a system of singular second order nonlocal boundary value problems*, J. Korean Math. Soc. **47**, **5** (2010), 985–1000.
- [6] A. CABADA, F. MINHÓS, *Fully nonlinear fourth-order equations with functional boundary conditions*, J. Math. Anal. Appl. **340** (2008), 239–251.
- [7] Y. CUI, J. SUN, *On existence of positive solutions of coupled integral boundary value problems for a nonlinear singular superlinear differential system*, Electron. J. Qual. Theory Differ. Equ. **41** (2012), 1–13.
- [8] T. H. FAY, S. D. GRAHAM, *Coupled spring equations*, Int. J. Math. Educ. Sci. Technol., vol. **34**, no. 1 (2003), 65–79.
- [9] J. GRAEF, L. KONG, F. MINHÓS, J. FIALHO, *On the lower and upper solution method for higher order functional boundary value problems*, Appl. Anal. Discrete Math., vol. **5**, no. 1 (2011), 133–146.
- [10] M. R. GROSSINHO, F. MINHÓS, A. I. SANTOS, *A note on a class of problems for a higher order fully nonlinear equation under one sided Nagumo type condition*, Nonlinear Anal. **70** (2009), 4027–4038.
- [11] J. HENDERSON, R. LUCA, *Boundary Value Problems for Systems of Differential, Difference and Fractional Equations, Positive Solutions*, Elsevier, 2015.
- [12] J. HENDERSON, R. LUCA, *Positive solutions for systems of nonlinear second-order multipoint boundary value problems*, Math. Meth. Appl. Sci. **37** (2014), 2502–2516.
- [13] G. INFANTE, F. M. MINHÓS, P. PIETRAMALA, *Non-negative solutions of systems of ODES with coupled boundary conditions*, Communications in Nonlinear Science and Numerical Simulations **17** (2012), 4952–4960.
- [14] T. JANKOWSKI, *Nonnegative solutions to nonlocal boundary value problems for systems of second-order differential equations dependent on the first-order derivatives*, Nonlinear Analysis **87** (2013), 83–101.
- [15] P. KANG, Z. WEI, *Existence of positive solutions for systems of bending elastic beam equations*, Electron. J. Differential Equations **19** (2012).
- [16] A. LEUNG, *A semilinear reaction-diffusion prey-predator system with nonlinear coupled boundary conditions: Equilibrium and Stability*, Indiana Univ. Math. J. **31** (1982), 223–241.
- [17] Y. LI, Y. GUO, G. LI, *Existence of positive solutions for systems of nonlinear third-order differential equations*, Commun Nonlinear Sci Numer Simulat **14** (2009), 3792–3797.
- [18] H. LÜ, H. YU, Y. LIU, *Positive solutions for singular boundary value problems of a coupled system of differential equations*, J. Math. Anal. Appl. **302** (2005), 14–29.
- [19] F. A. MEHMETI, S. NICAISE, *Nonlinear interaction problems*, Nonlinear Anal. **20**, no. 1 (1993), 27–61.
- [20] F. MINHÓS, R. DE SOUSA, *On the solvability of third-order three points systems of differential equations with dependence on the first derivatives*, Bulletin of the Brazilian Mathematical Society, New Series, vol. **48**, issue 3 (2017), 485–503.
- [21] F. MINHÓS, I. COXE, *System of coupled clamped beam equations: existence and localization results*, Nonlinear Analysis: Real World Applications **35** (2017), 45–60.
- [22] F. MINHÓS, J. FIALHO, *On the solvability of some fourth-order equations with functional boundary conditions*, Discrete Contin. Dyn. Syst., 2009, suppl., 564–573.
- [23] I. TALIB, N. A. ASIF, C. TUNC, *Coupled lower and upper solution approach for the existence of solutions of nonlinear coupled system with nonlinear coupled boundary conditions*, Proyecciones Journal of Mathematics, vol. **35**, no. 1 (2016), 99–117.
- [24] J. YANG, Z. WEI, *On existence of positive solutions of Sturm-Liouville boundary value problems for a nonlinear singular differential system*, J. Appl. Math. Comput. **217** (2011), 6097–6104.

- [25] F. YORUK, N. AYKUT HAMAL, *Second-order boundary value problems with integral boundary conditions on the real line*, Electronic Journal of Differential Equations, vol. **2014** (2014), no. 19, pp. 1–13.
- [26] A. WANG, J. SUN, A. ZETTL, *The classification of self-adjoint boundary conditions: separated, coupled*, J. Funct. Anal. **255** (2008), 1554–1573.
- [27] E. ZEIDLER, *Nonlinear Functional Analysis and Its Applications: Fixed-Point Theorems*, Springer, New York (1986).
- [28] F. ZHU, L. LIU, Y. WU, *Positive solutions for systems of a nonlinear fourth-order singular semipositive boundary value problems*, Applied Mathematics and Computation, **216** (2010), 448–457.

(Received March 31, 2017)

*Feliz Minhós*

*Departamento de Matemática, Escola de Ciências e Tecnologia  
Centro de Investigação em Matemática e Aplicações (CIMA)  
Instituto de Investigação e Formação Avançada  
Universidade de Évora  
Rua Romão Ramalho, 59, 7000-671 Évora, Portugal  
e-mail: fminhos@uevora.pt*

*Robert de Sousa  
Faculdade de Ciências e Tecnologia  
Núcleo de Matemática e Aplicações (NUMAT)  
Universidade de Cabo Verde. Campus de Palmarejo  
279 Praia, Cabo Verde  
e-mail: robert.sousa@docente.unicv.edu.cv*