



Contents lists available at ScienceDirect

Comptes Rendus Geoscience

www.sciencedirect.com



Petrology, Geochemistry (Geochronology)

U-Pb laser ablation ICP-MS zircon dating across the Ediacaran–Cambrian transition of the Montagne Noire, southern France



Maxime Padel ^{a,*}, J. Javier Álvaro ^b, Sébastien Clausen ^a, François Guillot ^c, Marc Poujol ^d, Martim Chichorro ^e, Éric Monceret ^f, M. Francisco Pereira ^g, Daniel Vizcaïno ^h

^a UMR 8198 EEP CNRS, université de Lille-1, bâtiment SN5, avenue Paul-Langevin, 59655 Villeneuve-d'Ascq cedex, France

^b Instituto de Geociencias (CSIC-UCM), Novais 12, 28040 Madrid, Spain

^c UMR 8187 LOG CNRS, université de Lille–université du Littoral Côte d'Opale, SN5 Sciences de la Terre, 59655 Villeneuve-d'Ascq cedex, France

^d Géosciences Rennes, UMR 6118, université de Rennes-1, campus de Beaulieu, 35042 Rennes, France

^e GEOBITEC/Departamento de Ciências da Terra, Universidade Nova de Lisboa, Portugal

^f 18, rue des Pins, 11570 Cazilhac, France

^g IDL/Departamento de Geociências, ECT, Universidade de Évora, Portugal

^h 7 c/o Jean-Baptiste Chardin, Maquens, 11090 Carcassonne, France

ARTICLE INFO

Article history:

Received 28 July 2016

Accepted after revision 25 November 2016

Available online 1 August 2017

Handled by Marc Chaussidon

Keywords:

Ediacaran

Cambrian

U–Pb dating

Montagne Noire

ABSTRACT

U–Pb laser ablation inductively coupled plasma mass spectrometry was used for dating zircon grains extracted from four sedimentary and volcano sedimentary rocks of the Montagne Noire encompassing the presumed Ediacaran–Cambrian boundary interval. Magmatic zircon from two samples from the basal and middle parts of the Riverous Formation (a rhyolitic tuff) were deposited at 542.5 ± 1 Ma and 537.1 ± 2.5 Ma, bracketing the 541 Ma age presently admitted as being at the Ediacaran–Cambrian boundary. In addition, a piece of sandstone from the underlying Riverous Formation containing mostly euhedral zircon grains, suggesting proximal magmatic sources, yields Neoproterozoic dates ranging from 574 Ma to 1 Ga, and subsidiary older dates from 1.25 to 2.75 Ga. Another piece of sandstone from the overlying Marcory Formation yielded mostly rounded zircon grains probably issued from more remote areas, with a large spectrum dominated by Neoproterozoic dates as well as older ages up to 3.2 Ga. A comparison of both kinds of sandstone suggests a significant change in provenance, changing from a restricted source area during the Ediacaran to a much larger source domain during the Cambrian Epoch 2 that recorded contributions from different cratons of Gondwana.

© 2017 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail addresses: maxime.padel@etudiant.univ-lille1.fr (M. Padel), jj.alvaro@csic.es (J.J. Álvaro), sebastien.clausen@univ-lille1.fr (S. Clausen), francois.guillot@univ-lille1.fr (F. Guillot), marc.poujol@univ-rennes1.fr (M. Poujol), ma.chichorro@fct.unl.pt (M. Chichorro), eric.monceret@orange.fr (É. Monceret), mpereira@uevora.pt (M.F. Pereira), daniel.vizcaino@wanadoo.fr (D. Vizcaïno).