

- Research Article
- [Published: 03 October 2020](#)

Monitoring system and in situ conservation of endemic and threatened *Beta patula* Aiton populations in Madeira Region

- [H. Nóbrega,](#)
- [G. Freitas,](#)
- [M. A. Zavattieri,](#)
- [C. Ragonezi,](#)
- [L. Frese](#) &
- [M. A. A. Pinheiro de Carvalho](#)

[Genetic Resources and Crop Evolution](#) (2020)[Cite this article](#)


- **49** Accesses
- [Metricsdetails](#)

Abstract

Madeira Archipelago is a hotspot for crop wild relatives (CWR) of crop cultures. Some of these CWR are present in very specific environments, such as, in Ponta de São Lourenço or Desertas Islands. One such species is *Beta patula* Aiton, a Critically Endangered species which belongs to the Gene Pool 1b of cultivated beets. A continuous effort has been developed for its ex situ conservation through the storage of accessions (ISOP2512 and ISOP1911) in the ISOPlexis GeneBank at the University of Madeira. Simultaneously, a series of studies have been carried out to understand the species' ecogeographic and ecological requirements, to validate populations' boundaries and sizes, and to establish population dynamics. This study includes a complete floristic survey at the *B. patula* locations of, Desembarcadouro islet (DI) with 12 (DI1–DI12) sampling sites, and Chão islet (CI) with 3 (CI1–CI3). Several Biodiversity indices were calculated for these locations. Plot DI3 exhibited the highest values for Corrected Evenness ($E' = 0.77 \pm 0.07$), Shannon–Weaver Diversity Index ($H' = 2.48 \pm 0.12$), and Hill's Index ($N_2 = 4.47 \pm 0.72$), with a total sum of 306 individuals of *B. patula*. The demographic status of *B. patula* populations in DI and CI was determined yearly between 2014 and 2018. The results show an average population size of 16,906 and

2917 plants, respectively. These data will be used for the establishment of a protocol to monitor and manage a genetic reserve for *B. patula* and other CWR. By doing so, our work will contribute to the implementation of the European genetic reserve network.

Graphic abstract



This is a preview of subscription content, [log in](#) to check access.

References

1. Adams J (2009) The holy grail of ecology: latitudinal gradients. Species Richness Patterns the Divers Life. https://doi.org/10.1007/978-3-540-74278-4_2

[Article Google Scholar](#)

2. Aguiar C, Capelo J, Fontinha S, Espirito-Santo D, Jardim R, Lousã M, Rivas-Martinez S, Mesquita S, Sequeira M, de Sousa J (2004) A paisagem vegetal da Ilha da Madeira. Quercetea 6:3–200

[Google Scholar](#)

3. Alves SF, Reis F, Henriques D, Freitas G, Gouveia CSS, Fernandes F, Carvalho JA, Pinheiro de Carvalho MAA (2019) *Solanum trisetum* Dunal: prospection, conservation and its ecogeographic baseline. *Asian J Plant Sci Res* 9(2):16–29

[Google Scholar](#)

4. Biancardi E, Panella L, Lewellen RT (2012) Source of useful traits. In: *Beta maritima*, Springer, New York, NY, pp 173–223
5. Borges PAV, Abreu C, Aguiar AMF, Carvalho P, Jardim R, Melo I, Oliveira P, Sérgio C, Serrano ARM, Vieira P (2008) A list of the terrestrial fungi, flora and fauna of Madeira and Selvagens archipelagos. Direcção Regional do Ambiente da Madeira, Universidade dos Açores, Funchal and Angra do Heroísmo
6. Brehm JM, Maxted N, Ford-Lloyd BV, Martins-Louçao MA (2008) National inventories of crop wild relatives and wild harvested plants: case-study for Portugal. *Genet Resour Crop Evol* 55(6):779–796

[Article Google Scholar](#)

7. Brozynska M, Furtado A, Henry RJ (2015) Genomics of crop wild relatives: expanding the gene pool for crop improvement. *Plant Biotechnol J* 14:1070–1085

[PubMed Article](#) [CAS](#) [PubMed Central](#) [Google Scholar](#)

8. Burnham KP, Overton WS (1979) Robust estimation of population size when capture probabilities vary among animals. *Ecol* 60(5):927–936

[Article Google Scholar](#)

9. Capistrano-Gossmann GG, Ries D, Holtgrawe D et al (2016) Crop wild relative populations of *Beta vulgaris* allow direct mapping of agronomically important genes. *Nat Commun* 8(15708):1–8

[Google Scholar](#)

10. Carvalho M, Frese L, Duarte MC, Magos Brehm J, Tavares M, Santos Guerra A, Draper D (2011) *Beta patula*. In: The IUCN red list of threatened species 2011:

e.T162088A5532483. <https://dx.doi.org/10.2305/IUCN.UK.2011-1.RLTS.T162088A5532483.en>. Accessed 31 Aug 2020

11. Costa JC, Neto CS, Aguiar C et al (2012) Vascular plant communities in Portugal (Continental, the Azores and Madeira). *Glob Geobot* 2:1–180

[Google Scholar](#)

12. De Hond L, Iriondo JM, Kell S (2004) Report of workshop 4: population management methodologies. PGR Forum, European Crop Wild Relative Diversity Assessment and Conservation Forum, Mahón, Menorca, Spain
13. De Vilmorin JL (1923) L'héritité chez la betterave cultivée. Thèse de doctorat, faculté des science, Paris, Gauthier-Villars et Cie, editeurs. Libraires du bureau des longitudes, de l'école polytechnique
14. Dempewolf H, Baute G, Anderson J, Kilian B, Smith C, Guarino L (2017) Past and future use of wild relatives in crop breeding. *Crop Sci* 57(3):1070–1082

[Article Google Scholar](#)

15. Dewar AM, Haylock LA, Ecclestone PMJ (1996) Strategies for controlling aphids and virus yellows in sugar beet. In: *Proceedings of the Brighton crop protection conference, pests and diseases, Brighton, England*, pp 190–195
16. Directive H (1992) Council directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. *Off J Eur Union* 206:7–50

[Google Scholar](#)

17. Drozd P (2010) Community ecology parameter calculator. ComEcoPaC <http://prf.osu.cz/kbe/dokumenty/sw/ComEcoPaC/ComEcoPaC.xls>
18. Dwivedi SL, Upadhyaya HD, Stalker HT, Blair MW, Bertoli DJ, Nielen S, Ortiz R (2008) Enhancing crop gene pools with beneficial traits using wild relatives. *Plant Breed Rev* 30:179

[CAS Article Google Scholar](#)

19. Eizenga GC, Agrama HA, Lee FN, Jia Y (2009) Exploring genetic diversity and potential novel disease resistance genes in a collection of rice (*Oryza* spp.) wild relatives. *Genet Resour Crop Evol* 56:65–76
-

[CAS Article](#) [Google Scholar](#)

20. FAO (2009) International treaty on plant genetic resources for food and agriculture. FAO, Rome
-

[Google Scholar](#)

21. FAO (2014) Genebank standards for plant genetic resources for food and agriculture. Commission on Genetic Resources for Food and Agriculture, Rome, Italy
22. FAO/UNESCO (1988) Soil map of the world: revised legend. FAO/UNESCO, Rome
-

[Google Scholar](#)

23. Fontinha S, Carvalho JA (1995) Evaluation of the vascular flora of Madeira's extreme East. *Bol Museu Municipal Funchal* 4:263–275
-

[Google Scholar](#)

24. Foolad MR, Panthee DR (2012) Marker-assisted selection in tomato breeding. *Crit Rev Plant Sci* 31:93–123
-

[Article](#) [Google Scholar](#)

25. Franklin IR (1980) Evolutionary change in small populations. In: Soule ME, Wilcox BA (eds) *Conservation biology: an evolutionary-ecological perspective*, Sunderland, Massachusetts, pp 135–149
26. Frese L (2003) Sugar beets and related wild species—from collecting to utilisation. *Schriften zu Genetischen Ressourcen* 22:170–181
-

[Google Scholar](#)

27. Frese L (2010) Conservation and access to sugar beet germplasm. *Sugar Tech* 12(3–4):207–219

[Article Google Scholar](#)

28. Frese L, Pinheiro de Carvalho MAA (2010) Case crop study Beta (including *Patellifolia*) part B: EU-level case study. EU Guidelines for the creation for *Beta* genetic resources conservation. Work paper. ECPGR, Roma
29. Frese L, Nachtigall M, Enders M, Pinheiro de Carvalho MAA (2012) *Beta patula* Aiton: genetic diversity analysis. In: Maxted N, Lothar F, Iriondo J, Dulloo E, Ford-Lloyd BV, Pinheiro de Carvalh (eds) *Agrobiodiversity conservation: securing the diversity of crop wild relatives and landraces*. CABI, Wallingford, pp 45–51
30. Frese L, Nachtigall M, Iriondo JM, Rubio Teso ML, Duarte MC, Pinheiro de Carvalho MAA (2018) Genetic diversity and differentiation in *Patellifolia* (Amaranthaceae) in the Macaronesian archipelagos and the Iberian Peninsula and implications for genetic conservation programmes. *Genet Resour Crop Evol* 66(1):225–241
-

[Article CAS Google Scholar](#)

31. Frese L, Nachtigall M, Iriondo JM, Teso MLR, Duarte MC, de Carvalho MAA (2019) Genetic diversity and differentiation in *Patellifolia* (Amaranthaceae) in the Macaronesian archipelagos and the Iberian Peninsula and implications for genetic conservation programmes. *Genet Resour Crop Evol* 66(1):225–241
-

[CAS Article Google Scholar](#)

32. Fuller DQ, Denham T, Arroyo-Kalin M et al (2014) Convergent evolution and parallelism in plant domestication revealed by an expanding archaeological record. *Proc Natl Acad Sci* 111(17):6147–6152
-

[CAS PubMed Article PubMed Central Google Scholar](#)

33. Gosselin F (2006) An assessment of the dependence of evenness indices on species richness. *J Theor Biol* 24(3):591–597
-

[Article](#) [Google Scholar](#)

34. Gregorius HR, Gillet EM (2015) Classifying measures of biological variation. PLoS ONE 10(3):e0115312

[PubMed](#) [PubMed Central](#) [Article](#) [CAS](#) [Google Scholar](#)

35. Hajjar R, Hodgkin T (2007) The use of wild relatives in crop improvement: a survey of developments over the last 20 years. Euphytica 156:1–13

[Article](#) [Google Scholar](#)

36. Harlan JR (1992) Origins and processes of domestication. In: Grass evolution and domestication, pp 159–175
37. Harlan J, de Wet J (1971) Towards a rational classification of cultivated plants. Taxon 20:509–517

[Article](#) [Google Scholar](#)

38. Hauer M, Hansen AL, Manderyck B, Olsson Å, Raaijmakers E, Hanse B, Märländer B (2017) Neonicotinoids in sugar beet cultivation in Central and Northern Europe: efficacy and environmental impact of neonicotinoid seed treatments and alternative measures. Crop Prot 93:132–142

[CAS](#) [Article](#) [Google Scholar](#)

39. Heijbroek W, Roelands AJ, De Jong JH (1983) Transfer of resistance to beet cyst nematode from *Beta patellaris* to sugar beet. Euphytica 32(2):287–298

[Article](#) [Google Scholar](#)

40. Heywood V, Kell S, Maxted N (2005) Global strategy for crop wild relative conservation and use. In: First international conference on crop wild relative conservation and use, PGR Forum, Agrigento, Italy
41. Hubalek Z (2000) Measures of species diversity in ecology: an evaluation. Folia Zool Praha 49(4):241–260

[Google Scholar](#)

42. Hufford MB, Lubinsky P, Pyhäjärvi T, Devengenzo MT, Ellstrand NC, Ross-Ibarra J (2013) The genomic signature of crop-wild introgression in maize. *PLoS Genet* 9(5):e1003477
-

[CAS](#) [PubMed](#) [PubMed Central](#) [Article](#) [Google Scholar](#)

43. Instituto Português do Mar e da atmosfera. <http://www.ipma.pt/pt/>. Accessed 01 Apr 2019
44. Iriondo JM, Ford-Lloyd B, DeHond L, Kell SP, Lefevre F, Korpelainen H, Lane A (2008) Plant population monitoring methodologies for the in situ genetic conservation of CWR. In: Iriondo JM, Maxted N, Dulloo ME (eds) *Conserving plant genetic diversity in protected areas: population management of crop wild relatives*. CABI, Oxfordshire, pp 89–116
-

[Google Scholar](#)

45. Iriondo JM, Maxted N, Kell SP, Ford-Lloyd B, Lara-Romero C, Labokas J, Pinheiro de Carvalho, MAA (2012) Identifying quality standards for genetic reserve conservation of CWR. In: Nigel (ed) *Agrobiodiversity conservation: securing the diversity of crop wild relatives and landraces*, pp 72–77
46. Kantar MB, Nashoba AR, Anderson JE, Blackman BK, Rieseberg LH (2017) The genetics and genomics of plant domestication. *Bioscience* 67(11):971–982
-

[Article](#) [Google Scholar](#)

47. Kell SP, Maxted N, Frese L, Iriondo JM (2012) In situ conservation of crop wild relatives: a strategy for identifying priority genetic reserve sites. In: Maxted N, Dulloo ME, Ford-Lloyd BV, Frese L, Iriondo JM, PinheirodeCarvalho MAA (eds) *Agrobiodiversity conservation: securing the diversity of crop wild relatives and landraces*. CAB International, Wallingford, pp 7–19
-

[Google Scholar](#)

48. Lange W, Brandenburg WA, De Bock TSM (1999) Taxonomy and culonomy of beet (*Beta vulgaris* L.). *Bot J Linn Soc* 130:81–96
-

[Article](#) [Google Scholar](#)

49. Letschert JPW (1993) *Beta* section *Beta*: biogeographical patterns of variation, and taxonomy. Dissertation, Wageningen Agricultural University
50. Luterbacher MC, Asher MJC, DeAmbrogio E, Biancardi E, Stevenato P, Frese L (2004) Sources of resistance to diseases of sugar beet in related *Beta* germplasm: I. Foliar diseases. *Euphytica* 139(2):105–121

[Article Google Scholar](#)

51. Marchante H, Morais M, Freitas H, Elizabete Marchante E (2008) Guia prático para a identificação de plantas invasoras em Portugal. Imprensa da Universidade de Coimbra, Coimbra, Portugal
52. Maxted N, Ford-loyd BV, Hawkes JG (1997) Complementary conservation strategies. In: Maxted N, Ford-loyd BV, Hawkes JG (eds) *Plant genetic conservation: the in situ approach*. Chapman & Hall, London, pp 20–55

[Google Scholar](#)

53. Maxted M, Ford-Lloyd BV, Jury S, Kell Sh, Scholten M (2006) Towards a definition of a crop wild relative. *Biodivers Conserv* 15:2673–2685

[Article Google Scholar](#)

54. Maxted N, Kell S, Toledo Á et al (2010) A global approach to crop wild relative conservation: securing the gene pool for food and agriculture. *Kew Bull* 65(4):561–576

[Article Google Scholar](#)

55. Maxted N, Avagyan A, Frese L, Iriondo JM, Magos-Brehm J, Singer A, Kell SP (2015) ECPGR Concept for in situ conservation of crop wild relatives in Europe. Wild species conservation in genetic reserves working group, European Cooperative Programme for Plant Genetic Resources. Rome, Italy
56. Mo Hill (1973) Diversity and evenness: a unifying notion and its consequences. *Ecol* 54:427–432

[Article Google Scholar](#)

57. Monteiro F, Frese L, Castro S, Duarte MC, Paulo OS, Loureiro J, Romeiras MM (2018) Genetic and genomic tools to assist sugar beet improvement: the value of the crop wild relatives. *Front Plant Sci* 9:74

[PubMed](#) [PubMed Central](#) [Article](#) [Google Scholar](#)

58. Morris EK, Caruso T, Buscot F, Fischer M, Hancock C, Maier TS, Socher SA (2014) Choosing and using diversity indices: insights for ecological applications from the German Biodiversity Exploratories. *Ecol Evol* 4(18):3514–3524

[PubMed](#) [PubMed Central](#) [Article](#) [Google Scholar](#)

59. Nair KP (2019) Utilizing crop wild relatives to combat global warming. In: *Advances in agronomy*, Academic Press, pp 175–258
60. Nevo E, Chen GX (2010) Drought and salt tolerances in wild relatives for wheat and barley improvement. *Plant, Cell Environ* 33:670–685

[CAS](#) [Article](#) [Google Scholar](#)

61. Nóbrega H, Freitas G, Pinheiro de Carvalho MAA (2020) *Beta patula* population assessment from 2014 to 2018 in Madeira Archipelago, Portugal. Version 1.3. University of Madeira. Sampling event dataset <https://doi.org/10.15468/j3qfk5>. Accessed via GBIF.org on 2020-06-09
62. Parra-Quijano M, Iriondo JM, Frese L, Torres E (2012) Spatial and ecogeographic approaches for selecting genetic reserves in Europe. In: Maxted N, Dulloo ME, Ford-Lloyd BV, Frese L, Iriondo JM, Pinheiro de Carvalho MAA (eds) *Agrobiodiversity conservation: securing the diversity of crop wild relatives and landraces*. CAB International, Wallingford, pp 20–28

[Google Scholar](#)

63. Pegot-Espagnet P, Guillaume O, Desprez B et al (2019) Discovery of interesting new polymorphisms in a sugar beet (elite × exotic) progeny by comparison with an elite panel. *Theor Appl Genet* 132(11):3063–3078
-

[CAS](#) [PubMed](#) [PubMed Central](#) [Article](#) [Google Scholar](#)

64. Pinheiro de Carvalho MAA, Frese L (2011) *Beta patula* Aiton genetic reserve action plan. Annex 10 des Abschlussberichtes zum AEGRO-Projekt
65. Pinheiro de Carvalho MAA, Nóbrega H, Frese L, Freitas G, Abreu U, Costa G, Fontinha S (2010) Distribution and abundance of *Beta patula* Aiton and other crop wild relatives of cultivated beets on Madeira. *J Kulturpflanzen* 62(10):357–366
-

[Google Scholar](#)

66. Pinheiro de Carvalho MAA, Nóbrega H, Freitas G, Fontinha S, Frese L (2012) Towards the establishment of a genetic reserve for *Beta patula* Aiton. In: Maxted N, Lothar F, Iriondo J, Dulloo E, Ford-Lloyd BV, Pinheiro de Carvalh MAA (eds) *Agrobiodiversity conservation: securing the diversity of crop wild relatives and landraces*. CAB International, Wallingford, pp 36–44
-

[Google Scholar](#)

67. Pinheiro de Carvalho MAA, Bebeli PJ, Barata da Silva A, Bettencourt E, Slaski JJ, Dias S (2016) Agro-biodiversity. the importance of inventories in the assessment of crop diversity and its time and spatial changes. In: Ahuja M, Jain SM (eds) *Genetic diversity and erosion in plants, sustainable development and biodiversity*. Springer, Heidelberg, pp 130–147
-

[Google Scholar](#)

68. Pinto R, Silva Câmara EM, Melo Ferreira MA (1992) *Carta dos Solos da Ilha da Madeira*. Gov Regional, Sec Reg. Economia, DRA
69. Pires P, Fontinha S (2008) *Plantas da Ponta de São Lourenço*. Edição do Serviço do Parque Natural da Madeira, Funchal, Portugal
70. Prescott-Allen C, Prescott-Allen R (1986) *The first resource*. Yale University Press, Yale
-

[Google Scholar](#)

71. Roberge JM, Angelstam PER (2004) Usefulness of the umbrella species concept as a conservation tool. *Conserv Biol* 18(1):76–85

[Article Google Scholar](#)

72. Romeiras MM, Vieira A, Silva DN et al (2016) Evolutionary and biogeographic insights on the macaronesian beta-*Patellifolia* species (Amaranthaceae) from a time-scaled molecular. *PLoS ONE* 11(3):1–17

[Article CAS Google Scholar](#)

73. Schliephake E, Graichen K, Rabenstein F (2000) Investigations on the vector transmission of the beet mild yellowing virus (BMYV) and the Turnip yellows virus (TuYV). *J Plant Dis Protect* 107(1):81–87

[Google Scholar](#)

74. Schliephake E, Habekuß A, Frese L (2010) Resistance of *Beta patula* Ait to Beet mild yellowing virus. In: CWR & Landraces in Europe: towards the establishment of genetic reserves for crop wild relatives and landraces in Europe' and the joint meeting of the ECPGR in situ and On-farm Conservation Network and the EU project AGRI GENRES 057–AEGRO. University of Madeira, Funchal

75. Seiler GJ, Qi LL, Marek LF (2017) Utilization of sunflower crop wild relatives for cultivated sunflower improvement. *Crop Sci* 57(3):1083–1101

[Article Google Scholar](#)

76. Spellerberg IF, Fedor PJ (2003) A tribute to Claude Shannon and a plea for more rigorous use of species richness, species diversity and the 'Shannon–Wiener' Index. *Glob Ecol Biogeogr* 12:177–179

[Article Google Scholar](#)

77. Stevanato P, Chiodi C, Broccanello C, Concheri G, Biancardi E, Pavli O, Skaracis G (2019) Sustainability of the sugar beet crop. In: *Sugar technology*, pp 1–14

78. Stevens M, Hallsworth PB, Smith HG (2004) The effects of Beet mild yellowing virus and Beet chlorosis virus on the yield of UK field-grown sugar beet in 1997, 1999 and 2000. *Ann Appl Biol* 144(1):113–119
-

[Article](#) [Google Scholar](#)

79. Traill LW, Brook BW, Frankham RR, Bradshaw CJ (2010) Pragmatic population viability targets in a rapidly changing world. *Biol Conserv* 143(1):28–34
-

[Article](#) [Google Scholar](#)

80. Urbaniak GC, Pious S (2011) Research randomizer. Social psychology network. <https://www.randomizer.org/>
81. Vincent H, Amri A, Castañeda-Álvarez NP, Dempewolf H, Dulloo E, Guarino L, Maxted N (2019) Modeling of crop wild relative species identifies areas globally for in situ conservation. *Commun Biol* 2(1):136
-

[PubMed](#) [PubMed Central](#) [Article](#) [Google Scholar](#)

82. Waples RS (2016) Tiny estimates of the N_e/N ratio in marine fishes: Are they real? *J Fish Biol* 89(6):2479–2504
-

[CAS](#) [PubMed](#) [Article](#) [PubMed Central](#) [Google Scholar](#)

83. Warschefsky E, Penmetsa RV, Cook DR, Von Wettberg EJ (2014) Back to the wilds: tapping evolutionary adaptations for resilient crops through systematic hybridization with crop wild relatives. *Am J Bot* 101(10):1791–1800
-

[PubMed](#) [Article](#) [PubMed Central](#) [Google Scholar](#)

84. Wricke G (1972) *Populationsgenetik*. Walter de Gruyter, Berlin
-

[Google Scholar](#)

85. Zhang H, Mittal N, Leamy LJ, Barazani O, Song BH (2017) Back into the wild-apply untapped genetic diversity of wild relatives for crop improvement. *Evol Appl* 10(1):5–24
-

Acknowledgements

The authors acknowledge the Forestry services and rangers of the Institute of Forests and Nature Conservation of Madeira for their support and assistance during the team fieldwork.

Funding

The authors acknowledge the financial support of this work that was provided by the funding programs Life and FEDER, Madeiran Operational Program, through the projects LIFE RECOVER NATURA—LIFE12 NAT/PT/000195 and CASBio, M1420-01-0145-FEDER-000011, in the collection, analysis, and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

Author information

Affiliations

- 1. ISOPlexis Center, University of Madeira, Campus da Penteada, 9020-105, Funchal, Portugal**
H. Nóbrega, G. Freitas, M. A. Zavattieri, C. Ragonezi & M. A. A. Pinheiro de Carvalho
- 2. Biology Department, Colégio da Mitra, University of Évora, Ap. 94, 7002-554, Évora, Portugal**
M. A. Zavattieri
- 3. Institute of Earth Sciences (ICT), Colégio Luís António Verney, University of Évora, Rua Romão Ramalho 59. 7002-554, Évora, Portugal**
M. A. Zavattieri
- 4. Julius Kühn-Institut (JKI), Bundesforschungsinstitut für Kulturpflanzen, Institut für Züchtungsforschung an landwirtschaftlichen Kulturen, Erwin-Baur-Str. 27, 06484, Quedlinburg, Germany**
L. Frese
- 5. Faculty of Life Sciences, University of Madeira, Campus da Penteada, 9020-105, Funchal, Portugal**

M. A. A. Pinheiro de Carvalho

Corresponding author

Correspondence to [C. Ragonezi](#).