

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/344031313>

# Cork Oak Vegetation Series of Southwestern Iberian Peninsula: Diversity and Ecosystem Services

Chapter · January 2021

DOI: 10.1007/978-3-030-48279-4\_119

CITATIONS

0

READS

103

7 authors, including:



**Ricardo Quinto-Canas**

Universidade do Algarve

23 PUBLICATIONS 88 CITATIONS

SEE PROFILE



**Mauro Raposo**

Universidade de Évora

25 PUBLICATIONS 18 CITATIONS

SEE PROFILE



**José Carlos Piñar Fuentes**

Universidad de Jaén

26 PUBLICATIONS 100 CITATIONS

SEE PROFILE



**Eusébio Cano**

Universidad de Jaén

229 PUBLICATIONS 1,720 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Special Issue "Plant Diversity Conservation and Rocky Habitats" on Diversity (MDPI) [View project](#)



OTALEX C [View project](#)



# Cork Oak Vegetation Series of Southwestern Iberian Peninsula: Diversity and Ecosystem Services

Ricardo Quinto-Canas<sup>1,2</sup> , Ana Cano-Ortiz<sup>3</sup> , Mauro Raposo<sup>4</sup>,  
José Carlos Piñar Fuentes<sup>3</sup> , Eusebio Cano<sup>3</sup> , Neuza Barbosa<sup>1</sup>,  
and Carlos José Pinto Gomes<sup>4</sup> 

<sup>1</sup> Faculty of Sciences and Technology, University of Algarve,  
Campus de Gambelas, 8005-139 Faro, Portugal  
r.jcanas@ualg.pt

<sup>2</sup> CCMAR – Centre of Marine Sciences (CCMAR), University of Algarve,  
Campus de Gambelas, 8005-139 Faro, Portugal

<sup>3</sup> Department of Animal and Plant Biology and Ecology, Section of Botany,  
University of Jaén, Las Lagunillas s/n, 23071 Jaén, Spain

<sup>4</sup> Department of Landscape, Environment and Planning, MED - Mediterranean  
Institute for Agriculture, Environment and Development, School of Science  
and Technology, University of Évora (Portugal), Rua Romão Ramalho, nº 59,  
7000-671 Évora, Portugal

**Abstract.** Currently, the occurrence of thermophile cork oak communities is becoming exceptionally rare, which can be attributed to the long-lasting impact of human agroforestry and grazing land-use practices. In this work we identified the thermophilous cork oak vegetation series of southwestern of the Iberian Peninsula and present an overview of ecologic factors and processes controlling the diversity of plant communities observed in their dynamic, as well as, the characteristic vascular flora, including species with special conservation interest, is here presented. This information allows the development of sustainable management, which may enhance both biodiversity and conservation. Moreover, the sustainable economic activities of the cork oak forests, regarding to enhance their capacity to provide benefits that are essential for ecosystem services is developed and presented.

**Keywords:** *Quercus suber* · Thermophilous cork oak vegetation series · Phytosociology · Sustainable management of cork oak forests

## 1 Introduction

Cork oak (*Quercus suber* L.) woodlands are distributed widely across the western and central parts of the Mediterranean basin, mostly on silicate substrates. In southwest of the Iberian Peninsula, cork oak woodlands have high biodiversity and support a wide range of plant communities and endangered species. Moreover, cork oak forests generate substantial economic benefits: from cork and firewood, to a framework tree for agroforestry and silvopastoral systems [1]. Nevertheless, large areas of cork oak forest

landscape, open farmland and shrubland turned into new forest plantations (mainly, *Eucalyptus sp.* and *Pinus sp.pl.*) modern permanent-crop farming systems [2], fires, pathogens, overgrazing and, therefore, have drastically affected the sustainability of Mediterranean woodland ecosystems, leading to their classification as at-risk native biodiversity hotspots [3]. For the correct management and conservation of biodiversity it is important to diagnose the cork oak vegetation series, as well as their dynamics and successional stages, which includes important habitats for many plant species of conservation importance. Hence, we provide a condensed description of the cork oak climatophilous series associated with the thermophilous areas of the southwestern Iberian Peninsula, already described within the *Quercus rotundifoliae-Oleion sylvestris* alliance. In addition to supporting biodiversity, cork oak woodlands provide important ecosystem services and sustainable economic activities based on renewable flow management, since this system can regenerate cork and other useful products with economic value. Therefore, we also present the importance to promote sustainable management of cork oak forests, and halt and reverse land degradation and halt biodiversity loss, and protect the extinction of threatened species. The particular objectives were: 1) to present the cork oak vegetation series, from an ecological and dynamic point of view; 2) to identify the habitats of community interest through the correspondence between plant communities and Directive Habitats types; 3) to indicate species with special conservation interest; 4) to present the sustainable economic activities of the cork oak forests, in order to enhance their capacity to provide benefits that are essential for both, sustainable development and ecosystems, and their services.

### **Study Area**

The research area covers the thermophilous territories with acidic soils of the southwestern Iberian Peninsula, linked to the sublittoral areas and the valley of the rivers Tagus and Sado. According to the most recent study on Iberian Peninsula bioclimatic classification by [4], these territories are classified as Mediterranean pluviseasonal oceanic, mostly thermomediterranean hyperoceanic to euoceanic. In a biogeographical context and following [5], the study area is included in the Coastal Lusitania and West Andalusian Province and westernmost part of the Lusitania and Extremadura Sub-province (West Iberian Mediterranean Province).


### **Materials and Methods**

The climatophilous series of *Q. suber* diagnosis followed past works from [6–26]. Sintaxonomical nomenclature followed [27–29]. Taxonomic nomenclature follows [29, 30]. The classification of habitats, through the evaluation of the dominant vegetation types and their diagnostic species, follows [31]. The biogeographical and bioclimatological analysis was carried out according to [4, 5]. Generally, the information of ecosystem services and sustainable based on the available published works, such as [1, 45, 47–49].

### **Discussion and Results**

The set of vegetation types which, by virtue of succession replace each other at a particular site has been called a series [42]. We can summarize the typical sequence of

stages occurring in the cork oak vegetation series from southwestern Iberian Peninsula (See Fig. 1), within the limits of a tessela, which includes the whole forest successional sequence, depending on its degree of degradation [26, 50–55].



**Fig. 1.** Typical sequence of stages occurring in the cork oak vegetation series from southwest of Iberian Peninsula: 1 - Cork oak woodland; 2 - Maquis scrubland; 3 - Broomland; 4 - Perennial grassland; 5 - Gorse scrubland; 6 - Cistaceous scrubland; 7 - Perennial grassland; 8 - Annual grasslands (adapted from [56]).

Therefore, the four *Q. suber* woodland series identified in the study area is presented briefly, highlighting the successional stages, as well as the natural or semi-natural habitats that it may incorporate and its asset value (Table 1).

#### A. *Lavandulo viridis-Quercus suberis* sigmetum

Vegetation series occurring along the mountainous areas of the southwest of Portugal mainland (Caldeirão, Monchique, Cercal, Grândola) (Monchique Sierran District), reaching the Aracena mountain (Aracena Sierran District), typical from silicates substrates, derived from schists and greywackes, in the thermomediterranean, hyperoceanic, lower sub-humid bioclimate areas. The mature stage *Lavandulo viridis-Quercetum suberis*, is characterized by the dominance of *Quercus suber* and the constant presence of *Lavandula viridis*, a southwestern Iberian endemic, considered a characteristic species of this association. Its fringe belongs to the *Arbutus unedo* maquis from *Cisto populifolii-Arbutetum unedonis*. Nevertheless, the removal of tree and shrub cover, leads to the *Cytisus striatus* broomlands (*Lavandulo viridis-Cytisetum striati*). In contrast, with the soil degradation and following the regressive dynamic, occurs the heathland from *Ulici argentei-Ericetum australis* and a gorse scrubland of *Cisto ladaniferi-Ulicetum argentei*, both exclusive to the Monchique Sierran District and well characterized by the endemic *Ulex argenteus*. In woodland clearings, the perennial grasslands

**Table 1.** Biogeographical, substratum, Habitats Directive, plant communities and species with conservation interest of the thermophilous cork oak vegetation series of southwestern the Iberian Peninsula (within the *Quercus rotundifoliae-Oleion sylvestris* alliance).

Climatophilous series	Woodland and successional stages	Biogeography	Habitats Directive	Substratum affinity	Main species with conservation interest
<i>Lavandulo viridis-Quercus suberis</i> sigmetum	<i>Lavandulo viridis-Quercetum suberis</i> ; <i>Cisto populifolii-Arbutetum unedonis</i> ; <i>Lavandulo viridis-Cytisetum striati</i> ; <i>Ulici argentei-Ericetum australis</i> ; <i>Cisto ladamiferi-Ulicetum argentei</i> ; community of <i>Dacrylis hispanica</i> subsp. <i>lusitanica</i> ; <i>Trifolio chertieri-Plantaginietum bellardii</i> ; <i>Holco annui-Brachypodiëtum distachyi</i>	Monchique Sierran and Aracena Sierran Districts	9330, 5330, *6220, 4030	Schists and greywackes	<i>Doronicum plantagineum</i> subsp. <i>tournefortii</i> (Annex V Habitats Directive), <i>Narcissus bulbocodium</i> (Annex V Habitats Directive), <i>Drosophyllum lusitanicum</i> , <i>Gagea lusitanica</i> , <i>Senecio minutus</i> , <i>Centaurea crocata</i>
<i>Aro neglecti-Quercus suberis</i> sigmetum	<i>Aro neglecti-Quercetum suberis</i> ; <i>Phillyreo angustifoliae-Arbutetum unedonis</i> ; <i>Cytisetum cabezudo</i> ; <i>Avenulo hackelii-Celticetum sterilis</i> ; <i>Armerio macrophyllae-Celticetum giganteae</i> ; <i>Euphorbio transaganae-Celticetum giganteae</i> ; <i>Hyacinthoides transaganae-Brachypodiëtum phoenicoidis</i> ; <i>Erico umbellatae-Ulicetum welwitschiani</i> ; <i>Halimio halimifolii-Stauracanthetum gemistoidis</i> ; <i>Genisto triacanthi-Stauracanthetum vicentini</i> ; <i>Tuberario majoris-Stauracanthetum botvini</i> ; <i>Thymo camphorati-Stauracanthetum spectabilis</i> ; <i>Cistetum libanotidis</i> ; Communities of <i>Corynephorus canescens</i> var. <i>maritimus</i> ; <i>Tolpido barbatae-Tuberarietum bupleurifoliae</i>	Coastal Lusitania and West Andalusian Province	*2150, 2230, 2260, 2330, 4030, 5330, *6220, 9330	Sandy soils (paleodunes)	i) *priority species from Annex II Directive Habitats: * <i>Thymus camphoratus</i> ; * <i>Thymus lotocephalus</i> ; * <i>Tuberaria major</i> ; ii) Annex II Habitats Directive: <i>Armeria rotuyana</i> , <i>Armeria velutina</i> , <i>Euphorbia transtaganana</i> , <i>Halimium verticillatum</i> , <i>Hyacinthoides vicentinas</i> subsp. <i>transtaganana</i> , <i>Thymus camphoratus</i> , <i>Santolina impressa</i> ; iii) Annex IV Habitats Directive: <i>Thymus caprellatus</i> , <i>Thymus villosus</i> ; iv) Annex V Habitats Directive: <i>Malcolmia laevis</i> subsp. <i>gracillima</i> , <i>Narcissus bulbocodium</i> ; v) other species: <i>Arenaria algarbiensis</i> , <i>Armeria macrophylla</i> , <i>Armeria apinifolia</i> , <i>Centaurea exarata</i> , <i>Dianthus broteri</i> subsp. <i>hinoxianus</i> , <i>Klasea algarbiensis</i> , <i>Lupinus cosentinii</i> , <i>Ononis cossoniana</i> , <i>Scilla odorata</i> , <i>Thesium humile</i> , <i>Thymus albicans</i> subsp. <i>albicans</i> , <i>Thymus albicans</i> subsp. <i>donyanae</i> , <i>Trisetaria duforei</i> , <i>Ulex argenteus</i> subsp. <i>subsericeus</i>

(continued)

Table 1. (continued)

Climatophilous series	Woodland and successional stages	Biogeography	Habitats Directive	Substratum affinity	Main species with conservation interest
<i>Asparago aphylli-Quercu suberis</i> <i>angustifoliae-Arbutetum unedonis</i> , <i>Erico scopariae-Quercetum lusitanicae</i> , <i>Erico umbellatae-Ulacetum welwitschianii</i> , <i>Lavandulo luisieri-Ulacetum jussiaei</i> , <i>Thymo villosae-Ulacetum airenis</i> , <i>Avenulo sulcatae-Celticetum giganteae</i>	Iberian Atlantic Mediter-ranean biogeographic territory	9330, 5330, *6220, 4030	Schist and sandstone	<i>Halimium verticillatum</i> (Annex II Habitats Directive), <i>Thymus villosus</i> (Annex IV Habitats Directive), <i>Drosophyllum lusitanicum</i>	
<i>Smilaco asperae-Quercu suberis</i> <i>angustifoliae-Arbutetum unedonis</i> , <i>Cytisetum multifloro-ericocampi</i> , <i>Erico australis-Cistetum populifolii</i> , <i>Halimio ocymoidis-Ericetum umbellatae</i> , <i>Avenulo sulcatae-Celticetum giganteae</i>	Oretana Range and Tajo Sector	9330, 5330, *6220, 4030	Schists and quartzites	<i>Ruscus aculeatus</i> (Annex V Habitats Directive), <i>Juniperus oxycedrus</i> subsp. <i>lagunae</i> , <i>Centaurea coutinhoi</i> , <i>Iris lusitanica</i>	

dominated by *Dactylis hispanica* subsp. *lusitanica* frequently occur where as ground clearance leads to the successive appearance of therophytic grasslands from *Tuberarion guttatae*. The *Picrido spiniferae-Cynarietum algarbiensis*, dominated by *Cynara algarbiensis*, represents the forest fringe of *Lavandulo viridis-Quercetum suberis*.

B. *Aro neglecti-Quercus suberis* sigmetum

According to [23], along the thermomediterranean, dry to subhumid bioclimatic belts of the Coastal Lusitania and West Andalusian Province, occurs the *Aro neglecti-Quercetum suberis*. This psammophilous cork oak forests develops on podzol sandy soils derived from Pleistocene paleodunes [26], and is dominated by *Quercus suber* and normally accompanied by *Arum italicum* subsp. *neglectum*, *Pistacia lentiscus* and *Stauracanthus genistoides* [9]. The fringe which occurs above sandy soils belongs to the strawberry tree from *Phillyrea angustifoliae-Arbutetum unedonis*. The elimination of woody and shrubby strata leads to the broomland from *Cytisetum cabezudo*, dominated by *Cytisus grandiflorus* subsp. *cabezudo*. Shifting to deeper soils, in forests and shrub clearings, occurs the perennial grasslands from *Agrostio castellanae-Celticion giganteae* alliance (*Avenulo hackelii-Celticetum sterilis*, *Armerio macrophyllae-Celticetum giganteae*, *Euphorbio transtaganae-Celticetum giganteae*) [13, 16] and the orchid-rich grasslands of *Brachypodium phoenicoides* (*Hyacinthoides transtaganae-Brachypodietum phoenicoidis*) [25]. In the regressive dynamic, the domain is separate from *Erico umbellatae-Ulicetum welwitschiani* heathlands, and a psammophilous community of *Halimio halimifolii-Stauracanthetum genistoidis*. For the San Vicente Coastal District, occurs the gorse scrubland of *Genisto triacanthi-Stauracanthetum vicentini*, with *Stauracanthus spectabilis* subsp. *vicentinus* as an important endemic element. In sandy soils with clay we observe the gorse scrubland from *Tuberario majoris-Stauracanthetum boivinii*, mainly in palaeopodzolic ferruginous lithosols, subject to water logging. Through its shift, in the San Vicente Coastal and San Vicente Cape Districts, in sandy soils without the ferruginous horizon, coastal gorse scrublands of *Thymo camphorate-Stauracanthetum spectabilis* emerge and also the psammophilous community of *Cistetum bourgaeani* (*libanotidis*). With more degradation, occurs the perennial grasslands dominated by *Corynephorus canescens* var. *maritimus* from *Corynephorion maritime* alliance, and the therophytic communities of *Tolpido barbatae-Tuberarietum bupleurifoliae* occur.

C. *Asparago aphylli-Quercus suberis* sigmetum

This climatophilous series occurs in the Iberian Atlantic Mediterranean biogeographic territory, on siliceous substrata (schist and sandstone). The climatic domain corresponds to a cork oak forest from *Asparago aphylli-Quercetum suberis*. The pre-forestall stage is dominated by *Arbutus unedo* and *Phillyrea angustifolia* (*Phillyrea angustifoliae-Arbutetum unedonis*). As this scrubland regresses, occurs the *Erico scopariae-Quercetum lusitanicae*. According to [26] the seral stage also includes gorse/heathland dominated by *Ulex australis* subsp. *welwitschianus* (*Erico umbellatae-Ulicetum welwitschianii*; low Tagus and Sado), *Ulex jussiaei* (*Lavandulo luisieri-Ulicetum jussiaei*; north of the Tagus) and *Ulex airensis*

(*Thymo villosae-Ulicetum airensis*). In woodland clearings, occurs the perennial grasslands dominated by *Celtica gigantea* (*Avenulo sulcatae-Celticetum giganteae*). The *Stachydo lusitanicae-Origanetum virentis* association represents the forest fringe community of *Asparago aphyllus-Quercetum suberis*.

D. *Smilaco asperae-Quercu suberis* sigmetum

This climatophilous series develops on siliceous substrata (schists and quartzites) of the lower subhumid areas of the central Tagus valley (Oretana Range and Tajo Sector). The climatic stage belongs to the *Smilaco asperae-Quercetum suberis*, characterized by the presence of *Juniperus oxycedrus* subsp. *lagunae*. The *Phillyreo angustifoliae-Arbutetum unedonis* represents the first substitution layer. The elimination of this shrubby strata leads to the *Cytisetum multifloro-eriocarpi*. Soil degradation promotes the development of the heathlands: *Erico australis-Cistetum populifolii* and *Halimio ocymoidis-Ericetum umbellatae*. In the woodland clearings, occurs the perennial grasslands from *Avenulo sulcatae-Celticetum giganteae*.




Fig. 2. Ecosystem services of cork oak forests and ecological economics tools diagram. (adapted from Ellen MacArthur Foundation [57]).

According to [32], as a direct result of human management, functional cork oak landscapes present a wide array of forest or woodland, shrubland, and open habitat components. As shown in the Table 1, a total of 8 habitats associated to European Habitats Directive (Directive 92/43/EEC), are recognized and, therefore, have a large set of rare, endemic, endangered and protected species under the Directive 92/43/EEC



(about 41 species with conservation interest). Besides the high biodiversity value, the cork oak vegetation series provide, from an ecological perspective, important ecosystem services, mainly associated to soil fertility, water retention, watershed protection, erosion control, fire risk prevention and carbon sequestration [32] (See Fig. 2). Among these factors, the capacity of cork oak woodland to contribute to carbon mitigation targets must be highlighted. In fact, several studies in Portugal reveal that they store a significant amount of carbon (reaching 14.7 tons of CO<sub>2</sub> per year per hectare) [36] in standing biomass (cork oaks) and soil. Despite the ecological importance, the cork oak forests are one of the few examples of fully sustainable forestry exploitations, comprising multiple land uses, including cork, acorn, firewood, fruits, oils, mushrooms, recreation and tourism activities [1]. Nevertheless, they are economically sustainable mostly to the high market value of cork, where bottle stoppers that drive the cork industry, represent almost 70% of cork's market value [35]. Therefore, several authors evidence that cork, being a natural and renewable material, which in itself is the result of a CO<sub>2</sub> absorption and sequestration, and also a storage of C, can also contribute to the design of the post-carbon city by reducing energy waste, improving the quality and the insulation of buildings [37–49]. Moreover, the amount of carbon sequestered in cork oaks according to [34] is 0.95–1.25 tonne C/tonne cork. Hence, if the cork oak was registered in the “National Register of Agro-Forestry Carbon Tanks”, then it would receive an allowance of 20 €/tonne/year [in 2005 the cost of a tonne of CO<sub>2</sub> fluctuated between 8 and 32 €/tonne/year] (Spampinato et al. 2019) [43].

Currently, cork oak forests are facing complex problems and are under increasing pressure, due to soil mobilization, intensification of agricultural activities, particularly overgrazing (mainly cattle grazing), climate change, which promote diseases, drought and water stress (further endangering viable cork production), forest fires, and inappropriate silvicultural management, that could seriously threatened the long-term of trees [33].

According to [32], to address the problems driving the degradation and loss of cork oak woodlands, there is a clear need to review national forestry education schemes and introduce innovative concepts for sustainable management and biodiversity conservation of cork oak woodlands and the development of sustainable markets, based on responsible management of natural resources.

## 2 Conclusions

We present the first approach to the cork oak vegetation series of southwestern Iberian Peninsula, through which it was possible to perceive the existence of a large diversity of plant communities and habitat type, displaying great value, with several rare endemic plants. Understanding vegetation series is of utmost importance for the diagnosis and determination of the vegetation's current conservation status. This information allows the landscape managers to present some priority measures that must be addressed strategically, to help biodiversity adapt to recent drivers of transformation and degradation, at the same time, accounting for the socio-economic activities that support the sustainable use of the existing natural resources. According to [1], a research goal for the future must be to apply environmental and ecological economics

tools to the valuation of biodiversity conservation and ecosystem services of cork oak woodlands and derived cultural systems. We also recognize that promote the implementation of this valuation in management and planning, is the greatest challenge and an indispensable requirement for sustainable development in its three dimensions – economic, social and environmental.

**Acknowledgments.** Diagram icons made by Freepik from [www.flaticon.com](http://www.flaticon.com).

## References

1. Aronson, J., Pereira, J.S., Pausas, J.G.: Challenges for the future. In: Aronson, J., Pereira, J. S., Pausas, J.G. (eds.) *Cork Oak Woodlands on the Edge. Ecology, Adaptive Management and Restoration*, pp. 201–202. Island Press, Washington, D.C. (2009)
2. Costa, A., Madeira, M., Lima Santos, J.: Is cork oak (*Quercus suber* L.) woodland loss driven by eucalyptus plantation? A case-study in southwestern Portugal. *iForest* **7**, 193–203 (2014)
3. Myers, N., Mittermeier, R.A., Mittermeier, C.G.: Biodiversity hotspots for conservation priorities. *Nature* **403**, 853–858 (2000)
4. Rivas-Martínez, S., Penas, A., Díaz González, T.E., Paloma Cantó, P., Río, S., Costa, J.C., et al.: Biogeographic units of the Iberian Peninsula and Balearic Islands to district level. A concise synopsis. In: Loidi, J. (ed.) *The Vegetation of the Iberian Peninsula. Plant and Vegetation*, vol. 12, pp. 131–188. Springer, Cham (2017)
5. Rivas-Martínez, S., Penas, A., Río, S., Díaz González, T.E., Rivas-Sáenz, S.: Bioclimatology of the Iberian Peninsula and the Balearic Islands. In: Loidi, J. (ed.) *The Vegetation of the Iberian Peninsula. Plant and Vegetation*, vol. 12, pp. 29–80. Springer, Cham (2017)
6. Rivas-Martínez, S., Costa, M., Castroviejo, S., Valdes, E.: Vegetación de Doñana (Huelva, España). *Lazaroa* **2**, 5–190 (1980)
7. Rivas-Martínez, S., Lousã, M., Díaz, T., Fernández-González, F., Costa, J.: La vegetación del sur de Portugal. *Itin. Geob.* **3**, 5–126 (1990)
8. Rivas-Martínez, S.: Brezales y jarales de Europa Occidental (revisión fitosociológica de las clases Calluno-Ulicetea y Cisto-Lavanduletea). *Lazaroa* **1**, 5–127 (1979)
9. Rivas-Martínez, S.: Mapas de series, geoseries y geopermaseries de vegetación de España (Memoria del mapa de vegetación potencial de España). *Itin. Geob.* **18**(1–2), 5–800 (2011)
10. Capelo, J.: *Nemorum Trantaganae Descriptio. Sintaxonomia numerica das comunidades florestais e preforestais do Baixo Alentejo. Dissertação de Doutoramento, Instituto Superior de Agronomia – Universidade Técnica de Lisboa, Lisboa, Portugal* (2007)
11. Costa, J.C., Capelo, J., Lousã, M., Espírito-Santo, M.D.: Os sobreirais do Sector Divisório Português: *Asparago aphylli-Quercetum suberis*. *Quercetea* **3**, 81–96 (2002)
12. Pinto-Gomes, C., Paiva-Ferreira, R.: Flora e vegetação do barrocal algarvio, Tavira-Portimão. Comissão de Coordenação e Desenvolvimento Regional do Algarve (2005)
13. Pinto-Gomes, C., Paiva Ferreira, R., Meireles, C.: New proposals on Portuguese vegetation. *Lazaroa* **28**, 66–77 (2007)
14. Pinto-Gomes, C., Paiva-Ferreira, R., Quinto-Canas, R., Rosa-Pinto, J., Meireles, C., Redondo García, M.M.: Guia Geobotânico ao Barrocal Algarvio. *Quercetea* **8**, 3–143 (2008)
15. Pinto-Gomes, C., Paiva-Ferreira, R., Meireles, C.: Os montados: bases para uma metodologia integrada de exploração, gestão e conservação. *A. Bot. Gallica* **156**(1), 149–151 (2009)

16. Pinto-Gomes, C., Paiva-Ferreira, R., Meireles, C.: New proposals on Portuguese Vegetation (II). *Lazaroa* **31**, 59–65 (2010)
17. Pinto-Gomes, C., Cano-Ortiz, A., Quinto-Canas, R., Vila-Viçosa, C., Martínez Lombardo, M.: Analysis of the *Cytisetia scopario-striati* scrubs in the south-west-centre of the Iberian Peninsula. *A. Bot. Gallica* **159**(2), 251–266 (2012)
18. Pinto-Gomes, C., Ladero, M., Gonçalves, P., Mendes, S., Lopes, M.C.: *Smilaco asperae-Quercetum suberis*: um novo sobreiral relíquo do Alto Tejo. *Quercetea* **4**, 23–29 (2004)
19. Neto, C., Costa, J.C., Capelo, J., Gaspar, N., Monteiro-Henriques, T.: Os sobreirais da Bacia ceno-Antropozóica do Tejo (Província Lusitano-Andalusa Litoral, Portugal). *Acta Bot. Malacitana* **32**, 201–210 (2007)
20. Neto, C., Arsénio, P., Costa, J.C.: Flora e vegetação do Sudoeste de Portugal continental. *Quercetea* **9**, 43–144 (2009)
21. Quinto-Canas, R.: Flora y vegetación de la Serra do Caldeirão. Ph.D. thesis, Facultad Ciencias Experimentales, Universidad de Jaén, Jaén (2014)
22. Quinto-Canas, R., Vila-Viçosa, C., Meireles, C., Paiva-Ferreira, R., Martínez-Lombardo, M., Cano, A., et al.: A contribute to the knowledge of the climatophilous cork-oak woodlands from Iberian southwest. *A. Bot. Gallica* **157**(4), 627–637 (2010)
23. Quinto-Canas, R., Vila-Viçosa, C., Paiva-Ferreira, R., Cano-Ortiz, A., Pinto-Gomes, C.: The Algarve (Portugal) climatophilous vegetation series: a base document to the planning, management and natureconservation. *A. Bot. Gallica* **159**(3), 289–298 (2012)
24. Quinto-Canas, R., Mendes, P., Cano-Ortiz, A., Musarella, C.M., Pinto-Gomes, C.: Forest fringe communities of the southwestern Iberian Peninsula. *Rev. Chapingo Ser. Cie.* **24**(3), 415–434 (2018)
25. Raposo, M., Mendes, P., Cano-Ortiz, A., Pinto-Gomes, C.: Séries de vegetação prioritárias para a conservação no centro e sul de Portugal Continental. *Botanique* **1**, 113–148 (2016)
26. Espírito-Santo, D., Capelo, J., Neto, C., Pinto-Gomes, C., Ribeiro, S., Quinto-Canas, R., Costa, J.C.: Lusitania. In: Loidi, J. (ed.) *The Vegetation of the Iberian Peninsula. Plant and Vegetation*, vol. 13, pp. 35–82. Springer, Cham (2017)
27. Rivas-Martínez, S., Fernández-González, F., Lodi, J., Lousã, M., Penas, A.: Syntaxonomical checklist of vascular plant communities of Spain and Portugal to association level. *Itin. Geob.* **14**, 5–341 (2001)
28. Rivas-Martínez, S., Díaz, T.E., Fernández-González, F., Izco, J., Loidi, J., Lousã, M., et al.: Vascular plant communities of Spain and Portugal: addenda to the syntaxonomical checklist of 2001. *Itin. Geob.* **15**(1, 2), 5–922 (2002)
29. Costa, J.C., Neto, C., Aguiar, C., Capelo, J., Espírito Santo, M.D., Honrado, J., et al.: Vascular plant communities in Portugal (continental, the Azores and Madeira). *Global Geobot. Int. J. Geobot. Sci.* **2**, 1–180 (2012)
30. Castroviejo, S.: Flora Ibérica. Plantas vasculares de la península Ibérica y Baleares (vols. I–VIII, X–XV, XVII, XVIII, XXI). Real Jardín Botánico, CSIC, España (1986–2012)
31. ALFA: Tipos de Habitat Naturais e Semi-Naturais do Anexo I da Directiva 92/43/CEE (Portugal continental): Fichas de Caracterização Ecológica e de Gestão para o Plano Sectorial da Rede Natura 2000. Associação Lusitana de Fitossociologia (2004)
32. Berrahmouni, N., Regato, P., Ellatifi, M., Daly-Hassen, H., Bugalho, M., Bensaid, S., Díaz, M., James Aronson, J.: Ecoregional planning for biodiversity conservation. In: Aronson, J., Pereira, J.S., Pausas, J.G. (eds.) *Cork Oak Woodlands on the Edge. Ecology, Adaptive Management and Restoration*, pp. 203–217. Island Press, Washington, D.C. (2009)
33. Costa, A., Pereira, H., Madeira, M.: Analysis of spatial patterns of oak decline in cork oak woodlands in Mediterranean conditions. *Ann. For. Sci.* **67**(2), 204 (2010)
34. Croezen, A., Bijleveld, M., Sevenster, M.: Natural cork bottle stoppers: a stopper on CO<sub>2</sub> emissions? Report, Delft, CE Delft (2013). 69 p.

35. APCOR: Cork – Environmental Importance. Cork Information Bureau (2015). 16 p.
36. APCOR: Cork – Environmental Importance. Cork Information Bureau (2019). 18 p.
37. Del Giudice, V., Massimo, D.E., De Paola, P., Forte, F., Musolino, M., Malerba, A.: Post carbon city and real estate market: testing the dataset of Reggio Calabria market using spline smoothing semiparametric method. In: Calabrò, F., Della Spina, L., Bevilacqua, C. (eds.) *New Metropolitan Perspectives: Local Knowledge and Innovation. Smart Innovation, Systems and Technologies*, vol. 100, pp. 206–214. Springer, Heidelberg (2019)
38. De Paola, P., Del Giudice, V., Massimo, D.E., Forte, F., Musolino, M., Malerba, A.: Isovalore maps for the spatial analysis of real estate market: a case study for a central urban area of Reggio Calabria, Italy. In: Calabrò, F., Della Spina, L., Bevilacqua, C. (eds.) *New Metropolitan Perspectives: Local Knowledge and Innovation. Smart Innovation, Systems and Technologies*, vol. 100, pp. 402–410. Springer, Heidelberg (2019)
39. Malerba, A., Massimo, D.E., Musolino, M., Nicoletti, F., De Paola, P.: Post carbon city: building valuation and energy performance simulation programs. In: Calabrò, F., Della Spina, L., Bevilacqua, C. (eds.) *New Metropolitan Perspectives: Local Knowledge and Innovation. Smart Innovation, Systems and Technologies*, vol. 101, pp. 513–531. Springer, Heidelberg (2019)
40. Massimo, D.E., Del Giudice, V., De Paola, P., Forte, F., Musolino, M., Malerba, A.: Geographically weighted regression for the post carbon city and real estate market analysis: a case study. In: Calabrò, F., Della Spina, L., Bevilacqua, C. (eds.) *New Metropolitan Perspectives: Local Knowledge and Innovation. Smart Innovation, Systems and Technologies*, vol. 100, pp. 142–149. Springer, Heidelberg (2019)
41. Massimo, D.E., Malerba, A., Musolino, M., Nicoletti, F., De Paola, P.: Valutazione energetica comparativa degli edifici per la Post Carbon City [Energy Comparative Assessment of Buildings, for the Post Carbon City], pp. 63–70. Laborest (2019)
42. Musolino, M., Malerba, A., De Paola, P., Musarella, C.M.: Building Efficiency Adopting Ecological Materials and Bio Architecture Techniques, pp. 707–717. ArcRHistoR (2019)
43. Spampinato, G., Massimo, D.E., Musarella, C.M., De Paola, P., Malerba, A., Musolino, M.: Carbon sequestration by cork oak forests and raw material to built up post carbon city. In: Calabrò, F., Della Spina, L., Bevilacqua, C. (eds.) *New Metropolitan Perspectives: Local Knowledge and Innovation. Smart Innovation, Systems and Technologies*, vol. 101, pp. 663–671. Springer, Heidelberg (2019)
44. Massimo, D.E., Malerba, A., Musolino, M.: Green district to save the planet. In: Mondini, G., Fattinanzi, E., Oppio, A., Bottero, M., Stanghellini, S. (eds.) *Integrated Evaluation for the Management of Contemporary Cities. Green Energy and Technology*, pp. 255–269. Springer, Heidelberg (2018)
45. Massimo, D.E., Malerba, A., Musolino, M.: Valuating historic centers to save the planet soil. In: Mondini, G., Fattinanzi, E., Oppio, A., Bottero, M., Stanghellini, S. (eds.) *Integrated Evaluation for the Management of Contemporary Cities. Green Energy and Technology*, pp. 297–311. Springer, Heidelberg (2018)
46. Massimo, D.E., Malerba, A., Musolino, M., et al.: A green district to save the planet. In: Mondini, G., Fattinanzi, E., Oppio, A., Bottero, M., Stanghellini, S. (eds.) *Integrated Evaluation for the Management of Contemporary Cities. Green Energy and Technology*, pp. 255–269. Springer, Cham (2017)
47. Massimo, D.E., Fragomeni, C., Malerba, A., Musolino, M.: Valuation supports green university: case action at Mediterranean campus in Reggio Calabria. *Procedia Soc. Behav. Sci.* **223**, 17–24 (2016)

48. Massimo, D.E.: Green building: characteristics, energy implications and environmental impacts. Case study in Reggio Calabria. In: Mildred, C.-S. (ed.) *Green Building and Phase Change Materials: Characteristics, Energy Implications and Environmental Impacts*. Energy Science, Engineering and Technology, pp. 71–101. Nova Science Publishers, Inc., New York (2015)
49. Massimo, D.E., Battaglia, L., Fragomeni, C., Guidara, M., Rudi, G., Scala, C.: Sustainability valuation for urban regeneration-the “Geomatic Valuation University Lab” research. In: *Advanced Engineering Forum*. TTP Publications, USA (2014)
50. Loidi, J.: Dynamism in vegetation. Vegetation changes on a short time scale. In: Loidi, J. (ed.) *The Vegetation of the Iberian Peninsula*. Plant and Vegetation, vol. 13, pp. 35–82. Springer, Cham (2017)
51. Piñar Fuentes, J.C., Cano-Ortiz, A., Musarella, C.M., Quinto Canas, R., Pinto Gomes, C.J., Spampinato, G., del Río, S., Cano E.: Bioclimatology, structure, and conservation perspectives of *Quercus pyrenaica*, *Acer opalus* subsp. *granatensis*, and *Corylus avellana* deciduous forests on Mediterranean bioclimate in the South-Central part of the Iberian Peninsula. *Sustainability* **11**, 6500 (2019)
52. Cano Ortiz, A., Pinto Gomes, C.J., Musarella, C.M., Cano, E.: Expansion of the *Juniperus* genus due to anthropic activity. In: Weber, R.P. (ed.) *Old-Growth Forest and Coniferous Forests*, pp. 55–65. Nova Science Publishers, New York (2015)
53. Cano, E., Musarella, C.M., Cano Ortiz, A., Piñar Fuentes, J.C., Rodríguez Torres, A., Del Río González, S., Pinto Gomes, C.J., Quinto Canas, R., Spampinato, G.: Geobotanical study of the microforests of *Juniperus oxycedrus* subsp. *badia* in the Central and Southern Iberian Peninsula. *Sustainability* **11**(4), 1111 (2019)
54. Mendes, P., Meireles, C., Vila-Viçosa, C., Musarella, C.M., Pinto-Gomes, C.: Best management practices to face degraded territories occupied by *Cistus ladanifer* shrublands – Portugal case study. *Plant Biosyst.* **149**(3), 494–502 (2015)
55. Piñar Fuentes, J.C., Cano-Ortiz, A., Musarella, C.M., Pinto Gomes, C.J., Spampinato, G., Cano, E.: Rupicolous habitats of interest for conservation in the central-southern Iberian Peninsula. *Plant Sociol.* **54**((2), Suppl. 1), 29–42 (2017)
56. Barbosa, N.: A experiência profissional do atelier greenflux e contributos para a adoção de práticas sustentáveis de projeto. Tese de Mestrado. Universidade do Algarve (2018). 164 p.
57. Ellen MacArthur Foundation: *Growth Within: A Circular Economy Vision for a Competitive Europe*. SUN and McKinsey Center for Business and Environment (2015)