

Oral Communication 10

Potato growth under environment chamber conditions

Pedro Barbosa<sup>1</sup>, Rui Machado<sup>2</sup>, Renato Coelho<sup>3</sup>, Cláudia Vicente<sup>1,4</sup>, A. Cristina Figueiredo<sup>5</sup> and Manuel Mota<sup>1,3,6</sup>

<sup>1</sup>NemaLab/MED – Mediterranean Institute for Agriculture, Environment and Development, Instituto de Investigação

e Formação Avançada, Universidade de Évora, Pólo da Mitra, Ap. 94, 7006-554 Évora, Portugal.

<sup>2</sup>MED – Mediterranean Institute for Agriculture, Environment and Development & Departamento de Fitotecnia, Escola

de Ciências e Tecnologia, Universidade de Évora, Pólo da Mitra, Ap. 94, 7006-554 Évora, Portugal.

<sup>3</sup>MED – Mediterranean Institute for Agriculture, Environment and Development & Departamento de Biologia, Escola de Ciências e Tecnologia, Universidade de Évora, Pólo da Mitra, Ap. 94, 7006-554 Évora, Portugal.

<sup>4</sup>Instituto Nacional de Investigação Agrária e Veterinária (INIAV, I.P.), Quinta do Marquês, Oeiras, Portugal. <sup>5</sup>Centro de Estudos do Ambiente e do Mar (CESAM Lisboa), Faculdade de Ciências da Universidade de Lisboa, Centro de Biotecnologia Vegetal (CBV), DBV, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal.

<sup>6</sup>Departamento Ciências da Vida, Universidade Lusófona de Humanidades e Tecnologias, EPCV, Campo Grande 376, 1749-024 Lisboa, Portugal.

Email: pbarbosa@uevora.pt

Potato (*Solanum tuberosum* L.) is one of the world's most common vegetable, source of calories, essential nutrients and vitamin C. Most studies with potatoes are performed in vitro, under strict environmental conditions and high levels of asepsis. Integrated in a larger research study that aims at simulating in vivo conditions for application of nematicides for potato pest control, this study intended to evaluate the best growth chamber conditions for potato growth. To optimize the best conditions, type of substrate, potato variety, seed depth, day/ night period, environment conditions and watering/ fertilization routine were evaluated between September 2018 and February 2020 in 14 trials with 20 plants each. In March and in June 2020, 16 potato seeds were placed in 3 L plastic pots, filled with pine forest sand and composted pine bark (9:1 ratio). Plants were grown in an Aralab© D1200PLH chamber at MED/ U. Évora facilities. Day period was set to 14 h, 23°C and 50% relative humidity and night period to 10 h, 12°C and 70% relative humidity. Plants were routinely watered and fertilized. Destructive sampling was performed 7, 14, 21 and 28 days after plant emergence (DAE). Non-destructive sampling was measured every 2 days, starting on the 7th day after emergence. Biomass related parameters (height, fresh weight, leaf area, etc.) increased as plant develops, with some overlay between consecutive time-points. Stomatal conductance, chlorophyll content and chlorophyll fluorescence values are more variable as plants grow. A plant with 14 DAE has a maximum quantum efficiency (FV/FM) value of 0.82 and an average Performance Index (PIabs) of 7.91; a

plant with 28 DAE has an average FV/FM value of 0.83 and an average PIabs of 8.10. Leaf chlorophyll content varied between 73.66 for a 14 DAE plant and 80.20 for the 28 DAE plant (SPAD index).

This work is funded by National Funds through FCT under the PhD grant SFRH/BD/134201/2017, PTDC/AGR PRO/2589/2014 and UID/AGR/00115/2013, CESAM UIDP/50017/2020 + UIDB/50017/2020, FEDER PT2020-Compete 2020.