

A review on the nematocidal activity of volatile allelochemicals against the pinewood nematode †

Jorge M. S. Faria ^{1,*}, Pedro Barbosa ¹, Dora Martins Teixeira ^{2,3} and Manuel Mota ^{1,3}

¹ NemaLab-MED, Mediterranean Institute for Agriculture, Environment and Development, Institute for Advanced Studies and Research, Évora University, Pólo da Mitra, Ap. 94, 7006-554 Évora, Portugal.

² HERCULES Laboratory, Évora University, Largo Marquês de Marialva 8, 7000-809 Évora, Portugal.

³ Science and Technology School of Évora University, Rua Romão Ramalho nº59, 7000-671 Évora, Portugal.

* Correspondence: Correspondence: fariajms@gmail.com

† Presented at the 1st International Electronic Conference on Forests, 15–30 November 2020;

Available online: <https://sciforum.net/conference/IECF2020>

Published: 25 October 2020

Abstract: The pinewood nematode (PWN), *Bursaphelenchus xylophilus*, induces shoot yellowing and wilting by injuring pine vascular tissues and resin canals. In Asia, it devastated wide forests of susceptible pine, causing drastic ecological, economic and cultural repercussions. In 1999, it reached Europe (Portugal) and despite the actions of the authorities, it rapidly advanced to the border areas of Spain, threatening the European pine forests. Chemical control is used in Asia with remarkable success, yet most nematocides are dangerous to human health and the environment. Natural volatile allelochemicals (VA) and derivatives are sustainable alternatives, providing many advantages to commercial nematocides. The present work summarizes available bibliographic information on VAs with activity against the PWN and reviews the chemical properties leading to nematotoxicity. Published works reported over 250 VAs in direct contact assays. Compounds highly active against the PWN belong to monoterpenoid, phenylpropanoid and aliphatic alcohols, aldehydes, ketones, carboxylic acids, esters and sulphide groups. The presence of highly electronegative elements in these structures appears to increase activity. Trisulphides, coumarins, medium carbon chain length aliphatic alcohols and derivatives composed the top 5 most active allelochemicals. An indication of their mechanisms of action can be obtained from studying the reported structure-activity relationships.

Keywords: *Bursaphelenchus xylophilus*; monoterpenoids; nematocide; oxygenated aliphatic compounds; phenylpropanoids; pinewood nematode; sulphides; toxicity

1. Introduction

Pine wilt disease (PWD) induces pine shoot yellowing (chlorosis) and wilting (caused by embolism), killing a mature tree in less than 50 days. It is caused by the pinewood nematode (PWN), *Bursaphelenchus xylophilus* (Steiner & Buhner 1934), an invasive microscopic plant parasitic nematode with a short life cycle and an impressive spread rate. It was introduced from North America to Japan around 1905, devastating vast pine forests. The PWN has since spread to neighboring China (1982) and Korea (1988). Susceptible Asian pine forests were heavily affected causing irreversible changes to native forest ecosystems and significant losses to national economies [1]. In 1999, Portugal became its entry point into Europe, most likely through imported goods from Asian countries [2]. It has since spread throughout Portugal and reached Spanish border cities, after only 10 years [3,4]. Due to the environmental conditions imposed by climate change, there is a very high risk for a European wide epidemic of this forest pathology [5]. The PWN is now classified as an A2 quarantine pest and the European Union (EU) has taken actions to ensure its control beyond the present geographic area and

to eradicate it from the EU territory. PWN containment strategies generally rely on removal of the insect vector through aerial application of synthetic insecticides, establishment of pine tree-free buffer zones, fumigation of infected trees or on the use of insect vector natural enemies [1]. In the most affected countries, chemical control is considered one of the most effective and reliable containment strategies within integrated management. Directly killing the nematode is performed by trunk injection of synthetic/hemisynthetic nematicides, like avermectins, carbamates, neonicotinoids, etc [6]. Unfortunately, in general, common use nematicides are toxic to other beneficial microorganisms, to humans and animals, and can accumulate in the soil and in food plants. Development of environmentally safer strategies to control invasive PWN populations has become paramount, due to the ban imposed on hazardous pesticides and the recent fear of drug resistance on the PWN [6]. Strong pressures on the development of improved ecological biopesticides has prompted researchers to screen for environmentally friendlier natural compounds with increased anti-PWN properties which are, at the same time, cost-effective. Primary breakthroughs have been reached by screening highly active volatile allelochemicals (VAs), commonly produced by plants and microbes [7,8]. These compounds have the advantage of not accumulating in the environment and having a broad range of activities, which diminishes the risk of developing resistant pathogenic strains. Mixtures of VAs, such as essential oils, have shown great promise due to their high nematicidal activities and for being easily obtained [9]. Research has been mainly performed in the most affected countries, Japan, China, Korea and recently in Portugal. Faria et al. [10] screened over 80 EOs and identified their chemical composition through gas chromatography coupled to mass spectrometry (GC-MS). Through EO fractionation, the authors were able to attribute anti-PWN activity to the oxygen containing EO components. Other authors have also identified EOs chemical composition and tested the main components of the active EOs by using commercially acquired chemicals, achieving noteworthy results [11,12].

In the present review, a comprehensive bibliographic research was performed to catalogue the tested VAs and identify groups with the highest activities against the PWN. A critical overview of their chemical structures allows highlighting the most successful anti-PWN chemical properties, in an effort to uncover novel highly nematicidal environmentally friendly volatiles and develop eco-friendly anti-PWN strategies.

2. Bibliographic data

Research was performed using Web of Science® search engine on published works in all available databases, using the topics “*Bursaphelenchus xylophilus*” and “nematicide” or “nematocide” or “nematodicide”. To ensure a direct causality between allelochemicals contact and PWN mortality, only works reporting direct contact bioassays were selected. Works were identified, catalogued and information on compound chemical nature and half maximal effective concentration (EC₅₀) against the PWN was collected when available. A total of 24 publications were identified reporting the activity of anti-PWN VAs and derivatives. The first identified report dated to 1993 and the latest to 2020 (with 2 reports by 10 September). The highest number of reports was published during 2007 (5 publications). Regarding the scientific fields covered by the journals that published these reports, 33% were in zoology, 17% in biochemistry and molecular biology and 17 % in applied chemistry (Figure 1a). The identified publications were cited 650 times (611 when excluding self-citations) by a total of 478 articles (462, excluding self-citations), with an average of 27 citations per work. From 1995 to 2005 this field showed a stable low interest, with an average 3 citations/year, yet since 2005, the study of VAs and derivatives as anti-PWN agents gained momentum and showed a linear growth, triplicating citations each year (slope=3) (Figure 1b). The pattern of increased citation since mid-2000s may be due to a similar increase in reports on the nematicidal activity of essential oils (EO) [13]. These complex mixtures of volatiles are mainly comprised of mono- and sesquiterpenes (and a few diterpenes), phenylpropanoids and other groups of volatile compounds, in relevant amounts. Their biological activity can generally result from the combined effect (synergistic or antagonistic) of compounds with direct activity and those with no direct activity towards the biological system, that can influence resorption, rate of reactions and bioavailability of the active compounds [10]. In the 24

publications identified, EOs were tested and profiled through gas chromatography coupled to mass spectrometry (GC-MS), to identify the component(s) responsible for the nematocidal activity against the PWN.




Figure 1. (a) Percentage of reports on the activity of volatile allelochemicals and derivatives against *Bursaphelenchus xylophilus* distributed by scientific journal areas and (b) yearly citation evolution of these reports.

3. Anti-PWN allelochemicals

Worldwide research efforts tested the activity of over 250 allelochemicals against the PWN. Publications report mainly on compounds identified in plant extracts [14–18] or essential oils [11,12,19–23], microbial extracts [24–26] and compounds synthesized based on highly active analogs [27–35]. The compounds assayed mainly belonged to highly active classes of chemical compounds, namely aliphatic alcohols, aldehydes, ketones, carboxylic acids, esters and sulphides (42%), phenylpropanoids (27%) and monoterpenoids (18%) (Figure 2a). Allelochemicals with direct activity against the PWN showed highly electronegative elements in their composition. The most active were oxygenated compounds, with alcohol, aldehyde or carboxylic acid active groups, or with sulphur, with disulfide or trisulphide functional groups.




Figure 2. (a) Percentage of the main chemical groups of the volatile allelochemicals and derivatives reported against the PWN; “less than 10” includes classes with less than 10 instances. (b) The top 3 compounds with the lowest reported half maximal effective concentrations (EC₅₀).

The top 3 compounds with the lowest EC_{50} values were trans-2-hexenal, undecanol and 2-(1-undecyloxy)-1-ethanol (Figure 2b). Trans-2-hexenal is an aliphatic aldehyde with a C6 length carbon chain, very common in fresh green plant EOs [10,17,36,37]. It's used as a flavoring compound in food to induce a leafy fruity note and in perfumes to obtain a green-leaf note. Against the PWN it appears to influence several biologic functions, such as reproduction, reducing rate of reproduction and number of eggs, and development, such as body length, respiration, movement, nutrition and enzymatic activity [17,18]. Undecanol is an aliphatic primary alcohol with a C11 length carbon chain. It's a colorless or pale-yellow lemon-scented liquid used for some typical aroma fragrance only at low concentrations. Its activity against the PWN appears to be a part of a mechanism of toxicity imposed by oxygenated aliphatic compounds. Aliphatic compounds with C8 to C11 carbon chain lengths containing alcohol, aldehyde or carboxylic acid functional groups showed increased activities against the PWN in comparison to lower or higher chain lengths or to hydrocarbons or alkanolic acetates. In this mechanism the presence of double bonds next to the functional group appears to play a role in anti-PWN activity [28]. 2-(1-undecyloxy)-1-ethanol shows similarities to undecanol by having a C11 carbon chain linked to an oxygenated functional group. Interestingly, this molecule is an aggregation pheromone that attracts both conspecific females and males of *Monochamus* species, the PWN insect vector in many countries. This pheromone, known also as monochamol, is commonly used in insect traps against the PWD [31].

Chemical characteristics of successful anti-PWN allelochemicals

The information gathered allowed crossing data on the most nematotoxic compounds and their chemical nature. Some preliminary insights into the chemical basis of anti-PWN efficacy were reported in the publications analyzed [11,23,28,29,31]. Concerning terpenoids, fundamentally monoterpenoids; compounds with phenol, alcohol or aldehyde functional groups had higher anti-PWN activities than ketones or hydrocarbons. Within compounds with the alcohol functional group, primary alcohols showed greater activity than secondary or tertiary alcohols. Molecule isomerism also appears to contribute to nematocidal activity. The position of the substituent in the geometrical isomer of monoterpenoid molecules influences anti-PWN activity. Regarding phenylpropanoids, besides most of the characteristics reported for monoterpenoids, aldehydes, carboxylic acids and alcohols are reported to be more active against the PWN than their corresponding saturated compounds. In the carboxylic acids of these compounds, allyl, ethyl and methyl substituents in the aromatic ring showed higher activities than their isopropyl and vinyl compounds. Phenylpropanoids with hydroxyl or methoxy groups showed higher activities than those with an acetyl group. The reported aliphatic alcohols, aldehydes, ketones, carboxylic acids, esters and sulphides showed also interesting structure/anti-PWN activity relationships. Compounds with alcohol, aldehyde and carboxylic acid functional groups were more active against the PWN than aliphatic hydrocarbons or acetates. The enhancement of anti-PWN activity in aldehydes is linked to unsaturated α,β -carbonyl compounds. More interestingly, aliphatic alcohols with specific shape and size showed the most successful activities against the PWN, C9 to C11 aliphatic chain alkanols and trans-2-alkenols showed substantially higher activities than those with lower or higher chain lengths. Finally, sulphides were strongly nematotoxic, showing a positive correlation between anti-PWN activity and sulphur number.

4. Conclusion

Research for highly active volatile allelochemicals against the PWN is relatively recent, less than 30 years, but has gained momentum in the last 10 years, promoted by an increasing search for alternative sustainable strategies to contain this highly damaging forest pest. The works reported so far highlight the success of compounds with oxygenated functional groups. Monoterpenoids, phenylpropanoids and aliphatic oxygenated compounds show the most promising activities. Future research must take advantage of this knowledge to uncover the specific sites and mechanisms of activity employed by these compounds against the PWN. This information will contribute for the development of effective sustainable strategies to control PWD spread.

Author Contributions: conceptualization, J.F.; methodology, J.F.; investigation, J.F. and P.B.; resources, J.F.; writing—original draft preparation, J.F.; writing—review and editing, J.F., P.B., D.T. and M.M.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

1. References

2. Vicente, C.; Espada, M.; Vieira, P.; Mota, M. Pine Wilt Disease: A threat to European forestry. *Eur. J. Plant Pathol.* **2012**, *133*, 89–99, doi:10.1007/s10658-011-9924-x.
3. Mota, M.; Braasch, H.; Bravo, M.A.; Penas, A.C.; Burgermeister, W.; Metge, K.; Sousa, E. First report of *Bursaphelenchus xylophilus* in Portugal and in Europe. *Nematology* **1999**, *1*, 727–734, doi:https://doi.org/10.1163/156854199508757.
4. Robertson, L.; Cobacho Arcos, S.; Escuer, M.; Santiago Merino, R.; Esparrago, G.; Abelleira, A.; Navas, A. Incidence of the pinewood nematode *Bursaphelenchus xylophilus* Steiner & Buhner, 1934 (Nickle, 1970) in Spain. *Nematology* **2011**, *13*, 755–757, doi:10.1163/138855411 × 578888.
5. Abelleira, A.; Picoaga, A.; Mansilla, J.P.; Aguin, O. Detection of *Bursaphelenchus xylophilus*, causal agent of Pine Wilt Disease on *Pinus pinaster* in Northwestern Spain. *Plant Dis.* **2011**, *95*, 776, doi:10.1094/PDIS-12-10-0902.
6. Ikegami, M.; Jenkins, T.A.R. Estimate global risks of a forest disease under current and future climates using species distribution model and simple thermal model – Pine Wilt disease as a model case. *For. Ecol. Manage.* **2018**, *409*, 343–352, doi:10.1016/j.foreco.2017.11.005.
7. Bi, Z.; Gong, Y.; Huang, X.; Yu, H.; Bai, L.; Hu, J. Efficacy of four nematicides against the reproduction and development of pinewood Nematode, *Bursaphelenchus xylophilus*. *J. Nematol.* **2015**, *47*, 126–132.
8. Kong, J.O.; Lee, S.M.; Moon, Y.S.; Lee, S.G.; Ahn, Y.J. Nematicidal activity of plant essential oils against *Bursaphelenchus xylophilus* (Nematoda: Aphelenchoididae). *J. Asia. Pac. Entomol.* **2006**, *9*, 173–178, doi:10.1016/S1226-8615(08)60289-7.
9. Yang, Z.; Yu, Z.; Lei, L.; Xia, Z.; Shao, L.; Zhang, K.; Li, G. Nematicidal effect of volatiles produced by *Trichoderma* sp. *J. Asia. Pac. Entomol.* **2012**, *15*, 647–650, doi:10.1016/j.aspen.2012.08.002.
10. Choi, I.H.; Park, J.Y.; Shin, S.C.; Kim, J.; Park, I.K. Nematicidal activity of medicinal plant essential oils against the pinewood nematode (*Bursaphelenchus xylophilus*). *Appl. Entomol. Zool.* **2007**, *42*, 397–401, doi:10.1303/aez.2007.397.
11. Faria, J.M.S.; Barbosa, P.; Bennett, R.N.; Mota, M.; Figueiredo, A.C. Bioactivity against *Bursaphelenchus xylophilus*: Nematotoxics from essential oils, essential oils fractions and decoction waters. *Phytochemistry* **2013**, *94*, 220–228, doi:10.1016/j.phytochem.2013.06.005.
12. Choi, I.H.; Kim, J.; Shin, S.C.; Park, I.K. Nematicidal activity of monoterpenoids against the pine wood nematode (*Bursaphelenchus xylophilus*). *Russ. J. Nematol.* **2007**, *15*, 35–40.
13. Kong, J.O.; Park, I.K.; Choi, K. sik; Shin, S.C.; Ahn, Y. joon Nematicidal and propagation activities of thyme red and white oil compounds toward *Bursaphelenchus xylophilus* (Nematoda: Parasitaphelenchidae). *J. Nematol.* **2007**, *39*, 237–242.
14. Andrés, M.F.; González-Coloma, A.; Sanz, J.; Burillo, J.; Sainz, P. Nematicidal activity of essential oils: A review. *Phytochem. Rev.* **2012**, *11*, 371–390, doi:10.1007/s11101-012-9263-3.
15. Suga, T.; Ohta, S.; Munesada, K.; Ide, N.; Kurokawa, M.; Shimizu, M.; Ohta, E. Endogenous pine wood nematicidal substances in pines, *Pinus massoniana*, *P. strobus* and *P. palustris*. *Phytochemistry* **1993**, *33*, 1395–1401, doi:10.1016/0031-9422(93)85098-C.

16. Alen, Y.; Nakajima, S.; Nitoda, T.; Baba, N.; Kanzaki, H.; Kawazu, K. Two Antinematodal Phenolics from *Knema hookeriana*, a Sumatran Rainforest Plant. *Zeitschrift für Naturforsch. C* **55**, 300–304, doi:https://doi.org/10.1515/znc-2000-3-426.
17. Hanawa, F.; Yamada, T.; Nakashima, T. Phytoalexins from *Pinus strobus* bark infected with pinewood nematode, *Bursaphelenchus xylophilus*. *Phytochemistry* **2001**, *57*, 223–228, doi:10.1016/S0031-9422(00)00514-8.
18. Cheng, L.; Xu, S.; Xu, C.; Lu, H.; Zhang, Z.; Zhang, D.; Mu, W.; Liu, F. Effects of trans-2-hexenal on reproduction, growth and behaviour and efficacy against the pinewood nematode, *Bursaphelenchus xylophilus*. *Pest Manag. Sci.* **2017**, *73*, 888–895, doi:10.1002/ps.4360.
19. Zhao, Y.; Xu, S.; Lu, H.; Zhang, D.; Liu, F.; Lin, J.; Zhou, C.; Mu, W. Effects of the plant volatile trans-2-hexenal on the dispersal ability, nutrient metabolism and enzymatic activities of *Bursaphelenchus xylophilus*. *Pestic. Biochem. Physiol.* **2017**, *143*, 147–153, doi:10.1016/j.pestbp.2017.08.004.
20. Park, I.K.; Park, J.Y.; Kim, K.H.; Choi, K.S.; Choi, I.H.; Kim, C.S.; Shin, S.C. Nematicidal activity of plant essential oils and components from garlic (*Allium sativum*) and cinnamon (*Cinnamomum verum*) oils against the pine wood nematode (*Bursaphelenchus xylophilus*). *Nematology* **2005**, *7*, 767–774, doi:10.1163/156854105775142946.
21. Park, J.-Y.; Choi, I.-H.; Park, I.-K.; Shin, S.-C. Nematicidal activity of medicinal plant extracts and two cinnamates isolated from *Kaempferia galanga* L. (Proh Hom) against the pine wood nematode, *Bursaphelenchus xylophilus*. *Nematology* **2006**, *8*, 359–365, doi:10.1163/156854106778493402.
22. Park, I.-K.; Kim, J.; Lee, S.-G.; Shin, S.-C. Nematicidal activity of plant essential oils and components from Ajowan (*Trachyspermum ammi*), Allspice (*Pimenta dioica*) and Litsea (*Litsea cubeba*) essential oils against pine wood nematode (*Bursaphelenchus Xylophilus*). *J. Nematol.* **2007**, *39*, 275–9.
23. Choi, I.-H.; Shin, S.-C.; Park, I.-K. Nematicidal activity of onion (*Allium cepa*) oil and its components against the pine wood nematode (*Bursaphelenchus xylophilus*). *Nematology* **2007**, *9*, 231–235, doi:10.1163/156854107780739018.
24. JunHeon, K.; SunMi, S.; SangGil, L.; SangChul, S.; IlKwon, P. Nematicidal activity of plant essential oils and components from coriander (*Coriandrum sativum*), Oriental sweetgum (*Liquidambar orientalis*), and valerian (*Valeriana wallichii*) essential oils against pine wood nematode (*Bursaphelenchus xylophilus*). *J. Agric. Food Chem.* **2008**, *56*, 7316–7320, doi:10.1021/jf800780f.
25. Yu, J.; Du, G.; Li, R.; Li, L.; Li, Z.; Zhou, C.; Chen, C.; Guo, D. Nematicidal activities of bacterial volatiles and components from two marine bacteria. *Nematology* **2015**, *17*, 1011–1025, doi:10.1163/15685411-00002920.
26. Bang, H.B.; Lee, Y.H.; Kim, S.C.; Sung, C.K.; Jeong, K.J. Metabolic engineering of *Escherichia coli* for the production of cinnamaldehyde. *Microb. Cell Fact.* **2016**, *15*, 16, doi:10.1186/s12934-016-0415-9.
27. Huang, Z.; Dan, Y.; Huang, Y.; Lin, L.; Li, T.; Ye, W.; Wei, X. Sesquiterpenes from the mycelial cultures of *Dichomitus squalens*. *J. Nat. Prod.* **2004**, *67*, 2121–2123, doi:10.1021/np0497144.
28. Kohno, T.; Togashi, K.; Fukamiya, N. The nematocidal activity and the structure-activity relationships of stilbenes. *Nat. Prod. Res.* **2007**, *21*, 606–615, doi:10.1080/14786410701369730.
29. Seo, S.M.I.; Junheon, K.; Eunae, K.; Park, H.M.I.; Kim, Y.J.; Park, I.L.K. Structure-Activity relationship of aliphatic compounds for nematicidal activity against pine wood nematode (*Bursaphelenchus xylophilus*). *J. Agric. Food Chem.* **2010**, *58*, 1823–1827, doi:10.1021/jf902575f.

30. Seo, S.M.; Kim, J.; Koh, S.H.; Ahn, Y.J.; Park, I.K. Nematicidal activity of natural ester compounds and their analogues against pine wood nematode, *bursaphelenchus xylophilus*. *J. Agric. Food Chem.* **2014**, *62*, 9103–9108, doi:10.1021/jf503631e.
31. Pan, L.; Li, X.Z.; Sun, D.A.; Jin, H.; Guo, H.R.; Qin, B. Design and synthesis of novel coumarin analogs and their nematicidal activity against five phytonematodes. *Chinese Chem. Lett.* **2016**, *27*, 375–379, doi:10.1016/j.ccl.2016.01.029.
32. Kim, J.; Lee, S.M.; Park, C.G. *Bursaphelenchus xylophilus* is killed by homologues of 2-(1-undecyloxy)-1-ethanol. *Sci. Rep.* **2016**, *6*, 2–6, doi:10.1038/srep29300.
33. Lee, Y.J.; Bashyal, P.; Pandey, R.P.; Sohng, J.K. Enzymatic and microbial biosynthesis of novel violacein glycosides with enhanced water solubility and improved anti-nematode activity. *Biotechnol. Bioprocess Eng.* **2019**, *24*, 366–374, doi:10.1007/s12257-018-0466-3.
34. Li, Y.; Feng, Y.; Wang, X.; Cui, J.; Deng, X.; Zhang, X. Adaptation of pine wood nematode *Bursaphelenchus xylophilus* to β -pinene stress. *BMC Genomics* **2020**, *21*, 1–16, doi:10.1186/s12864-020-06876-5.
35. Cheng, F.; Wang, J.; Song, Z.; Cheng, J.; Zhang, D.; Liu, Y. Nematicidal effects of 5-aminolevulinic acid on plant-parasitic nematodes. *J. Nematol.* **2017**, *49*, 295–303.
36. Cui, J.; Li, Y.-X.; Zhang, W.; Wang, X.; Pan, L.; Feng, Y.-Q.; Zhang, X.-Y. The population structure and sex ratios of *Bursaphelenchus xylophilus* under α -pinene stress. *J. For. Res.* **2020**, *31*, 921–926, doi:10.1007/s11676-018-0847-7.
37. Barbosa, P.; Faria, J.M.S.; Mendes, M.D.; Dias, L.S.; Tinoco, M.T.; Barroso, J.G.; Pedro, L.G.; Figueiredo, A.C.; Mota, M. Bioassays against pinewood nematode: Assessment of a suitable dilution agent and screening for bioactive essential oils. *Molecules* **2012**, *17*, doi:10.3390/molecules171012312.
38. Zhao, Y.; Xu, S.; Lu, H.; Zhang, D.; Liu, F.; Lin, J.; Zhou, C.; Mu, W. Effects of the plant volatile trans-2-hexenal on the dispersal ability, nutrient metabolism and enzymatic activities of *Bursaphelenchus xylophilus*. *Pestic. Biochem. Physiol.* **2017**, *143*, 147–153, doi:10.1016/j.pestbp.2017.08.004.

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).