



OPEN ACCESS

EDITED AND REVIEWED BY
Jeroen Van Hunen,
Durham University, United Kingdom

*CORRESPONDENCE
João Fontiela,
✉ jfontiela@uevora.pt

RECEIVED 11 May 2024
ACCEPTED 20 May 2024
PUBLISHED 03 June 2024

CITATION

Fontiela J, Silveira G, Dias NA, Pimentel A, Ramalho RS and Sigloch K (2024), Editorial: Challenges in seismology to our understanding of volcanic islands and their magmatic unrest.
Front. Earth Sci. 12:1431193.
doi: 10.3389/feart.2024.1431193

COPYRIGHT

© 2024 Fontiela, Silveira, Dias, Pimentel, Ramalho and Sigloch. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Editorial: Challenges in seismology to our understanding of volcanic islands and their magmatic unrest

João Fontiela^{1*}, Graça Silveira^{2,3}, Nuno Afonso Dias^{2,3}, Adriano Pimentel⁴, Ricardo S. Ramalho^{2,5,6} and Karin Sigloch⁷

¹Institute of Earth Sciences, University of Évora, Évora, Portugal, ²Instituto Dom Luiz, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal, ³Instituto Superior de Engenharia de Lisboa, Instituto Politécnico de Lisboa, Lisboa, Portugal, ⁴Instituto de Investigação em Vulcanologia e Avaliação de Riscos (IVAR), Universidade dos Açores, Ponta Delgada, Portugal, ⁵School of Earth and Environmental Sciences, Cardiff University, Park Place, Cardiff, United Kingdom, ⁶Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY, United States, ⁷Université Côte d'Azur, CNRS, Observatoire Côte d'Azur, IRD, Géoazur, Sophia Antipolis Cedex, Nice, France

KEYWORDS

volcanic islands, geodynamics, seismic anisotropy, seismic interferometry, seismic tomography, volcano monitoring

Editorial on the Research Topic

Challenges in seismology to our understanding of volcanic islands and their magmatic unrest

Volcanic islands are some of the most intriguing geological features on Earth, yet the processes that lead to their formation and evolution remain largely enigmatic. Such islands exhibit complex, heterogeneous and anisotropic subsurface structures, characterized by pronounced and abrupt changes in seismic velocities, attenuation, anisotropy, and interfaces. Volcanic islands are amongst the planet's largest and fastest-forming geological features. They are sites of significant geohazards including volcanic eruptions, earthquakes, landslides, and tsunamis. Seismic monitoring is therefore crucial for detecting magmatic movements and issuing pre-eruptive warnings, as well as for understanding the internal structure of islands, thereby aiding in hazard and risk mitigation.

Studying volcanic islands poses challenges due to the complex processes occurring at depth and the logistical constraints imposed by the islands' size, shape and/or location relative to other islands. To overcome this, there is a critical need to densify seismic networks with land-based and ocean-bottom seismometers to enhance geographical coverage. This will enable better imaging of the subsurface and detection and classification of seismic signals associated with volcanic and tectonic activity. Generating 3D models from the shallow crust to the deep mantle, combining multiple seismic and other geophysical techniques is essential for understanding the geodynamic processes driving magma ascent and volcanism.

Our Research Topic, "Challenges in seismology to our understanding of volcanic islands and their magmatic unrest" features four original research articles. These articles explore dynamic subsurface changes and fluid migration during eruptions, innovate volcano-monitoring methodologies, and examine the complex mantle structures beneath

volcanic islands, enhancing our understanding of the processes occurring before and during eruptions, and geodynamic constraints.

Civiero et al. conducted a comprehensive review of studies focusing on the complex mantle structure and transition zone beneath the Macaronesian volcanic islands in the central-eastern Atlantic (comprising the Azores, Madeira, Canaries, and Cape Verde). Their analysis underscored regional models which illustrate low-velocity features in the asthenosphere and a notably thinned transition zone, suggesting that these characteristics are due to mantle plumes. Additionally, these tomographic models were found to be consistent with global mantle tomography models. The authors also explored alternative theories that attribute volcanism to mechanisms related to mantle upwellings and plate tectonics, offering a broader perspective on the underlying geodynamic processes.

Continuous monitoring of active volcanoes for early warning and eruption forecasting continues to pose significant challenges. Addressing this, Fenner et al. utilized the Adaptive-Window Volcanic Event Selection Analysis Module (AWESAM) on several island volcanoes, including Stromboli and Etna in Italy, Yasur in Vanuatu, and Whakaari in New Zealand. Their study aimed to detect characteristic patterns in the recurrence times and overall volcanic activity. The findings revealed that, despite geographical differences, similar traits were consistently observed across all studied volcanoes in terms of amplitude-frequency relations, inter-event time distributions, and amplitude distributions. This consistency underscores the effectiveness of AWESAM in providing reliable data for volcanic monitoring.

Significant changes in the structure of volcanoes, driven by magma movement, are crucial to the forecasting of future eruptions. Shear-Wave Splitting (SWS) is an effective technique for assessing crustal and mantle anisotropy. Schlaphorst et al. used crustal SWS in an attempt to detect changes in the anisotropy pattern linked to volcanic unrest. The authors document the temporal variations in seismic anisotropy before, during, and after the 2011–2012 eruption offshore El Hierro and the 2021 eruption in La Palma in the Canary Islands. The two years following the El Hierro eruption show clear evidence of intrusive events. Following a period of relative calm, an uptick in results from El Hierro since 2018 suggests renewed subsurface activity in deeper parts of the system, impacting overall seismicity rates but not vertical GPS measurements concurrently. In La Palma, subsurface dynamic changes were observed in the week leading up to the 2021 eruption.

Ambient seismic noise interferometry is a cost-effective and high-sensitivity tool to detect and analyze changes in velocity related to subsurface changes over time and space. In their study, Carvalho et al. applied auto- and cross-correlation functions of ambient seismic noise to assess temporal structural changes induced by magma movement in La Palma between 2020 and 2022. Their findings indicate a significant decorrelation prior to the eruption, which is primarily attributed to the ascent of hydrothermal fluids. A marked increase in decorrelation was also noted at the onset of the eruption. Post-eruption observations revealed that, while some deeper parts of the plumbing system remained pressurized by hydrothermal fluids, the shallow structures experienced minimal changes and quickly returned to the pre-eruptive state.

This Research Topic underscores the ongoing efforts and advances in understanding and monitoring volcanic islands, demonstrating the critical role of integrated seismological studies in managing volcanic hazards and associated risks, and improving forecasting capabilities.

Author contributions

JF: Writing–original draft, Writing–review and editing, Validation. GS: Writing–original draft, Writing–review and editing. ND: Writing–review and editing. AP: Writing–review and editing. RR: Writing–review and editing. KS: Writing–review and editing.

Funding

The author(s) declare that financial support was received for the research, authorship, and/or publication of this article. This work is a contribution to the three projects funded by FCT I.P./ MCTES through national funds, SIGHT PTDC/CTA-GEF/30264/2017, RESTLESS PTDC/CTA-GEF/ 6674/2020 (<https://doi.org/10.54499/PTDC/CTA-GEF/6674/2020>) and GEMMA PTDC/CTA-GEO/2083/2021 (<https://doi.org/10.54499/PTDC/CTA-GEO/2083/2021>). This work was also funded by the Portuguese Fundação para a Ciência e a Tecnologia (FCT) I.P./MCTES through national funds PIDDAC-UIDB/04683/2020 (<https://doi.org/10.54499/UIDB/04683/2020>), UIDP/04683/2020 (<https://doi.org/10.54499/UIDP/04683/2020>), UIDB/50019/2020 (<https://doi.org/10.54499/UIDB/50019/2020>), UIDP/50019/2020 (<https://doi.org/10.54499/UIDP/50019/2020>), and LA/P/0068/2020 (<https://doi.org/10.54499/LA/P/0068/2020>). AP is supported by project <https://doi.org/10.54499/CEECINST/00024/2021/CP2780/CT0003>. GS, ND, and JF are grateful for support from European Research Council (ERC) UPFLOW consolidator grant under the European Union's Horizon 2020 research and innovation programme (grant agreement No 10100160).

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.